

BIBLIOGRAPHY

1. S. A. Amitsur, *The identities of PI-rings*, Proc. Amer. Math. Soc. vol. 4 (1953) pp. 27-34.
2. ———, *Algebras over infinite fields*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 35-48.
3. I. Kaplansky, *Rings with a polynomial identity*, Bull. Amer. Math. Soc. vol. 54 (1948) pp. 575-580.
4. W. Krull, *Jacobson'sches Radical und Hilbertscher Nullstellensatz*, International Congress of Mathematicians, Cambridge, 1950, Conference in Algebra, pp. 56-64.

HEBREW UNIVERSITY, JERUSALEM

A NOTE ON THE STONE-WEIERSTRASS THEOREM FOR QUATERNIONS¹

JOHN C. HOLLADAY

A result of M. H. Stone [1, p. 466], which is nicely presented by N. Dunford [2, p. 23], is as follows: Let A be a closed subalgebra of the B -algebra $C(X)$ of all continuous real-valued functions on the compact Hausdorff space X . Then $A = C(X)$ if and only if A distinguishes between every pair of distinct points of X , i.e., for every pair $x_1 \neq x_2$ of points in X , there is an f in A such that $f(x_1) \neq f(x_2)$.

If one substitutes the word complex for the word real in the above statement, it becomes false. A well known counter example is obtained by letting X be the set of complex numbers z such that $|z| \leq 1$ and letting A be the subalgebra of functions which are analytic in the interior of X .

The purpose of this note is to show that if the word quaternion is substituted for the word real in the above statement, it remains valid. To be specific, let A be a set of continuous quaternion-valued functions which satisfy the following conditions:

1. A is complete.
2. Given a quaternion q , the function $f(x) \equiv q$ is in A .
3. If f and g are in A , then fg and $f+g$ are in A .

Received by the editors November 18, 1956.

¹ A small part of the work done under an AEC Predoctoral Fellowship at Yale University, year 1952-1953, under the kind and patient guidance of Dr. Charles E. Rickart.

If A contains all continuous quaternion-valued functions, it obviously distinguishes between points. Letting A distinguish between points, consider two arbitrary distinct points x_1 and x_2 . Choose an element of A which takes a different value at x_2 than at x_1 . Multiply this function by an appropriate quaternion to obtain a function f such that real part $[f(x_1)] \neq \text{real part } [f(x_2)]$. But the real part of f is $[f - ifi - jfj - kfk]/4$ which is an element of A . Therefore, A contains real valued functions which distinguish between points.

Since A is complete, and is closed under multiplication, addition and subtraction, it follows that the set of all real-valued functions in A is also complete and closed under these arithmetic operations. The Stone-Weierstrass Theorem implies that A contains all continuous real-valued functions on X . Therefore, A contains all continuous quaternion-valued functions on X .

BIBLIOGRAPHY

1. M. H. Stone, *Applications of the theory of Boolean rings to general topology*, Trans. Amer. Math. Soc. vol. 41 (1937) pp. 375-481.
2. N. Dunford, *Spectral theory in abstract spaces and Banach algebras*, Symposium on Spectral Theory and Differential Equations, pp. 1-26.

LOS ALAMOS SCIENTIFIC LABORATORY AND
YALE UNIVERSITY