

THE RING $C(X, R)$ CONSIDERED AS A SUBRING OF THE RING OF ALL REAL-VALUED FUNCTIONS

LYLE E. PURSELL

Conditions under which the ring $C(X, R)$ of all real-valued continuous functions on a completely regular space X determines X have been investigated in a number of papers [1-5]. Gelfand and Kolmogoroff [2] have shown that if X is a compact completely regular space, then the space of maximal ideals in $C(X, R)$ with the Stone topology is homeomorphic to X and hence $C(X, R)$ determines X . The most general result for noncompact spaces has been obtained by F. W. Anderson [1] who has shown that the lattice of all real-valued continuous functions on a topological space determines this space if it is a completely regular space all of whose points are G -delta sets. Since $f(x) \geq g(x)$ for all x if and only if $f - g$ has square root in $C(X, R)$, a ring isomorphism of $C(X, R)$ implies a lattice isomorphism and Anderson's result is equivalent to the statement that the ring $C(X, R)$ determines X if X is a completely regular space all of whose points are G -delta sets.

In this paper we show that if X is completely regular, then X is topologically determined by the pair of rings consisting of the ring R^X of all real-valued functions on X and its subring $C(X, R)$.

LEMMA 1. *An ideal I in $C(X, R)$ is fixed if and only if there exists a function g in R^X such that $0 \leq g(x) \leq 1$ for all x in X , $g(x_0) = 0$ for some x_0 and if f is in I and $-1 \leq f(x) \leq 1$ for all x , then $-g(x) \leq f(x) \leq g(x)$ for all x .*

PROOF. If all functions in I vanish at some point x_0 , then we observe that if we define $g(x) = 1$ for $x \neq x_0$ and $g(x_0) = 0$, then g is the desired function.

Conversely, assume that for an ideal I such a function g exists. If f is continuous, then the function h defined by $h(x) = f(x)$ if $-1 \leq f(x) \leq 1$ and $h(x) = 1/f(x)$ otherwise is continuous. If f is in I , then fh is in I , $-1 \leq (fh)(x) \leq 1$ for all x , and $(fh)(x) = 0$ if and only if $f(x) = 0$. Hence $-g(x) \leq (fh)(x) \leq g(x)$ for all x , $(fh)(x_0) = 0$, and, accordingly, $f(x_0) = 0$. Therefore I is a fixed ideal.

Since every automorphism of R^X preserves the relation $f(x) \leq g(x)$

Presented to the Society, November 23, 1956 under the title *The ring of all real-valued continuous functions considered as a subring of the ring of all real-valued functions*; received by the editors November 13, 1956.

for all x , then every automorphism of R^X which maps $C(X, R)$ onto itself will map a fixed ideal of $C(X, R)$ onto a fixed ideal of $C(X, R)$.

THEOREM 1. *If X is a completely regular topological space, then the space of fixed maximal ideals with the Stone topology (i.e. a collection of fixed maximal ideals is closed if and only if their intersection J is equal to the intersection of all fixed maximal ideals which contain J) is homeomorphic to X .*

The proof is similar to the proof [2] that the set of maximal ideals in $C(X, R)$ with the Stone topology is homeomorphic to X if X is compact. One observes that if $C(X, R)$ is a separating class for X (i.e. if $x_1 \neq x_2$, then there is a function f in $C(X, R)$ such that $f(x_1) \neq f(x_2)$) but X is not completely regular, then there is a one-to-one correspondence between X and the space of fixed maximal ideals but the latter space will have fewer closed sets.

From the above results we obtain:

THEOREM 2. *If X and Y are two completely regular spaces and if R^X and R^Y are isomorphic under an isomorphism which maps $C(X, R)$ onto $C(Y, R)$, then X and Y are homeomorphic.*

Lemma 1 and Theorems 1 and 2 also hold if $C(X, R)$ is replaced by the ring $C^*(X, R)$ of all bounded continuous functions on X .

BIBLIOGRAPHY

1. F. W. Anderson, *A lattice characterization of completely regular G -delta spaces*, Proc. Amer. Math. Soc. vol. 6 (1955) pp. 757-765.
2. I. Gelfand and A. N. Kolmogoroff, *On rings of continuous functions on topological spaces*, C. R. (Doklady) Acad. Sci. URSS. vol. 22 (1939) pp. 11-15.
3. L. Gillman, M. Henriksen and M. Jerison, *On a theorem of Gelfand and Kolmogoroff concerning maximal ideals in rings of continuous functions*, Proc. Amer. Math. Soc. vol. 5 (1954) pp. 447-455.
4. E. Hewitt, *Rings of real-valued continuous functions*. I, Trans. Amer. Math. Soc. vol. 64 (1948) pp. 45-99.
5. L. E. Pурсell, *An algebraic characterization of fixed ideals in certain function rings*, Pacific Journal of Mathematics vol. 5 (1955) pp. 963-969.

GRINNELL COLLEGE