

# ON A TAUBERIAN THEOREM OF LANDAU

BASIL GORDON

1. **Introduction.** It was the plan of Tschebyschef [1] and Sylvester [2] to deduce the prime number theorem<sup>1</sup>

$$\psi(x) \sim x$$

from the formula

$$T(x) = \sum_{n \leq x} \psi\left(\frac{x}{n}\right) = \log [x]!$$

However they succeeded only in proving the existence of positive constants  $c_1$  and  $c_2$  such that

$$c_1x < \psi(x) < c_2x.$$

Landau discusses this problem in his Handbuch [3, pp. 79–83], where he proves that if  $A(x)$  is any monotone nondecreasing function of  $x$  such that

$$T(x) = \sum_{n \leq x} A\left(\frac{x}{n}\right) = x \log x + bx + o(x)$$

with  $b$  a constant, then

$$(1) \quad c_1x < A(x) < c_2x,$$

where  $c_1$  and  $c_2$  are positive constants. This gives Tschebyschef's theorem if we take

$$A(x) = \psi(x)$$

and use the fact that

$$\log [x]! = x \log x - x + O(\log x).$$

Landau states that if only the condition

$$T(x) = x \log x + O(x)$$

is assumed, then (1) does not follow, a remark whose incorrectness was proved by H. N. Shapiro [4]. Landau then goes on to prove [3, pp. 598–604] that if

Received by the editors June 14, 1957 and, in revised form, November 25, 1957 and February 24, 1958.

<sup>1</sup> Here and in the sequel,  $\psi(x)$ ,  $\wedge(n)$ ,  $\mu(n)$ , and  $\theta(x)$  denote the usual functions of prime number theory.

$$T(x) = x \log x + bx + o\left(\frac{x}{\log^2 x}\right),$$

then  $A(x) \sim x$ . The proof requires the formula

$$(2) \quad \sum_{n=1}^{\infty} \frac{\mu(n) \log n}{n} = -1,$$

and since this is “deeper” than the prime number theorem, he concludes that Tschebyschef was foredoomed to failure. In the present paper we shall correct this statement by showing that a slightly stronger theorem than Landau’s can be obtained using only the prime number theorem. The theorem to be proved is

**THEOREM 1.** *If  $A(x)$  is nondecreasing, and*

$$(3) \quad T(x) = x \log x + bx + o\left(\frac{x}{\log x}\right),$$

then  $A(x) \sim x$ .

It should be mentioned that the best theorem in this direction has been obtained by A. E. Ingham [5], but the proof makes use of Wiener’s Tauberian theorem, and therefore cannot be considered elementary.

**2. A Selberg-like formula.** We shall require the following elementary lemma:

**LEMMA 1.** *If  $F(x)$  is any function defined for  $x \geq 1$ , and if*

$$G(x) = \log x \sum_{n \leq x} F\left(\frac{x}{n}\right),$$

then

$$F(x) \log x + \sum_{n \leq x} F\left(\frac{x}{n}\right) \Lambda(n) = \sum_{n \leq x} \mu(n) G\left(\frac{x}{n}\right).$$

The proof is a straight-forward application of the Möbius inversion formula. For details, see Trost [6, p. 66]. Now let  $A(x)$  be a function for which the hypothesis (3) is satisfied, and apply Lemma 1 with  $F(x) = A(x)$ . The corresponding  $G(x)$  is

$$G(x) = \log x \sum_{n \leq x} A\left(\frac{x}{n}\right) = T(x) \log x = x(\log x)^2 + bx \log x + o(x),$$

and so by Lemma 1,

$$(4) \quad A(x) \log x + \sum_{n \leq x} A\left(\frac{x}{n}\right) \Lambda(n) \\ = \sum_{n \leq x} \mu(n) \frac{x}{n} \left(\log \frac{x}{n}\right)^2 + \sum_{n \leq x} \mu(n) b \frac{x}{n} \log \frac{x}{n} + \sum_{n \leq x} \mu(n) o\left(\frac{x}{n}\right).$$

It is well known that the right side of (4) equals  $2x \log x + o(x \log x)$  so that we obtain

**THEOREM 2.** *If  $A(x)$  satisfies condition (3), then*

$$(5) \quad A(x) \log x + \sum_{n \leq x} A\left(\frac{x}{n}\right) \Lambda(n) = 2x \log x + o(x \log x).$$

It can be seen that this result is very similar to the Selberg formula for  $\theta(x)$  with a weaker error term (cf. [7], especially the remark on p. 313).

**3. Proof of the main theorem.** We shall now derive Theorem 1 from the identity (5), and it is here that the monotonicity of  $A(x)$  becomes important. Note first that (5) and the fact that  $A(x)$  is nondecreasing imply that  $A(x) > 0$  for all sufficiently large  $x$ . This fact together with (5) then yields that for  $x = y + o(x)$  we have  $A(x) = A(y) + o(x)$ . Thus it is easily seen that without loss of generality we may assume that for all  $x > 0$ , we have  $A(x) = A([x])$ . Since  $A(x)$  is nondecreasing we can then write

$$A(x) = \sum_{m \leq x} a_m,$$

where  $a_m \geq 0$ . We can then rewrite (5) as

$$A(x) \log x + \sum_{mn \leq x} a_m \Lambda(n) = 2x \log x + o(x \log x),$$

or equivalently

$$(6) \quad A(x) \log x + \sum_{m \leq x} a_m \psi\left(\frac{x}{m}\right) = 2x \log x + o(x \log x).$$

From the prime number theorem and the result of [4] (in particular Equation (6) therein) it follows that

$$\sum_{m \leq x} a_m \psi\left(\frac{x}{m}\right) = (1 + o(1))x \sum_{m \leq x} \frac{a_m}{m} \sim x \log x,$$

which together with (6) above yields  $A(x) \sim x$ , the desired result.

The following application of Theorem 1 is interesting. Let  $A(x) = [x] + M(x)$ , where  $M(x) = \sum_{n \leq x} \mu(n)$ . Since  $1 + \mu(n) \geq 0$ ,  $A(x)$  is

monotone. The corresponding  $T(x)$  is

$$\begin{aligned} T(x) &= \sum_{n \leq x} \left( \left[ \frac{x}{n} \right] + M\left(\frac{x}{n}\right) \right) \\ &= x \log x + cx + O(x^{1/2}) + \sum_{n \leq x} \mu(n) \left[ \frac{x}{n} \right] \\ &= x \log x + cx + O(x^{1/2}) \end{aligned}$$

since  $\sum_{n \leq x} \mu(n) [x/n] = 1$  (cf. [3, pp. 576–577]). Thus the hypothesis of Theorem 1 is satisfied and we conclude that  $A(x) \sim x$ , which is here the same as asserting  $M(x) = o(x)$ . The above thus provides another proof of the result of Landau [8] which asserts that  $\psi(x) \sim x$  implies  $M(x) = o(x)$ .

#### REFERENCES

1. P. L. Tschebyschef, *Mémoire sur les nombres premiers*, J. Math. Pures Appl. series 1, vol. 17 (1852) pp. 366–390.
2. J. J. Sylvester, *On Tschebyschef's theorem of the totality of prime numbers comprised within given limits*, Amer. J. Math. vol. 4 (1881) pp. 230–247.
3. E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen*, Leipzig, 1909.
4. H. N. Shapiro, *On the number of primes less than or equal to  $x$* , Proc. Amer. Math. Soc. vol. 1 (1950) pp. 346–348.
5. A. E. Ingham, *Some Tauberian theorems connected with the prime number theorem*, J. London Math. Soc. vol. 20 (1945) pp. 171–180.
6. E. Trost, *Primzahlen*, Basel, 1953.
7. A. Selberg, *An elementary proof of the prime number theorem*, Ann. of Math. vol. 50 (1949) pp. 305–313.
8. E. Landau, *Über den Zusammenhang einiger neuerer Sätze der analytischen Zahlentheorie*, Wiener Sitzungsberichte, Math. Klasse, vol. 115 (1906) pp. 589–632.
9. P. Erdős, *On a new method in elementary number theory which leads to an elementary proof of the prime number theorem*, Proc. Nat. Acad. Sci. U.S.A. vol. 35 (1949) pp. 374–384.

CALIFORNIA INSTITUTE OF TECHNOLOGY