

## PROXIMITY MAPS FOR CONVEX SETS

WARD CHENEY AND ALLEN A. GOLDSTEIN

The method of successive approximation is applied to the problem of obtaining points of minimum distance on two convex sets. Specifically, given a closed convex set  $K$  in Hilbert space, let  $P$  be the map which associates with each point  $x$  the point  $Px$  of  $K$  closest to  $x$ . That  $P$  is well-defined is proved in [1, p. 6].  $P$  will be called the *proximity map* for  $K$ . If there are two such sets,  $K_1$  and  $K_2$ , let  $Q$  denote the composition  $P_1P_2$  of their proximity maps. It is shown that every fixed point of  $Q$  is a point of  $K_1$  closest to  $K_2$ , and that the fixed points of  $Q$  may be obtained by iteration of  $Q$  when one of the sets is compact or when both are polytopes in  $E_n$ . An application to the solution of linear inequalities is cited. Our thanks are due the referee for having suggested substantial simplifications.

**THEOREM 1.** *Let  $Q$  be a map of a metric space into itself such that*

- (i)  $d(Qx, Qy) \leq d(x, y)$ ,
- (ii) if  $x \neq Qx$ , then  $d(Qx, Q^2x) < d(x, Qx)$ ,
- (iii) for each  $x$ , the sequence  $Q^n x$  has a cluster point. Then for each  $x$ , the sequence  $Q^n x$  converges to a fixed point of  $Q$ .

**PROOF.** By (i), the sequence  $d(Q^n x, Q^{n+1} x)$  is nonincreasing. Let  $y$  be a cluster point of  $Q^n x$ , say  $y = \lim_k Q^{n_k} x$ . By (i),  $Q$  is continuous; therefore  $d(y, Qy) = \lim_k d(Q^{n_k} x, Q^{n_k+1} x) = \lim_n d(Q^n x, Q^{n+1} x) = \lim_k d(Q^{n_k+1}, Q^{n_k+2} x) = d(Qy, Q^2 y)$  contrary to (ii) unless  $y = Qy$ . From (i) it follows that for all  $n$ ,  $d(Q^{N+n} x, y) \leq d(Q^N x, y)$  whence  $Q^n x \rightarrow y$ .

**COROLLARY.** *Let  $Q$  be a map of a normed linear space into itself having the property  $\|Qx - Qy\| \leq \|x - y\|$ , equality holding only if  $x = y$ . Let  $R = \alpha Q + (1 - \alpha)I$ , ( $0 < \alpha \leq 1$ ). If the range of  $R$  is compact, then  $Q$  has a unique fixed point which is the limit of every sequence  $R^n x$  with  $x$  arbitrary. (For related results, see [2].)*

**LEMMA.** *Let  $K$  be a convex set in Hilbert space. A point  $b \in K$  is nearest a point  $a \notin K$  if and only if  $s \equiv (x - b, b - a) \geq 0$  for all  $x \in K$ .*

**PROOF.** Suppose  $b$  nearest  $a$ , and let  $x$  be arbitrary in  $K$ . When  $0 \leq t \leq 1$ ,  $tx + (1 - t)b \in K$ . Thus  $0 \leq \|a - tx - (1 - t)b\|^2 - \|a - b\|^2 = t^2 \|b - x\|^2 + 2ts$ . But this inequality would be violated by small  $t$

---

Presented to the Society, January 22, 1959; received by the editors July 26, 1958 and, in revised form, October 4, 1958.

unless  $s \geq 0$ . For the converse, suppose  $s \geq 0$ . Then  $\|x - a\|^2 - \|a - b\|^2 = (x, x) - 2(a, x) + 2(a, b) - (b, b) = (x - b, x - b) + 2s \geq 0$ .

**THEOREM 2.** *Let  $K_1$  and  $K_2$  be two closed convex sets in Hilbert space. Let  $P_i$  denote the proximity map for  $K_i$ . Any fixed point of  $P_1P_2$  is a point of  $K_1$  nearest  $K_2$ , and conversely.*

**PROOF.** Suppose  $y = P_2x$  and  $x = P_1y$ . If  $x = y$ , the distance between  $K_1$  and  $K_2$  is thereby attained. Otherwise  $x \notin K_2$  and  $y \notin K_1$ . If  $u$  is arbitrary in  $K_1$ , then by the Lemma,  $(u - x, x - y) \geq 0$  whence  $(u, x - y) \geq (x, x - y)$ . Similarly for arbitrary  $v \in K_2$ ,  $(v, y - x) \geq (y, y - x)$ . Addition yields  $(u - v, x - y) \geq (x - y, x - y)$  from which via the Schwartz inequality,  $\|u - v\| \geq \|x - y\|$ . For the converse, suppose that  $\|x - P_2x\| \leq \|z - P_2z\|$  for all  $z \in K_1$ . Setting  $z = P_1P_2x$  we have  $\|z - P_2z\| \leq \|z - P_2x\| \leq \|x - P_2x\| \leq \|z - P_2z\|$  whence  $z = x$  by the uniqueness of  $z$ .

**THEOREM 3.** *The proximity map  $P$  for a closed convex set  $K$  in Hilbert space satisfies the Lipschitz condition  $\|Px - Py\| \leq \|x - y\|$ , equality holding only if  $\|x - Px\| = \|y - Py\|$ .*

**PROOF.** By the Lemma,  $A \equiv (Px - Py, Py - y) \geq 0$  and  $B \equiv (Py - Px, Px - x) \geq 0$ . Regrouping terms in the inequality  $A + B \geq 0$  and using the Schwartz inequality, one has  $\|Py - Px\| \cdot \|y - x\| \geq (Py - Px, y - x) \geq (Py - Px, Py - Px) = \|Py - Px\|^2$ , whence  $\|y - x\| \geq \|Py - Px\|$ . Equality holds here only if  $A = B = 0$  and if  $Py - Px = \lambda(y - x)$ . Thus  $C \equiv (y - x, Py - y) = 0$  and  $D \equiv (y - x, Px - x) = 0$ . A computation shows then that  $0 = A - B + C + D = \|Px - x\|^2 - \|Py - y\|^2$ .

**THEOREM 4.** *Let  $K_1$  and  $K_2$  be two closed convex sets in Hilbert space and  $Q$  the composition  $P_1P_2$  of their proximity maps. Convergence of  $Q^n x$  to a fixed point of  $Q$  is assured when either (a) one set is compact, or (b) one set is finite dimensional and the distance between the sets is attained.*

**PROOF.** Theorem 3 implies that  $Q$  satisfies (i) of Theorem 1. If  $y \equiv Qx \neq x \in K_1$ , then  $\|y - P_2y\| \leq \|y - P_2x\| < \|x - P_2x\|$ . By Theorem 3,  $\|Qx - Qy\| \leq \|P_2x - P_2y\| < \|x - y\|$ . Thus  $Q$  satisfies (ii) of Theorem 1. If the distance between the sets is attained, then  $Q$  has a fixed point  $y$  by Theorem 2, and  $\|Q^n x\| \leq \|Q^n x - Qy\| + \|y\| \leq \|x - y\| + \|y\|$ . Boundedness of  $\|Q^n x\|$  and finite-dimensionality of  $K_1$  suffice for (iii) of Theorem 1.  $Q' = P_2P_1$  replaces  $Q$  in these arguments when  $K_2$  replaces  $K_1$ . But if  $y$  is a fixed point of  $Q'$ , then  $P_1y$  is a fixed point of  $Q$ .

**THEOREM 5.** *In a finite dimensional Euclidean space, the distance*

between two polytopes is attained, a polytope being the intersection of a finite family of halfspaces.

PROOF. First, the case when both polytopes are linear manifolds. Let  $K_1$  be the linear span of  $\{x_1, \dots, x_m\}$  and  $K_2$  the linear span of  $\{y_1, \dots, y_n\}$  translated by a vector  $cy_0$ . Assume the  $x$ 's and  $y$ 's each form orthonormal sets. A point  $x = \sum \xi_i x_i$  is sought which minimizes  $G = \|cy_0 + \sum (x, y_i)y_i - x\|^2$ .  $G$  is a positive definite quadratic function of  $\xi_1, \dots, \xi_m$ , and therefore attains a minimum.

Now assume the validity of the theorem when one polytope is of dimension less than  $n$  and the other is a linear manifold. (The validity for  $n=1$  being established by the above.) Let  $K_1$  be a polytope of dimension  $n$  and  $K_2$  a linear manifold. On each proper face of  $K_1$  there is a point nearest  $K_2$ . There being only a finite number of faces, either one of these points is the required one or there is a point  $x_0 \in K_1$  such that  $d(x_0, K_2) < d(F, K_2)$  for all proper faces  $F$  of  $K_1$ . In the latter case, let  $y_0$  be chosen nearest to  $K_2$  in the least linear manifold containing  $K_1$ . Since  $d(x, K_2)$  is a convex function of  $x$ , the line segment  $x_0 y_0$  contains no point of any proper face of  $K_1$ . Therefore  $y_0 \in K_1$  and must be the required point. The proof for the remaining case is as above, mutatis mutandis.

LINEAR INEQUALITIES. Consider the system of linear inequalities  $(A^i, x) \leq b_i$  where  $x \in E_n$  and  $1 \leq i \leq m$ . Let  $K_1$  denote the range of the matrix  $A$  and  $K_2$  the orthant  $\{y \in E_m: y_i \leq b_i \text{ all } i\}$ . If  $K_1$  and  $K_2$  have a point in common, then any  $x$  for which  $Ax \in K_1 \cap K_2$  is a solution of the system. Even if the system is inconsistent, a point on  $K_1$  closest to  $K_2$  may be obtained by iteration of the map  $Q = P_1 P_2$ . These proximity maps are defined as follows. If  $A$  is of rank  $n$ ,  $P_1 = A(A^T A)^{-1} A^T$ . If  $A$  is not necessarily of rank  $n$ , one may write  $P_1 = BB^T$  where  $B$  is a column-orthogonal matrix whose range is that of  $A$ .  $P_2 y = z$  if and only if  $z_i = y_i$  when  $y_i \leq b_i$  and  $z_i = b_i$  otherwise. Convergence of each sequence  $Q^n x$  is guaranteed by Theorem 4.

#### BIBLIOGRAPHY

1. B. v.-Sz. Nagy, *Spektralardarstellung Linearer Transformationen des Hilbertschen Raumes*, Berlin, 1942.
2. H. Schaefer, *Über die Methode Sukzessiver Approximationen*, Jber. Deutsch. Math. Verein. vol. 59 (1957) pp. 131-140.

CONVAIR ASTRONAUTICS, SAN DIEGO