

COMPACTIFICATIONS OF DIMENSION ZERO

L. J. HEIDER¹

1. Introduction. A c.r. space X is here understood to be a completely regular T_1 space. A continuous function on X is understood to be real-valued, and the symbol $C(X)$ denotes the collection of all such functions. For each element f of $C(X)$, let $N(f) = [x: f(x) < 0]$. Consider the following requirements with respect of the set $N(f)$:

- (I) $N(f)$ is closed (and open);
- (II) $\overline{N}(f)$ is open (and closed);
- (III) $N(f)$ is a countable union of open-and-closed (cloven) sets.

The c.r. spaces, in which each set $N(f)$ satisfies (I), are the P spaces of [2; 3] and enjoy many interesting alternate characterizations. The c.r. spaces, in which each set $\overline{N}(f)$ satisfies (II), are characterized as the c.r. spaces X for which the lattice $C(X)$ is conditionally σ -complete, or, equivalently, for which the Stone-Ćech compactification space βX is the Boolean representation space of a σ -complete Boolean algebra [3; 7].

Requirement (III) drastically weakens (I). Yet, the first purpose of this note is to show that (III) characterizes an important class of c.r. spaces, viz., those c.r. spaces X for which βX is of dimension zero, in the sense of possessing a base of cloven sets. From this it will follow that (I) \Rightarrow (II) \Rightarrow (III).

If X is of dimension zero, attention focuses on the field of cloven subsets of X . If, moreover, βX is of dimension zero, it is the Boolean representation space of this field of sets, viewed as partially ordered by the inclusion relation. These are set-theoretic considerations. Opposed to this, requirement (III) involves the consideration of continuous functions. In addition to (III), there are two established [3; 5] characterizations of the c.r. spaces X for which βX is of dimension zero: (i) for each pair A and B of separated sets in X , there is a cloven set in X containing A and disjoint from B ; (ii) each finite normal open covering of X possesses a refinement which is a finite partition of X into cloven sets. Condition (i), in the concept of separated sets, refers to continuous functions. The concept of a normal covering, employed in condition (ii), is difficult of expression in any form and is, perhaps, best expressed in terms of continuous functions [4]. The second purpose of this note is, therefore, to replace condi-

Presented to the Society, November 28, 1958; received by the editors July 16, 1958.

¹ This research was supported by the National Science Foundation.

tions (i) and (ii) and likewise requirement (III) by an equivalent condition based exclusively on set-theoretic concepts. This is accomplished for all c.r. spaces, but with the most notable simplicity for normal spaces.

For c.r. spaces X with βX of dimension zero, the preceding emphasis on set-theoretic techniques should extend to the consideration of continuous functions. The concluding section of this note describes a countable, purely set-theoretic technique for construction of all continuous functions pertaining to a c.r. space X with βX of dimension zero.

2. Compactifications of dimension zero. Let X be a c.r. space, and let f be an element of $C(X)$ with finite bounds a and b . Then, in terms of Stone's spectral representation of continuous functions [7], to each real number r there is assigned an open subset $E_f(r)$ of X such that:

- (1) $E_f(r) = \emptyset$ for all $r \leq a$,
- (2) $E_f(r) = \bigcup_{r' < r} E_f(r')$ for all r ,
- (3) $\overline{E}_f(r') \subseteq E_f(r)$ for $r' < r$,
- (4) $E_f(r) = X$ for $r > b$.

Conversely, any family $[E(r)]_{r \in \mathbb{R}}$ of open subsets of X , satisfying conditions (1)–(4) for some pair $a \leq b$ of real numbers, determines uniquely a bounded continuous function f on X with $[x: f(x) < r] = E(r)$.

The significance of requirement (III) of the introduction is now partly revealed.

THEOREM 2.1. *If a c.r. space X is such that βX is of dimension zero, then each spectral set $E_f(r)$, and thus the set $N(f)$, of each element f of $C(X)$ is a countable union of cloven subsets of X .*

PROOF. Let f be a bounded continuous function on X , and let \tilde{f} be its unique extension over βX , the latter being assumed of dimension zero. For arbitrary real numbers $r_1 < r_2$, $E_{\tilde{f}}(r_1) \subseteq [p \in \beta X: \tilde{f}(p) \leq r_1] \subseteq E_{\tilde{f}}(r_2)$. With βX compact and of dimension zero, a cloven subset G of βX may be chosen with $[p \in \beta X: \tilde{f}(p) \leq r_1] \subseteq G \subseteq E_{\tilde{f}}(r_2)$. Then $E_f(r_1) \subseteq 0 \subseteq E_f(r_2)$, where 0 is the cloven residue in X of the cloven subset G of βX .

Finally, with $\{r_n\}$ a monotone increasing sequence of numbers approaching the number r as a limit, from $E_f(r) = \bigcup_{r' < r} E_f(r') = \bigcup_{n=1}^{\infty} E_f(r_n)$, one concludes that $E_f(r)$ is a countable union of cloven subsets of X .

Assume, now, that the c.r. space X is such that each set $N(f)$, for f in $C(X)$, is a countable union of cloven subsets of X . Note that $N(-f) = P(f) = [x: f(x) > 0]$ and $E_f(r) = N(f - r)$. With $r_1 < r_2$, for any

element f of $C(X)$, the set $[x: f(x) \leq r_1]$ is a countable intersection of cloven subsets and is contained in the countable union of cloven sets $[x: f(x) < r_2]$. The following lemma is now useful:

LEMMA 2.2. *If $\{0_n\}$ and $\{G_n\}$ are two sequences of cloven subsets of a topological space X , with $\bigcap_{n=1}^{\infty} 0_n \subseteq \bigcup_{n=1}^{\infty} G_n$, then there exists a cloven subset L of X with $\bigcap_{n=1}^{\infty} 0_n \subseteq L \subseteq \bigcup_{n=1}^{\infty} G_n$.*

PROOF. Form a sequence $\{H_n\}$ of cloven sets, with $H_{2n-1} = G_n$ and $H_{2n} = \tilde{0}_n$, where $\tilde{0}_n$ denotes the cloven complement in X of 0_n . Each point of X is in some set H_n . Next form $\{H_n^*\}$ with $H_n^* = H_n - \bigcup_{1 \leq i < n} H_i$. The sets H_n^* are cloven, pairwise disjoint and cover X . Finally, let L denote the point-set union of all H_n^* containing points of the set $\bigcap_{n=1}^{\infty} 0_n$, while M denotes the union of the remaining H_n^* . Then L and M , as disjoint unions of open sets with $L \cup M = X$, are seen to be cloven subsets of X . Finally, $\bigcap_{n=1}^{\infty} 0_n \subseteq L \subseteq \bigcup_{n=1}^{\infty} G_n$ as desired.

Let X continue to denote a c.r. space for which all sets $N(f)$, with f in $C(X)$, are countable unions of cloven sets. It is clear that the cloven sets then constitute a base for the topology on X . These cloven sets, as ordered by the inclusion relation, form a Boolean algebra. Let ηX denote the Boolean representation space of this algebra. The points of ηX will here be regarded as prime dual ideals of cloven sets of X , i.e., proper subsets of the collection of all cloven sets, closed under finite intersections of their elements, and maximal with respect to these properties. Each point of X determines, uniquely, such a prime dual ideal, and X is seen to be imbedded as a dense subspace of ηX .

In general, for any c.r. space X of dimension zero, the space ηX serves as a maximal compact space of dimension zero, containing X as a dense subspace [1]. However, under the present assumptions, $\eta X = \beta X$, the Stone-Čech compactification space of X . Thus, as appears from the preceding discussion, for any (bounded) element f of $C(X)$ and any pair of numbers $r_1 < r_2$, there exists a cloven subset 0 of X with $E_f(r_1) \subseteq 0 \subseteq E_f(r_2)$. However, each prime dual ideal (point) of ηX contains either this cloven set 0 or its complement. Then, using only the fact that a sequence of closed, nested intervals of the real line, of widths approaching zero, determines a unique real number, for each prime dual ideal (point) α of ηX and for each bounded element f of $C(X)$, it is possible to assign a real number $\tilde{f}(\alpha)$ such that, for each $\epsilon > 0$, there is a cloven set 0 in α with $|\tilde{f}(\alpha) - f(x)| < \epsilon$ for each point x of X in 0 . Clearly the function \tilde{f} thus defined on ηX is a continuous extension of the given bounded element f of $C(X)$.

In view of this discussion and of Theorem 2.1, our first contention is now established.

THEOREM 2.3. *The c.r. spaces X for which βX is of dimension zero are the spaces X for which each set $N(f)$, f in $C(X)$, is a countable union of cloven sets.*

3. A set-theoretic characterization. The purpose of this section is to reduce the characterization of the c.r. spaces X for which βX is of dimension zero to the simplest possible set-theoretic terms. To this end, a definition and theorem of recent origin [6] will be of service.

DEFINITION. Let \mathfrak{M} be any family of open subsets of a c.r. space X . Then \mathfrak{M} will be called a *completely regular system of open sets* if, for each element M of \mathfrak{M} , there are sequences $\{M'_n\}$ and $\{M''_n\}$ of elements of \mathfrak{M} with $M'_n \subseteq \tilde{M}''_n \subseteq M$ and $M = \bigcup_{n=1}^{\infty} M'_n$. (Here \tilde{M}''_n denotes the complement in X of the set M''_n .)

For any element f of $C(X)$, with X a c.r. space, the system of all subsets $f^{-1}(0)$ of X , with 0 an open subset of the real line, is clearly a completely regular system of open subsets of X . However, even more can be asserted.

THEOREM. *Every element of any completely regular system of open sets of a c.r. space X has the form $f^{-1}(0)$ for some element f of $C(X)$ and some open subset 0 of the real line.*

The proof of this theorem, as it appears in [6], is an interpretation of involved properties of certain abstract systems. However, the theorem itself has a simple application to our present task.

THEOREM 3.1. *The c.r. spaces X for which βX is of dimension zero are the spaces in which the system of all countable unions of cloven sets includes every completely regular system of open sets.*

To establish this theorem, it needs only to be noted that sets of the type $f^{-1}(0)$ include all spectral sets, while each set of the type $f^{-1}(0)$ is a countable union of finite intersections of spectral sets $E_f(r)$ and $E_g(r)$, where $g = -f$ and f is an element of $C(X)$. For any c.r. space X , the system of all countable unions of cloven sets obviously constitutes a completely regular system of open sets. Thus, the theorem simply requires that this particular system be the greatest such completely regular system of open sets.

If βX is to be of dimension zero, the topology of X must be based on the field \mathfrak{F} of cloven subsets of X . This field must be *reduced*, in the sense that distinct points of X are contained in complementary elements of \mathfrak{F} , and it must be *union-intersection closed*, in the sense that any subset of X which is both an intersection and a union of elements of \mathfrak{F} is itself an element of \mathfrak{F} .

Now let \mathfrak{F} be such a reduced, union-intersection closed field of subsets of a set X . The symbol 0 , with or without subscript, will be reserved to denote elements of \mathfrak{F} . Let $X(\mathfrak{F})$ indicate the c.r. space obtained by using the elements of \mathfrak{F} as a base for open sets in X . The symbol $\bigcup_0 \mu$ will denote a general open set of $X(\mathfrak{F})$, while $\bigcup_r 0_\mu$ will indicate an open set contained in a completely regular system of open sets, and $\bigcap_r 0_r$ will indicate the complement in X of such a set. Finally, let $\eta X(\mathfrak{F})$ represent the Boolean representation space of \mathfrak{F} . The basic question concerns set-theoretic conditions under which $\eta X(\mathfrak{F}) = \beta X(\mathfrak{F})$.

THEOREM 3.2. *The condition $\{\bigcap_r 0_r \subseteq \bigcup_r 0_\mu \Rightarrow \exists 0 \text{ with } \bigcap_r 0_r \subseteq 0 \subseteq \bigcup_r 0_\mu\}$ is equivalent to the condition that $\eta X(\mathfrak{F}) = \beta X(\mathfrak{F})$.*

THEOREM 3.3. *The condition $\{\bigcap 0_r \subseteq \bigcup 0_\mu \Rightarrow \exists 0 \text{ with } \bigcap 0_r \subseteq 0 \subseteq \bigcup 0_\mu\}$ is equivalent to the double condition that $X(\mathfrak{F})$ be normal and $\eta X(\mathfrak{F}) = \beta X(\mathfrak{F})$.*

As regards Theorem 3.2, if $\eta X(\mathfrak{F}) = \beta X(\mathfrak{F})$, then, by Theorem 3.1, each $\bigcap_r 0_r$ is a countable intersection and each $\bigcup_r 0_\mu$ is a countable union, and Lemma 2.2 applies. Conversely, if $\bigcap_r 0_r \subseteq \bigcup_r 0_\mu$ always allows $\bigcap_r 0_r \subseteq 0 \subseteq \bigcup_r 0_\mu$, then each $\bigcup_r 0_\mu$, as a member of a completely regular system of open sets, is seen to be a countable union of cloven sets, and Theorem 3.1 applies.

As regards Theorem 3.3, if $\bigcap 0_r \subseteq \bigcup 0_\mu$ always allows $\bigcap 0_r \subseteq 0 \subseteq \bigcup 0_\mu$, then $X(\mathfrak{F})$ is obviously normal and, by Theorem 3.2, $\eta X(\mathfrak{F}) = \beta X(\mathfrak{F})$. Conversely, with $\beta X(\mathfrak{F}) = \eta X(\mathfrak{F})$ and thus of dimension zero, from the normality of $X(\mathfrak{F})$ it is clear that $\bigcap 0_r \subseteq \bigcup 0_\mu$ always allows $\bigcap 0_r \subseteq 0 \subseteq \bigcup 0_\mu$.

As an application of Theorem 3.3, with X , \mathfrak{F} , $X(\mathfrak{F})$ and $\eta X(\mathfrak{F})$ as described above, certain cases wherein $X(\mathfrak{F})$ is normal and $\eta X(\mathfrak{F}) = \beta X(\mathfrak{F})$ may be noted:

- (a) $X(\mathfrak{F})$ possesses a countable base of open sets;
- (b) each $\bigcup 0_\mu$ may be represented as a countable union of cloven sets;
- (c) $X(\mathfrak{F})$ has the Lindelöf property;
- (d) each binary covering of $X(\mathfrak{F})$ by open sets may be refined by a partition of X into cloven sets.

As an application of Theorem 3.2, the following case is of some importance. Let \mathfrak{F} continue to denote a reduced, union-intersection closed field of subsets of a set X . However, now also assume that \mathfrak{F} is a σ -field, in the sense that any countable union of elements of \mathfrak{F} is again an element of \mathfrak{F} . Let \mathfrak{M} denote a completely regular system of

open subsets of $X(\mathfrak{F})$, and let M be an element of \mathfrak{N} . Then, by definition, there exists a sequence $\{M_n''\}$ of elements of \mathfrak{N} with $M = \bigcup_{n=1}^{\infty} \tilde{M}_n''$. Each element \tilde{M}_n'' is of the form $\tilde{M}_n'' = \bigcap_{\nu \in I_n} 0_{n,\nu}$ where each $0_{n,\nu}$ is an element of \mathfrak{F} and each I_n is an index set. Then

$$M = \bigcup_{n=1}^{\infty} \left[\bigcap_{\nu \in I_n} 0_{n,\nu} \right] = \bigcap_{h \in H} \left[\bigcup_{n=1}^{\infty} 0_{n,h(n)} \right]$$

where H is the collection of all functions h on the positive integers with $h(n)$ in I_n . However, each $\bigcup_{n=1}^{\infty} 0_{n,h(n)}$, as a countable union of elements of \mathfrak{F} , is an element 0_h of \mathfrak{F} . Hence $M = \bigcap_{h \in H} 0_h$, and M is seen to be a cloven set in $X(\mathfrak{F})$ and thus an element of \mathfrak{F} . The c.r. spaces $X(\mathfrak{F})$ constructed with elements of a reduced, union-intersection closed, σ -field \mathfrak{F} of subsets of X as a base for open sets are identical with the P spaces of [2; 3], i.e., the c.r. spaces X for which each set $N(f)$, f in $C(X)$, is closed (and open).

4. The construction of continuous functions. Let the c.r. space X be such that βX is of dimension zero. Let f be an element of $C(X)$ and, for the sake of simplicity, assume that f is bounded with $0 \leq f(x) \leq 1$ throughout X . As indicated in the proof of Theorem 2.1, for each pair $r_1 < r_2$ of real numbers, there is a cloven subset 0 of X with $E_f(r_1) \subseteq 0 \subseteq E_f(r_2)$. Now let $r_2 = p/2^n$ where p is an odd positive integer $1 \leq p < 2^n$ and let $r_1 = (2p - 1)/2^{n+1}$. Let $[p/2^n]$ symbolize a cloven set such that $E_f(2p - 1/2^{n+1}) \subseteq [p/2^n] \subseteq E_f(p/2^n)$.

As $r_2 = p/2^n$ exhausts the sets $1/2; 1/4, 3/4; 1/8, \dots$, an array of cloven sets is formed:

$$\begin{aligned} & [1/2] \\ & [1/4] \subseteq [1/2] \subseteq [3/4] \\ & [1/8] \subseteq [1/4] \subseteq [3/8]; [1/2] \} \subseteq [5/8] \subseteq [3/4] \subseteq [7/8] \\ & [1/16] \subseteq [1/8] \subseteq [3/16]; [1/4] \} \subseteq [5/16] \subseteq [3/8] \subseteq [7/16]; \\ & [1/2] \} \subseteq [9/16] \subseteq \dots \\ & \dots \end{aligned}$$

Here \subseteq denotes set inclusion. The symbol $\} \subseteq [p/2^n]$ indicates that the set $[p/2^n]$ contains every set to the left of the bracket in the row under consideration. Finally the presence of the semi-colon, rather than the inclusion symbol, indicates that no inclusion relation is asserted in regard to the sets immediately adjacent to the semi-colon.

BIBLIOGRAPHY

1. B. Banaschewski, *Über nulldimensionale Räume*, Math. Nachr. vol. 13 (1955) pp. 129–140.
2. L. Gillman and M. Henriksen, *Concerning rings of continuous functions*, Trans. Amer. Math. Soc. vol. 77 (1954) pp. 340–362.
3. ———, *Rings of continuous functions in which every finitely generated ideal is principal*, *ibid.* vol. 82 (1956) pp. 366–391.
4. M. Henriksen and J. Isbell, *Local connectedness in the Stone-Čech compactification*, Illinois J. Math. vol. 1 (1957) pp. 574–582.
5. J. Isbell, *Zero-dimensional spaces*, Tôhoku Math. J. vol. 7 (1955) pp. 1–8.
6. J. Kerstan, *Eine Charakterisierung der vollständig regulären Räume*, Math. Nachr. vol. 17 (1958) pp. 27–46.
7. M. Stone, *Boundedness properties in function lattices*, Canad. J. Math. vol. 1 (1949) pp. 176–186.

INSTITUTE FOR ADVANCED STUDY AND
MARQUETTE UNIVERSITY