

ON THE CONSTRUCTION OF PERIODIC MAPS WITHOUT FIXED POINTS¹

P. E. CONNER AND E. E. FLOYD

1. Introduction. We will consider the existence problem for fixed points of periodic maps. When the period is prime power, the theorem of P. A. Smith is available [5]. That is, if X is a locally compact, finite dimensional space whose mod p Čech cohomology groups are trivial, and if $T: X \rightarrow X$ is of period p^n where p is a prime, then T has a fixed point. It is shown here that such propositions are false in case the period is not prime power.

In §4 we show that if r is a positive integer greater than one which is not prime power then there exists a compact, finite dimensional space X with trivial integral Čech cohomology groups and a map $T: X \rightarrow X$ of period r which is without fixed points. In §6, with r as above, it is shown that there exists a contractible manifold M and a map $T: M \rightarrow M$ of period r and without fixed points. It seems reasonable to conjecture that the manifold M constructed is actually a Euclidean space E^n . If this could be proved, the Smith conjecture that periodic maps on E^n have a fixed point would be answered in the negative.

2. Commuting maps; the prime power case. Our constructions will be based on the study of maps $f: X \rightarrow X$ which commute with a given periodic map T . In this section we point out some of the facts in the prime power case.

We say that an endomorphism h of the group G is nilpotent if for each $x \in G$ we have $h^n(x) = 0$ for n sufficiently large. In the following, $H^i(X, K)$ denotes the Čech cohomology group with coefficients in K , and $H^0(X; K)$ denotes the reduced 0-dimensional group.

(2.1) *Suppose that T is a periodic map of prime power period p^n on the compact, nonempty, finite dimensional space X . Suppose also that $f: X \rightarrow X$ is a map such that $fT = Tf$ and that $f^*: H^i(X, Z_p) \rightarrow H^i(X, Z_p)$ is nilpotent for $i = 0, 1, \dots$. If F denotes the fixed point set of T , then F is nonempty and $(f|F)^*: H^i(F, Z_p) \rightarrow H^i(F, Z_p)$ is also nilpotent for $i = 0, 1, \dots$.*

PROOF. Consider the inverse system

Received by the editors October 11, 1958.

¹ This research was supported in part by the United States Air Force through the Air Force Office of Scientific Research of the Air Research and Development Command. under contract AF 49(638)-72 at the University of Virginia

$$X \xleftarrow{f} X \xleftarrow{f} X \xleftarrow{f} \dots$$

Let Y denote the inverse limit space whose points are the sequences $(x_i: x_i \in X \text{ and } fx_{i+1} = x_i)$. Since f^* is nilpotent, the direct limit of

$$H^i(X, Z_p) \xrightarrow{f^*} H^i(X, Z_p) \xrightarrow{f^*} \dots$$

is trivial, so that $H^i(Y, Z_p) = 0$ for $i = 0, 1, \dots$. Also $\dim Y \leq \dim X$, so that Y is finite dimensional. There is the periodic map $T': Y \rightarrow Y$ which maps (x_i) into (Tx_i) . By the Smith theorems referred to in the introduction [5], the fixed point set F' of T' is nonempty and $H^i(F', Z_p) = 0$ for $i = 0, 1, \dots$. Since F' is nonempty, then F is nonempty.

Also F' is the inverse limit space of

$$F \xleftarrow{f|F} F \xleftarrow{f|F} \dots$$

so that $H^i(F', Z_p)$ is the direct limit of

$$H^i(F, Z_p) \xrightarrow{(f|F)^*} H^i(F, Z_p) \xrightarrow{(f|F)^*} \dots$$

Since $H^i(F', Z_p) = 0$, then $(f|F)^*$ is nilpotent. The theorem follows.

We now show by example that $(f|F)^*$ is not necessarily nilpotent when coefficients are in the group Z of integers. The example is closely related to the deeper examples of §3.

(2.2) Consider the involution $T: S^2 \rightarrow S^2$ which is reflection in a great circle, and whose fixed point set F is therefore a circle. There exists a map $f: S^2 \rightarrow S^2$ with $fT = Tf$, with f homotopic to a constant, and with $f|F: S^1 \rightarrow S^1$ of degree 2.

PROOF. Consider S^2 as the join $S^0 \circ S^1$ of a 0-sphere S^0 and a circle S^1 . Let $T': S^0 \rightarrow S^0$ denote the involution interchanging the two points of S^0 . Let $T: S^2 \rightarrow S^2$ denote the join $T' \circ id$ which maps the line segment from $x_0 \in S^0$ to $x_1 \in S^1$ linearly onto the segment from $T'x_0$ to x_1 . Let $f': S^1 \rightarrow S^1$ denote an arbitrary map of degree 2. We extend f' to a map $f: S^2 \rightarrow S^2$ satisfying the assertions of our theorem. Define f so as to map the first third of the segment from $x_0 \in S^0$ to $x_1 \in S^1$ linearly onto the segment from x_0 to x_1 , the middle third linearly onto the segment from $x_1 \in S^1$ to $T'x_0 \in S^0$, and the final third linearly onto the interval from $T'x_0 \in S^0$ to $f'x_1 \in S^1$. It may be seen that $fT = Tf$ and that $f|F = f'$, which is of degree 2. It remains to be seen that f is of degree 0. We leave this to the reader, since we do not use the example further.

3. Commuting maps; the composite case. In this section we give examples to show that the results of the preceding section cannot be extended to maps whose periods are not prime power. These examples form the basis for the remainder of the paper.

(3.1) *Suppose that r is a positive integer greater than one which is not prime power. There exists a periodic map $T: S^3 \rightarrow S^3$ of the 3-sphere S^3 which is of period r and is without fixed points, and there exists a map $f: S^3 \rightarrow S^3$ with $fT = Tf$ and with f homotopic to a constant.*

PROOF. Suppose $r = p_1 p_2$, where p_1 and p_2 are relatively prime and greater than one. Let $T_i: S^1 \rightarrow S^1$ denote rotation through $2\pi/p_i$, $i = 1, 2$. Consider S^3 as the join $S^1 \circ S^1$, consisting of a union of intervals from $x_1 \in S^1$ to $x_2 \in S^1$, one joining each x_1 in the first copy of S^1 in $S^1 \circ S^1$ to each x_2 in the second copy of S^1 in $S^1 \circ S^1$. Let $T = T_1 \circ T_2$; that is, if $x = (1-t)x_1 + tx_2$ let $Tx = (1-t)T_1x_1 + tT_2x_2$. Then T is of period r , and without fixed points. Of course, T is a well-known periodic map of S^3 .

We now construct $f: S^3 \rightarrow S^3$. Since p_1 and p_2 are relatively prime, there exist integers n_1 and n_2 with $n_1 p_1 + n_2 p_2 = -1$. Consider the maps $f_i: S^1 \rightarrow S^1$ given in complex coordinates by $f_i(z) = z^{m_i}$. Then $f_i T_i = T_i f_i$ if $m_i = 1 \pmod{p_i}$. Set $m_i = n_i p_i + 1$, $i = 1, 2$. Define $f: S^3 \rightarrow S^3$ so as to map the first third of the segment from x_1 in the first copy S^1_1 of S^1 in $S^1 \circ S^1$ to x_2 in the second copy S^1_2 of S^1 in $S^1 \circ S^1$ linearly onto the segment from $f_1 x_1$ to x_2 , the middle third linearly onto the segment from $x_2 \in S^1_2$ to $x_1 \in S^1_1$, and the final third linearly onto the interval from $x_1 \in S^1_1$ to $f_2 x_2$. Explicitly

$$f((1-t)x_1 + tx_2) = \begin{cases} (1-3t)f_1x_1 + 3tx_2, & 0 \leq t \leq 1/3, \\ (3t-1)x_1 + (2-3t)x_2, & 1/3 \leq t \leq 2/3, \\ (3-3t)x_1 + (3t-2)f_2x_2, & 2/3 \leq t \leq 1. \end{cases}$$

It can be seen that $fT = Tf$. The remainder of the proof will be devoted to showing that $f: S^3 \rightarrow S^3$ is of degree 0.

If $a \in S^1$ is represented by $e^{2\pi i \alpha}$, where α is an irrational number, then $f^{-1}(a/2 + a/2)$ consists of three types of points

1. $5x_1/6 + a/6$; $f_1(x_1) = a$, $x_1 \in S^1$;
2. $a/2 + a/2$;
3. $5a/6 + x_2/6$; $f_2(x_2) = a$, $x_2 \in S^1$.

We observe that f is differentiable near each point of $f^{-1}(a/2 + a/2)$; it is also orientation preserving except at the point 2. Hence $\deg f = \deg f_1 - 1 + \deg f_2 = n_1 p_1 + n_2 p_2 + 1 = 0$. The theorem is proved.

² We are indebted to the referee for suggesting the counting argument, which is shorter than our original proof.

4. A fixed point free map on a compact acyclic space. A special case of the theorem of Smith [5] implies that if $T: X \rightarrow X$ is of prime power period where X is a compact, finite dimensional space with $H^i(X, Z) = 0$ for $i = 0, 1, \dots$, then T has a fixed point. We show that this holds *only* if the period is prime power.

(4.1) *Suppose that r is a positive integer greater than one which is not prime power. There exists a compact finite dimensional space X with $H^i(X, Z) = 0$ for $i = 0, 1, \dots$ and a map $T: X \rightarrow X$ of period r such that T has no fixed points.*

PROOF. We use the technique of the proof of (2.1). Consider the map $T: S^3 \rightarrow S^3$ of period r and the map $f: S^3 \rightarrow S^3$ constructed in (3.1). Then T is without fixed points, $fT = Tf$, and f is homotopic to a constant. Consider the inverse mapping system

$$S^3 \xleftarrow{f} S^3 \xleftarrow{f} S^3 \xleftarrow{f} \dots$$

The limit space X of this system has $H^i(X, Z) = 0$ for $i = 0, 1, \dots$, since $f^* = 0$. The map $T: S^3 \rightarrow S^3$ generates a map $T': X \rightarrow X$ which maps (x_i) into (Tx_i) . T' is of period r and without fixed points. The conclusion follows.

5. Fixed point free maps on contractible complexes. We construct in this section a star-finite, finite dimensional contractible complex X together with a periodic simplicial map $T: X \rightarrow X$ which has no fixed points. The period r of T may be any positive integer greater than one which is not prime power. The fundamental device (other than (3.1)) used in the construction is that of an iterated mapping cylinder, a concept to which we have been led by study of an example of Eilenberg [1].

The mapping cylinders that we use are possibly slightly different from the usual, due to a desire to obtain polyhedra.

(5.1) DEFINITION. Suppose X and Y are finite complexes, and that $f: X \rightarrow Y$ is simplicial. Suppose also that the vertices of X are partially ordered by an ordering $<$ under which the vertices of any simplex of X are simply ordered. It will be convenient to suppose that X and Y are embedded as disjoint subcomplexes of a simplex s . The *simplicial mapping cylinder* C of f denotes the finite complex consisting of the union of X , of Y , and of all faces of simplices spanned by $x_0, \dots, x_k, fx_k, \dots, fx_n$ (repetitions are possible) where $x_0 < \dots < x_n$ are vertices of a simplex of X and where $0 \leq k \leq n$. Call X the *beginning* of C and Y the *end* of C .

(5.2) *Suppose $f: X \rightarrow Y$ as in (5.1). The simplicial mapping cylinder C may be contracted (in C) into its end Y . If f is homotopic to a con-*

stant, then the beginning X of C may be contracted to a point in C .

PROOF. The set of vertices of C is the union of the vertices of X and the vertices of Y . Partially order the vertices of C by taking the given partial ordering on X , by simply ordering the vertices of Y , and by defining each vertex of X to be less than each vertex of Y . Triangulate $C \times I$. If $v_0 < v_1 < \dots < v_n$ span a simplex of C and $0 \leq k \leq h$, then

$$v_0 \times 0, \dots, v_k \times 0, v_k \times 1, \dots, v_n \times 1$$

span a simplex of $C \times I$. We define now the simplicial contracting map F . Define $F(v \times 0) = v$ for all vertices v of C . If v is a vertex of X , let $F(v \times 1) = fv$; if v is a vertex of Y , let $F(v \times 1) = v$. It may be checked that F extends to a simplicial map $F: C \times I \rightarrow C$, and that F contracts C into Y .

If we consider $F|X \times I$ we obtain a homotopy of the inclusion $X \subset C$ to the composition $X \rightarrow Y \subset C$ in C . If f is homotopic to a constant, then the beginning X of C may be contracted to a point in C .

(5.3) DEFINITION. Suppose that X is a finite polyhedron with triangulation Δ_1 , and that $f: X \rightarrow X$ maps an n th barycentric subdivision Δ_2 of Δ_1 simplicially into Δ_1 . Suppose $<$ is the natural partial ordering of a barycentric subdivision: $x_0 < x_1$ if the simplex of which x_0 is barycenter is a face of the simplex of which x_1 is barycenter. In the simplicial mapping cylinder C , the beginning of C occurs with triangulation Δ_2 and the end of C with triangulation Δ_1 . Let C' denote the n th barycentric subdivision of C modulo the beginning of C . Then, in C' , the beginning and end both occur with triangulation Δ_2 .

Let now C_1, C_2, \dots be a sequence of disjoint copies of C' . Let C_∞ , the iterated mapping cylinder of f , denote the infinite complex obtained from $C_1 \cup C_2 \cup \dots$ by identifying the end of C_i and the beginning of C_{i+1} . C_∞ is a star-finite, finite dimensional complex.

(5.4) If $f: X \rightarrow X$, as in (5.3), is homotopic to a constant, then C_∞ is contractible.

PROOF. Consider the subset $C_1 \cup \dots \cup C_n$ of C_∞ , with the end of C_i identified with the beginning of C_{i+1} . Repeated use of (5.2) yields a contraction, in $C_1 \cup \dots \cup C_n$, of $C_1 \cup \dots \cup C_n$ into the end of C_n . Since f is homotopic to a constant, the end of C_n is contractible to a point in C_{n+1} . Hence $C_1 \cup \dots \cup C_n$ is contractible to a point in $C_1 \cup \dots \cup C_{n+1}$. All the homotopy groups of C_∞ are therefore trivial, and C_∞ is contractible.

(5.5) THEOREM. Suppose that r is a positive integer greater than one which is not prime power. There exists a star-finite, four dimensional

contractible complex X and a simplicial map $T: X \rightarrow X$ of period r and without fixed points.

PROOF. Consider the map $T: S^3 \rightarrow S^3$ constructed in (3.1) as $T = T_1 \circ T_2$. Now T_1 and T_2 are simplicial in certain triangulations D_1 and D_2 of S^1 . There is a natural join triangulation Δ_1 of $S^1 \circ S^1$ under which T is simplicial. It is no restriction to suppose that if v is a vertex of Δ_1 for which v and $T^i v$ span an edge of Δ_1 , then $v = T^i v$.

Consider next the map $f: S^3 \rightarrow S^3$ of (3.1) with $fT = Tf$ and f homotopic to a constant. There exists an n th barycentric subdivision Δ_2 of Δ_1 such that f has a simplicial approximation from Δ_2 to Δ_1 . That is, for each vertex v of Δ_2

$$(*) \quad f(Sl_{\Delta_2} v) \subset Sl_{\Delta_1} w$$

for some vertex w of Δ_1 . Select one vertex v from each T -orbit of vertices of Δ_2 . For such a v , select any vertex w of Δ_1 , for which (*) holds and define $w = f^i v$. Operating on (*) with T^i and using $fT^i = T^i f$,

$$(**) \quad f(Sl_{\Delta_2} T^i v) \subset Sl_{\Delta_1} T^i f'(v).$$

Define $f'(T^i v) = T^i f' v$. We must show that f' is uniquely defined. Suppose $T^i v = T^j v$. Then

$$f(Sl_{\Delta_2} T^i v) \subset Sl_{\Delta_1} T^i f'(v) \cap Sl_{\Delta_1} T^j f'(v).$$

Hence $T^i f'(v)$ and $T^j f'(v)$ span an edge of Δ_1 . Hence, by the requirement imposed on Δ_1 , $T^i f'(v) = T^j f'(v)$ and f' is uniquely defined.

Hence we have obtained a simplicial approximation $f': \Delta_2 \rightarrow \Delta_1$ of f with $f'T = Tf'$. Also f' is homotopic to a constant, being homotopic to f . Let X denote the iterated mapping cylinder C_∞ of f' . Now $T: S^3 \rightarrow S^3$ is simplicial in both Δ_1 and Δ_2 . It also preserves the order of (5.3) on vertices of Δ_2 . T induces a periodic map on the simplicial mapping cylinder C of $f': X \rightarrow X$ and also of the subdivision C' of C . Hence T induces a map of period r of C_∞ which has no fixed points. The theorem follows.

6. Fixed point free maps on contractible manifolds. In this section we use a regular neighborhood technique used before [3; 4] to expand the example of (5.5) into an example on a manifold. It is plausible that the manifold constructed is E^n ; however, the J. H. C. Whitehead theory of regular neighborhoods [6] has been developed in full only for finite complexes so that we are not able to draw any immediate conclusion from it.

(6.1) THEOREM. *Suppose that r is a positive integer greater than one*

which is not prime power. There exists a contractible manifold M and a periodic map $S: M \rightarrow M$ of period r which has no fixed points.

PROOF. According to (5.5) there exists a star-finite, four dimensional contractible complex K and a simplicial map $T: K \rightarrow K$ without fixed points. It is no restriction [3, p. 431] to suppose K embedded as a subcomplex of triangulated Euclidean space E^{nr} for some n , and to suppose that T on K is $S|K$ where S is the cyclic permutation

$$(x_1, \dots, x_r) \rightarrow (x_2, \dots, x_r, x_1), \quad x_i \in E^n.$$

One may also suppose S simplicial on all of E^{nr} . Consider the regular neighborhood M of K ; that is, the union of all open stars of Sd^2K relative to Sd^2E^{nr} . Since $S(K) \subset K$ then $S(M) \subset M$. Now the fixed point set F of S is a subcomplex of E^{nr} , disjoint from K . Hence F is disjoint from M . Therefore $S: M \rightarrow M$ is of period r and without fixed points. Also K is a deformation retract of M [2, p. 70]; since K is contractible, so is M . The theorem follows.

BIBLIOGRAPHY

1. S. Eilenberg, *On a theorem of P. A. Smith concerning fixed points for periodic transformations*, Duke Math. J. vol. 6 (1940) pp. 428–437.
2. S. Eilenberg and N. Steenrod, *Foundations of algebraic topology*, Princeton, 1952.
3. E. E. Floyd, *Fixed point sets of compact abelian Lie groups of transformations*, Ann. of Math. vol. 66 (1957) pp. 30–35.
4. E. E. Floyd and R. W. Richardson, Jr., *An action of a finite group on an n -cell without stationary points*, Bull. Amer. Math. Soc. vol. 65 (1959) pp. 73–76.
5. P. A. Smith, *Fixed point theorems for periodic maps*, Amer. J. Math. vol. 63 (1941) pp. 1–8.
6. J. H. C. Whitehead, *Simplicial spaces, nuclei and m -groups*, Proc. London Math. Soc. vol. 45 (1939) pp. 243–327.

UNIVERSITY OF VIRGINIA