

APPLICATIONS OF DUALITY

EBEN MATLIS

Let R and S be two rings with units; then Cartan and Eilenberg have established the following duality identities (1) and (2) [3, Chapter 5, §5]. In the situation described by the symbol $({}_R A, {}_S B_R, {}_S C)$, where C is S -injective, we have an identity:

$$(1) \quad \text{Ext}_R(A, \text{Hom}_S(B, C)) \cong \text{Hom}_S(\text{Tor}^R(B, A), C).$$

In the situation described by the symbol $({}_R A, {}_R B_S, C_S)$, if R is left Noetherian, A is finitely R -generated, and C is S -injective, then we have a second identity:

$$(2) \quad \text{Hom}_S(\text{Ext}_R(A, B), C) \cong \text{Tor}^R(\text{Hom}_S(B, C), A).$$

In this note we shall apply these identities to the solution of two entirely dissimilar problems. First we shall prove that if R is a left Noetherian ring, then $\text{f.l.inj. dim } R = \text{f.r.w. dim } R$. Secondly, we shall give a new, short proof of a theorem of Kaplansky [5] which states that a ring R is a Prufer ring if and only if, the torsion submodule of any finitely generated R -module is a direct summand.

DEFINITION. Let R be a ring with unit. If A is a left R -module, we shall denote the injective dimension of A as an R -module by $\text{inj. dim}_R A$. If B is a right R -module, we shall denote the weak dimension of B as an R -module by $\text{w. dim}_R B$. We then define the finitistic, left, injective, dimension of R by $\text{f.l.inj. dim } R = \sup. \text{inj. dim}_R A$, where A ranges over all left R -modules of finite injective dimension. Similarly, we define the finitistic, right, weak dimension of R by $\text{f.r.w. dim } R = \sup. \text{w. dim}_R B$, where B ranges over all right R -modules of finite weak dimension.

LEMMA 1. *Let A be a left module over a ring R . Then $\text{inj. dim}_R A \leq n$ if and only if, $\text{Ext}_R^{n+1}(R/I, A) = 0$ for every left ideal I of R .*

PROOF. If $\text{inj. dim}_R A \leq n$, then of course $\text{Ext}_R^{n+1}(X, A) = 0$ for all left R -modules X . Conversely, suppose that $\text{Ext}_R^{n+1}(R/I, A) = 0$ for all left ideals I of R . Let

$$0 \rightarrow A \rightarrow C_0 \rightarrow C_1 \rightarrow \cdots \rightarrow C_{n-1} \rightarrow D \rightarrow 0$$

be an exact sequence of left R -modules, where C_i is R -injective for $i = 0, \dots, n-1$. Then $0 = \text{Ext}_R^{n+1}(R/I, A) \cong \text{Ext}_R^1(R/I, D)$ for all left

Received by the editors November 12, 1958.

ideals I of R . Hence D is R -injective [3, Proposition 1.3.2], and thus $\text{inj. dim}_R A \leq n$.

Lemma 1 provides an easy proof of the following theorem of M. Auslander [1, Theorem 1]. (It has been brought to my attention that this proof was known, but unpublished, by S. Eilenberg).

COROLLARY. *For any ring R , $\text{l.gl. dim } R = \sup. \text{hd}_R R/I$, where I ranges over the left ideals of R .*

PROOF. If $\sup. \text{hd}_R R/I = \infty$, then $\text{l.gl. dim } R = \infty$. Hence assume that $\sup. \text{hd}_R R/I = n$. Then $\text{Ext}_R^{n+1}(R/I, A) = 0$ for all left R -modules A and all left ideals I of R . Thus by Lemma 1 $\text{inj. dim}_R A \leq n$ for all left R -modules A , and thus $\text{l.gl. dim } R \leq n$.

The following lemma is well known, but we will prove it for the sake of completeness.

LEMMA 2. *Let B be a right module over a ring R . Then $\text{w. dim}_R B \leq n$ if and only if, $\text{Tor}_{n+1}(B, R/I) = 0$ for every left ideal I of R .*

PROOF. If $\text{w. dim}_R B \leq n$, then by definition $\text{Tor}_{n+1}^R(B, X) = 0$ for every left R -module X . Conversely, assume that $\text{Tor}_{n+1}^R(B, R/I) = 0$ for every left ideal I of R . We assume by induction that $\text{Tor}_{n+1}^R(B, Y) = 0$ for every left R -module Y generated by fewer than m elements. Let E be a left R -module generated by m elements, and E_1 a submodule of E generated by one of these elements. Then E/E_1 is generated by fewer than m elements. Hence from the exact sequence:

$$0 \rightarrow E_1 \rightarrow E \rightarrow E/E_1 \rightarrow 0$$

we obtain the exact sequence:

$$0 = \text{Tor}_{n+1}^R(B, E_1) \rightarrow \text{Tor}_{n+1}^R(B, E) \rightarrow \text{Tor}_{n+1}^R(B, E/E_1) = 0.$$

Thus $\text{Tor}_{n+1}^R(B, E) = 0$ for all finitely generated left R -modules E . Since Tor commutes with direct limits, $\text{Tor}_{n+1}^R(B, X) = 0$ for all left R -modules X . Thus $\text{w. dim}_R B \leq n$.

DEFINITION. Let Z denote the integers, and K the rationals (mod one). If X is a left (or right) module over a ring R , then $X^* = \text{Hom}_Z(X, K)$ is a right (or left) module over R . It is easily seen that $X = 0$, if and only if $X^* = 0$.

LEMMA 3. *Let R be any ring. Then $\text{f.r.w. dim } R \leq \text{f.l.inj. dim } R$.*

PROOF. Let B be any right R -module, and A any left R -module. Then by (1) $\text{Ext}_R(A, B^*) \cong \text{Tor}^R(B, A)^*$. Thus $\text{w. dim}_R B$

$= \text{inj. dim}_R B^*$. It follows immediately that $\text{f.r.w. dim } R \leq \text{f.l.inj. dim } R$.

THEOREM 1. *Let R be a left Noetherian ring. Then $\text{f.l.inj. dim } R = \text{f.r.w. dim } R$.*

PROOF. Let I be a left ideal of R , and A a left R -module. Then by (2) $\text{Ext}_R(R/I, A)^* \cong \text{Tor}^R(A^*, R/I)$. Hence by Lemmas 1 and 2, $\text{inj. dim}_R A = \text{w. dim}_R A^*$. Thus $\text{f.l.inj. dim } R \leq \text{f.r.w. dim } R$. The opposite inequality is provided by Lemma 3.

Of course if R is a commutative, Noetherian ring, then there is no distinction between left and right modules; and then $\text{f.inj. dim } R = \text{f.w. dim } R$. It then follows from [2, Theorem 2.4] that $\text{f.inj. dim } R \leq \text{Krull dim } R$.

As a final application of duality we give a new proof of a theorem of Kaplansky [5].

DEFINITION. A *Prufer ring* is an integral domain in which every finitely generated ideal is invertible.

THEOREM 2. *Let R be an integral domain. Then R is a Prufer ring if and only if the torsion submodule of every finitely generated R -module is a direct summand.*

PROOF. If R is a Prufer ring, a finitely generated, torsion-free R -module is projective, and the condition follows. Conversely, assume that the torsion submodule of every finitely generated R -module is a direct summand. Let S be a finitely generated, torsion-free R -module, and I an ideal of R . By [4, Theorem 2] and Lemma 2 it is sufficient to prove that $\text{Tor}_1^R(S, R/I) = 0$.

Suppose that $\text{Tor}_1^R(S, R/I) \neq 0$. Since $\text{Tor}_1^R(S, R/I)$ is a torsion module, there exists a torsion, injective R -module C such that $\text{Hom}_R(\text{Tor}_1^R(S, R/I), C) \neq 0$. By (1) we have $\text{Ext}_R^1(S, \text{Hom}_R(R/I, C)) \cong \text{Hom}_R(\text{Tor}_1^R(S, R/I), C)$. Since $\text{Hom}_R(R/I, C)$ is a torsion module, to establish a contradiction it will be sufficient to prove that $\text{Ext}_R^1(S, R) = 0$ for any torsion module T .

Let A be an extension of T by S . Now S is generated by a finite number of elements $\{y_1, \dots, y_n\}$. Choose $x_i \in A$ such that $x_i \rightarrow y_i$, and let $B = (x_1, \dots, x_n) \subset A$. Then $T \cap B$ is the torsion submodule of B . Hence by assumption $B = (T \cap B) \oplus S'$, where $S' \cong S$. Then $A = T \oplus S'$, and thus $\text{Ext}_R^1(S, T) = 0$.

BIBLIOGRAPHY

1. M. Auslander, *On the dimension of modules and algebras* (III), Nagoya Math. J. vol. 9 (1955) pp. 67-77.

2. M. Auslander and D. Buchsbaum, *Homological dimension in Noetherian rings*, II, Trans. Amer. Math. Soc. vol. 88 (1958) pp. 194–206.
3. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton University Press, 1956.
4. A. Hattori, *On Prufer rings*, J. Math. Soc. Japan vol. 9 no. 4 (1957) pp. 381–385.
5. I. Kaplansky, *A characterization of Prufer rings*, J. Indian Math. Soc. (forthcoming).

NORTHWESTERN UNIVERSITY