

A GENERAL RUDIN-CARLESON THEOREM

ERRETT BISHOP¹

1. **Introduction.** Rudin [7] and Carleson [4] have independently proved that if S is a closed set of Lebesgue measure 0 on the unit circle

$$L = \{z: |z| = 1\}$$

and if f is a continuous function on L then there exists a continuous function F on

$$D = \{z: |z| \leq 1\}$$

which is analytic on $D-L$ such that $F(z)=f(z)$ for all z in S . It is the purpose of this paper to generalize this theorem. Before stating the generalization, we remark that the Rudin-Carleson theorem is closely related to a theorem of F. and M. Riesz, which states that any (finite, complex-valued, Baire) measure on L which is orthogonal to all continuous functions F on D which are analytic on $D-L$ is absolutely continuous with respect to Lebesgue measure $d\theta$ on L . The proofs of the two theorems show that the results of Rudin-Carleson and of F. and M. Riesz are closely related. We shall state an abstract theorem which shows that the Rudin-Carleson theorem is a direct consequence of the F. and M. Riesz theorem. This abstract theorem will permit an automatic generalization of the Rudin-Carleson result to any situation to which the F. and M. Riesz result can be generalized. The theorem to be proved reads as follows:

THEOREM 1. *Let $C(X)$ be the uniformly-normed Banach space of all continuous complex-valued functions on a compact Hausdorff space X . Let B be a closed subspace of $C(X)$. Let B^\perp consist of all (finite, complex-valued, Baire) measures μ on X such that $\int f d\mu = 0$ for all f in B . Let $\hat{\mu}$ be the regular Borel extension of the Baire measure μ . Let S be a closed subset of X with the property that $\hat{\mu}(T) = 0$ for every Borel subset T of S and every μ in B^\perp . Let f be a continuous complex-valued function on S and Δ a positive function on X such that $|f(x)| < \Delta(x)$ for all x in S . Then there exists F in B with $|F(x)| < \Delta(x)$ for all x in X and $F(x) = f(x)$ for all x in S .*

If X is taken to be the set L defined above and B is taken to be those functions in $C(L)$ which are restrictions to L of functions in

Received by the editors December 13, 1960.

¹ This work was supported by the Sloan Foundation.

$C(D)$ analytic on $D-L$, then by the F. and M. Riesz result any closed subset S of L of Lebesgue measure 0 satisfies the hypotheses of Theorem 1.

Therefore from Theorem 1 the Rudin-Carleson theorem follows.

By a general F. and M. Riesz theorem we mean a theorem to the effect that for certain closed subspaces B of certain $C(X)$ there exists a non-negative measure μ_0 on X with respect to which all measures in B^\perp are absolutely continuous. From Theorem 1 it follows that to each such general F. and M. Riesz theorem corresponds a general Rudin-Carleson theorem, which states that if S is a closed subset of X with $\mu_0(S) = 0$ then every continuous function on S is the restriction to S of some function in B .

There are various general F. and M. Riesz theorems which exist in the literature. We mention three of these.

(1) Bochner [3] and Helson and Lowdenslager [5] have proved an F. and M. Riesz theorem for $X = L \times L$, where B is the subspace of $L \times L$ generated by all functions $z^m w^n$ with (m, n) belonging to a sector of lattice points of opening greater than π . Here μ_0 is Lebesgue measure on $L \times L$.

(2) Bishop [1; 2] has proved a general F. and M. Riesz theorem for the boundary X of a compact set C in the complex plane whose complement is connected. Here B consists of all continuous functions on X which have extensions to C analytic on $C-X$.

(3) Wermer [9] and Royden [6] (see also Rudin [8]) have proved a general F. and M. Riesz theorem for X the boundary of a finite Riemann surface R . Here B again is the set of continuous functions on X which can be extended to be analytic on R and continuous on $X \cup R$.

2. Proof of Theorem 1. We first prove a lemma from which the proof of Theorem 1 will be trivial.

LEMMA. *Assume that $|f(x)| < r < 1$ for all x in S . Then under the hypotheses of Theorem 1 there exists F in B with $\|F\| < 1$ and $F(x) = f(x)$ for all x in S .*

PROOF. Let U_r be the set of all g in B of norm less than r . Let ϕ be the restriction mapping of B into $C(S)$. The lemma states that $f \in \phi(U_1)$. We begin by showing that $f \in V_r$, where V_r is the closure of $\phi(U_r)$. Assume otherwise. It follows from one of the many variants of the Hahn-Banach theorem that there exists a bounded linear functional α on $C(S)$ with $\alpha(f) > 1$ and $|\alpha(h)| < 1$ for all h in V_r . By the Riesz representation theorem, there exists a measure μ_1 on S such that

$$\int g d\mu_1 = \alpha(g)$$

for all g in $C(S)$. Define the linear functional β on B by $\beta(g) = \alpha(\phi(g))$. Since $\phi(g) \in V_r$ for all g in U_r it follows that $|\beta(g)| < 1$ for all g in U_r so that $\|\beta\| \leq r^{-1}$. By the Hahn-Banach theorem and the Riesz representation theorem it follows that there exists a measure μ_2 on X with $\|\mu_2\| \leq r^{-1}$ and $\beta(g) = \int g d\mu_2$ for all g in B . Thus the measure

$$\mu = \mu_1 - \mu_2$$

is in B^\perp . Also

$$\left| \int_S f d\mu \right| \geq \int_S f d\mu_1 - r \|\mu_2\| > 1 - r r^{-1} = 0,$$

contradicting the fact that μ vanishes on all subsets of S . This contradiction shows that $f \in V_r$, so that there exists F_1 in B with $\|F_1\| < r$ and $|f(x) - F_1(x)| < \lambda/2$ for all x in S , where $\lambda = 1 - r$. If we write $f_1(x) = f(x) - F_1(x)$, it follows by the result just proved, with f replaced by f_1 , that there exists F_2 in B with $\|F_2\| < \lambda/2$ and $|f_1(x) - F_2(x)| < \lambda/4$ for all x in S . Thus by induction we find a sequence $\{F_n\}$ of functions in B with

$$\|F_n\| < 2^{-n+1}\lambda, \quad n \geq 2,$$

and

$$\left| f(x) - \sum_{k=1}^n F_k(x) \right| < 2^{-n}\lambda, \quad n \geq 2, x \in S.$$

We define the function F in $C(X)$ by

$$F = \sum_{n=1}^{\infty} F_n.$$

Thus $F \in B$ and

$$\|F\| < r + \sum_{n=2}^{\infty} 2^{-n+1}\lambda = r + \lambda = 1.$$

Clearly also $F(x) = f(x)$ for all x in S . This proves the lemma.

Proof of Theorem 1. Let B_0 be the closed subspace

$$\{g: \Delta g \in B\}$$

of $C(X)$. Then

$$B_0^\perp = \{ \Delta\mu : \mu \in B^\perp \}.$$

Thus $\nu(T) = 0$ for all Borel sets $T \subset S$ and all ν in B_0^\perp . It follows from the lemma that there exists F_0 in B_0 with $\|F_0\| < 1$ and $F_0(x) = (\Delta(x))^{-1}f(x)$ for all x in S . Let F be the element ΔF_0 of B . Thus $|F(x)| < |\Delta(x)|$ for all x in X and $F(x) = \Delta(x)F_0(x) = f(x)$ for all x in S . This completes the proof.

REFERENCES

1. Errett Bishop, *The structure of certain measures*, Duke Math. J. vol. 25 (1958) pp. 283-290.
2. ———, *Boundary measures of analytic differentials*, Duke Math. J. vol. 27 (1960) pp. 331-340.
3. Salomon Bochner, *Boundary values of analytic functions in several variables and of almost periodic functions*, Ann. of Math. vol. 45 (1944) pp. 708-722.
4. Lennart Carleson, *Representations of continuous functions*, Math. Z. vol. 66 (1957) pp. 447-451.
5. Henry Helson and David Lowdenslager, *Prediction theory and Fourier series in several variables*, Acta Math. vol. 99 (1958) pp. 165-202.
6. Halsey Royden, *The boundary values of analytic and harmonic functions*, O.O.R. Technical Report no. 19, DA-04-200-ORD-704, Stanford University.
7. Walter Rudin, *Boundary values of continuous analytic functions*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 808-811.
8. ———, *Analyticity and the maximum modulus principle*, Duke Math. J. vol. 20 (1953) pp. 449-458.
9. John Wermer, *Subalgebras of the algebra of all complex-valued continuous functions on the circle*, Amer. J. Math. vol. 78 (1956) pp. 225-242.

UNIVERSITY OF CALIFORNIA, BERKELEY