

## ON 3-MANIFOLDS THAT ARE NOT SIMPLY CONNECTED

KYUNG WHAN KWUN

Let  $X$  be a connected 3-manifold. Then  $X$  is simply connected if and only if each simple closed curve in  $X$  lies in a simply connected neighborhood. On the other hand, each point of  $X$  trivially lies in a simply connected neighborhood with or without  $X$  itself simply connected. However, we will show that arcs are big enough to reflect the global property of  $X$  as to simple connectedness in the sense that  $X$  is simply connected if and only if each arc in  $X$  lies in a simply connected neighborhood.

**THEOREM.** *Let  $X$  be a 3-manifold that is not simply connected. Then there exists an arc in  $X$  that does not lie in any simply connected neighborhood.*

**PROOF.** There exists a simple closed polygon  $L$  that is not deformable to a point over  $X$ , i.e., the inclusion map of  $L$  into  $X$  is essential. To see this, we choose a loop representing a nontrivial element of the fundamental group of  $X$  and apply thereto the simplicial approximation theorem followed by obvious modifications if necessary. Let  $T$  be a polyhedral solid torus which is a closed tubular neighborhood of  $L$ . Evidently,  $T$  does not lie in a simply connected neighborhood.

We now proceed to construct a Cantor set of Antoine type [1]. Let  $T_1, T_2, \dots, T_k$  be a finite number of polyhedral solid tori of diameter less than  $1/2$  that are situated in  $\text{Int } T$  as suggested by Figure 1. We show that the sum  $T^1$  of the  $T_i$  does not lie in a simply connected neighborhood. Suppose it does.




FIGURE 1




FIGURE 2

Received by the editors March 16, 1961.

Let  $U$  be a simply connected open set containing  $T^1$ . We cut a polyhedral cube  $R$  containing  $T_1$  and  $T_2$  out of  $T$  as suggested by Figures 1 and 2.  $T_1$  and  $T_2$  are linked in  $R$  in the sense that neither of the two can be deformed to a point over the complement in  $R$  of the other. We show that they can be joined by an arc in  $\text{Int } R \cdot U$ . Let  $L_i$  be simple closed polygon that circle on  $\text{Bd } T_i$  longitudinally once and  $x_i$  be points on  $L_i$ ,  $i=1, 2$ . Since  $U$  is simply connected, there exists a singular disk  $D$  bounded by  $L_1$ ,  $D \subset U$ . We may suppose that  $D$  meets  $\text{Bd } R$  at a 1-dimensional set. Let  $D'$  be the component of  $D$ .  $\text{Int } R$  that contains  $L_1$ . Then  $D' + \text{Bd } R$  contains a singular disk  $D''$  bounded by  $L_1$ . Since  $D''$  meets  $L_2$  and  $L_2$  does not meet  $\text{Bd } R$ , the existence of the required arc is established.

Let  $C_1$  be an arc in  $\text{Int } R \cdot U$  that joins  $x_1$  and  $x_2$ . Similarly, we construct arcs  $C_i$  joining  $x_i$  and  $x_{i+1}$ ,  $x_i \in T_i$ , subscripts being taken modulo  $k$ , such that  $\sum C_i$  is a simple closed curve in  $U$  which is deformable to  $L$  in  $\text{Int } T$ . Since  $U$  is simply connected,  $L$  is deformable to a point over  $X$ , contradictory to the choice of  $L$ . Thus  $T^1$  does not lie in a simply connected neighborhood.

We now construct polyhedral solid tori  $T_{i1}, T_{i2}, \dots$  in  $\text{Int } T_i$  such that each  $T_{ij}$  is of diameter less than  $1/4$ ,  $T_{ij}$  are situated in  $\text{Int } T_i$  as  $T_j$  are in  $\text{Int } T$ . Let  $T^2$  be the sum of  $T_{ij}$ . Then  $T^2$  does not lie in a simply connected neighborhood. For if it did, then the simply connected neighborhood must contain for each  $i$  a simple closed polygon in  $\text{Int } T_i$  that is deformable within  $T_i$  to a simple closed polygon that lies on  $\text{Bd } T_i$  and circles  $T_i$  longitudinally once, and which in turn would imply that it contained a curve which is deformable to  $L$  in  $X$ , thus the same contradiction.

Similarly, we construct sets  $T^3, T^4, \dots$  such that no  $T^i$  lies in a simply connected neighborhood, each  $T^i$  contains  $T^{i+1}$  in its interior and the intersection of  $T^i$  is a Cantor set  $C$ . There exists in  $\text{Int } T$  an arc  $P$  containing  $C$ . Then  $P$  does not lie in any simply connected neighborhood in  $X$ .

We wish to thank the referee for his suggestion which made the proof considerably shorter.

#### REFERENCE

1. L. Antoine, *Sur l'homotopie de deux figures et de leur voisinages*, J. Math. Pures Appl. **86** (1921), 221-325.

THE SEOUL NATIONAL UNIVERSITY, SEOUL, KOREA