

REPRODUCING KERNELS AND PRINCIPAL FUNCTIONS¹

BURTON RODIN²

1. Introduction. Using the notation and terminology of Ahlfors-Sario [3], we shall denote by Γ_h the Hilbert space of square integrable harmonic differentials on a Riemann surface W . Let γ be a 1-chain on W and let ψ_γ be the element in Γ_h with the property, $(\omega, \psi_\gamma) = \int_\gamma \omega$ for all $\omega \in \Gamma_h$. We refer to ψ_γ as a reproducing kernel for periods for Γ_h . Let ζ be a point of W and $z = x + iy$ be a local parameter near ζ such that $z(\zeta) = 0$. A Γ_h -kernel for n th derivatives at ζ is a differential $\psi_\zeta \in \Gamma_h$ which satisfies $(\omega, \psi_\zeta) = (\partial^n / \partial x^n)u(0)$ for all $\omega \in \Gamma_h$ where $\omega = du(z)$ near ζ . This kernel is uniquely determined by ζ , the uniformizer z , and the positive integer n . If in the above definitions we replace Γ_h by one of its subspaces then we shall refer to the corresponding kernels as reproducing kernels for that subspace. It is easily seen that such a kernel is the orthogonal projection of a Γ_h -kernel.

The existence of these kernels is well known (see [1; 3]). The possibility of expressing them explicitly in terms of principal harmonic functions (the existence of which has been proved constructively by Sario [5]) was first investigated by Weill [6]. In this paper the kernels for the spaces $\Gamma_h, \Gamma_{hse}, \Gamma_{he}, \Gamma_{hm}, \Gamma_{h0}, \Gamma_{he} \cap \Gamma_{hse}^*, \Gamma_{h0} \cap \Gamma_{hse}^*, \Gamma_a, \Gamma_{ase}$, and some spaces associated with a regular partition of the ideal boundary of the surface are found in terms of principal functions.

§§9 and 10 deal with some applications of these results. A characterization of Ahlfors' class of quasi-rational functions is given in terms of principal functions. A Riemann-Roch type theorem is proved for arbitrary Riemann surfaces which reduces to a theorem of Royden [4] in case the surface is of class O_{KD} .

2. Reproducing kernels. We shall apply the general method of normal operators [3] to the following situation. Let h be a harmonic function defined in a boundary neighborhood of a Riemann surface W' and assume that h has zero flux along the ideal boundary of W' . The principal operator L_0 maps h into a function p_{0h} which is harmonic on W' and is characterized up to an additive constant by the

Received by the editors October 24, 1961.

¹ Work supported by the N.S.F., the O.O.R. contract number DA-04-495-ORD-722, and the A.F.O.S.R. contract number AF49(638)-574.

² Parts of this paper are contained in a doctoral dissertation submitted to the University of California, Los Angeles. The author is indebted to Professor Leo Sario for his helpful guidance.

property that the normal derivative of $p_{0h} - h$ vanishes identically along the ideal boundary of W' . When W' is the complement of a finite subset $\{\zeta_1, \dots, \zeta_n\}$ of a Riemann surface W then $p_{0h} - h$ has a removable singularity at each ζ_i . We say that it has L_0 behavior at the ideal boundary of W . Let P be a regular partition of this ideal boundary [3, p. 165]. The $(P)L_1$ principal operator associates to h a function p_{Ph} , harmonic on W' with the singularity h at each ζ_i . Furthermore, $p_{Ph} - h$ is constant on each ideal boundary component of W determined by P and has zero flux over each P -dividing cycle.

Let I and Q denote, respectively, the identity and the canonical partitions of the ideal boundary of W in the sense of [3]. In terms of a uniformizer z near $\zeta \in W$ such that $z(\zeta) = 0$, we let $s = \text{Re } z^{-n}$ and $t = \text{Im } z^{-n}$ for a positive integer n , and define $s \equiv t \equiv 0$ outside of a compact subset of W . The differential $dp_{I_s} + dp_{I_t}^*$ has a removable singularity at ζ and is an element of Γ_h .

THEOREM 1. *The differential $dp_{I_s} + dp_{I_t}^*$ reproduces for the space Γ_h . Explicitly, we have*

$$(\omega, dp_{I_s} + dp_{I_t}^*) = - \frac{2\pi}{(n - 1)!} \frac{\partial^n}{\partial x^n} u(0)$$

where $\omega = du(z)$ near ζ .

3. Before proving the theorem we shall first establish a lemma. A subregion $\Omega \subset W$ is called canonical if it is relatively compact, its boundary consists of a finite number of analytic Jordan curves, and each component of $W - \Omega$ is noncompact and bounded by a single contour. For a first order differential ω let $\int_{\beta\omega}$ indicate the limit as $\Omega \rightarrow W$ of integrals taken along the boundaries of exhausting canonical subregions Ω .

LEMMA. *Let p be the $(I)L_1$ principal function for any singularity. Then $\int_{\beta} p\omega = 0$ for all $\omega \in \Gamma_h$.*

If $\Omega \subset W$ is a canonical subregion with boundary $\partial\Omega$ then $|\int_{\partial\Omega} p\omega| = |\int_{\partial\Omega} (p - p_\Omega)\omega|$, where p_Ω is the $(I)L_1$ principal function for the surface Ω which has the same singularity as that of p . By Stokes' theorem the right side of this equation is the inner product $(dp - dp_\Omega, \bar{\omega}^*)_\Omega$ taken over the region Ω . By Schwarz's inequality this inner product is dominated by $\|dp - dp_\Omega\|_\Omega \cdot \|\omega\|_\Omega$. We therefore obtain $|\int_{\beta} p\omega| \leq \lim_{\Omega \rightarrow W} \|dp - dp_\Omega\|_\Omega \cdot \|\omega\|_W$. The uniform convergence $p_\Omega \rightarrow p$ on compact subsets of W implies that $\|dp - dp_\Omega\|_\Omega \rightarrow 0$.

For the proof of Theorem 1 we may assume that ω is real. Let Δ be a relatively compact neighborhood of ζ which is mapped by z onto a

disk $\{|z| < r\}$ and let $\partial\Delta = \alpha$. In order to simplify the notation we shall temporarily denote the functions p_{I_s} and p_{I_t} by p and q respectively. In Δ let $\omega = du(z)$ and $p = \operatorname{Re} z^{-n} + v(z)$ where u and v are harmonic in $\{|z| < r\}$. We then obtain

$$\begin{aligned} (\omega, dp + dq^*) &= \iint_W \omega(dp^* - dq) \\ &= \iint_{W-\Delta} \omega dp^* - \omega dq + \iint_{\Delta} \omega(dp^* - dq). \end{aligned}$$

The integral over $W - \Delta$ is the limit as $\Omega \rightarrow W$ of integrals over $\Omega - \Delta$. We may apply Stokes' theorem to these integrals and, after simplification, we find

$$(\omega, dp + dq^*) = \int_{\beta} (p\omega^* + q\omega) - \int_{\alpha} (p\omega^* + p^*\omega).$$

The integral along β vanishes by the lemma. Hence

$$\begin{aligned} (\omega, dp + dq^*) &= - \int_{|z|=r} (\operatorname{Re} z^{-n} + v)du^* + (\operatorname{Im} z^{-n} + v^*)du \\ &= \operatorname{Re} \int (u^* - iu)d(z^{-n}) - \int vdu^* - u dv^* \\ &= \operatorname{Re} ni \int \frac{u + iu^*}{z^{n+1}} dz \\ &= \frac{2\pi}{(n-1)!} \frac{\partial^n}{\partial x^n} u(0). \end{aligned}$$

4. We now seek an expression for the reproducing differential for periods. Let Δ be a parametric disk on W and γ a 1-simplex contained in Δ . In terms of the parameter z which maps Δ onto the unit disk we define a singularity function $\sigma = \log |(z - \zeta_2)/(z - \zeta_1)|$, where $\partial\gamma = \zeta_2 - \zeta_1$. We set $\sigma \equiv 0$ outside of a compact set. The corresponding $(P)L_1$ principal function is $p_{P\sigma}$. Let τ be the singularity function $\tau = \arg (z - \zeta_2)/(z - \zeta_1)$ in $\Delta - \gamma$ and $\tau \equiv 0$ near the ideal boundary of W . On the surface $W - \gamma$ we choose the normal operator which is composed of $(P)L_1$ for a boundary neighborhood of W and of the Dirichlet operator for $\Delta - \gamma$. This Dirichlet operator maps a continuous function on $\partial\Delta$ into the restriction to $\Delta - \gamma$ of the harmonic function in Δ with these boundary values. The direct sum of these operators yields a function $p_{P\tau}$ harmonic on $W - \gamma$. The differential $dp_{P\tau}$ can be ex-

tended harmonically to all of $W - \{\zeta_1, \zeta_2\}$. We shall continue to denote the extension by dp_{Pr} , even though it is not exact. If γ is an arbitrary 1-chain it is homologous to a finite sum $\sum n_i \gamma_i$ where each γ_i is a 1-simplex contained in a parametric disk and each n_i is an integer. We extend the definitions of $dp_{P\sigma}$ and dp_{Pr} to arbitrary γ by letting $dp_{P\sigma} = \sum n_i dp_{P\sigma_i}$ and similarly for dp_{Pr} . That these differentials are well defined will follow from Theorem 2.

THEOREM 2. *The differential $dp_{I\sigma} + dp_{I\tau}^*$ reproduces for Γ_h . Specifically, if σ and τ correspond to a 1-chain γ then*

$$(\omega, dp_{I\sigma} + dp_{I\tau}^*) = 2\pi \int_{\gamma} \omega$$

for all $\omega \in \Gamma_h$.

Because of the linearity it suffices to prove Theorem 2 for the case that γ is a 1-simplex contained in a parametric disk $\Delta: \{|z| < 1\}$ and $\partial\gamma = \zeta_2 - \zeta_1$. We shall shorten the notation and write p and q for $p_{I\sigma}$ and $p_{I\tau}$ respectively. As in the proof of Theorem 1 it can be shown that

$$(\omega, dp + dq^*) = - \int_{\alpha} (p\omega^* + p^*\omega),$$

where $\partial\Delta = \alpha$. In Δ let $\omega = du$ and $p = \log |(z - \zeta_2)/(z - \zeta_1)| + v$, where u and v are harmonic. Let α_1 and α_2 be disjoint circles in Δ with centers at $z(\zeta_1)$ and $z(\zeta_2)$ respectively. After applying Green's formula one obtains

$$(\omega, dp + dq^*) = \int_{\alpha_2} ud \log |z - \zeta_2|^* - \int_{\alpha_1} ud \log |z - \zeta_1|^*.$$

By the mean value formula the last two integrals reduce to $2\pi(u(\zeta_2) - u(\zeta_1))$. We have shown that $(\omega, dp + dq^*) = 2\pi \int_{\zeta_1}^{\zeta_2} du = 2\pi \int_{\alpha} \omega$. This completes the proof of Theorem 2.

5. For a regular partition P of the ideal boundary of W the spaces $(P)\Gamma_{hm}$ and $(P)\Gamma_{hse}$ were introduced in [3], and a proof of the decomposition $\Gamma_h = (P)\Gamma_{hm} \dot{+} (P)\Gamma_{hse}^*$ was sketched. It follows that $\Gamma_h = \Gamma_{he}^* \dot{+} (P)\Gamma_{hm} \dot{+} (P)\Gamma_{hse}^* \cap \Gamma_{h0}$ is an orthogonal direct sum. Consider the identity

$$(1) \quad -(dp_{I\sigma} + dp_{I\tau}^*) = (dp_{0\tau} - dp_{I\tau})^* + (dp_{P\sigma} - dp_{I\sigma}) - (dp_{0\tau}^* + dp_{P\sigma}),$$

where the zero subscript refers to the L_0 principal function. The first term on the right side of (1) is evidently in Γ_{he}^* . The second term is

the limit as $\Omega \rightarrow W$ of $dp_{P_s\Omega} - dp_{I_s\Omega}$, where the subscript Ω indicates a principal function for the subsurface Ω . These approximating differentials belong to the space $(P)\Gamma_{hm}(\Omega)$ of harmonic differentials du on Ω with u constant on each set of contours of the boundary of Ω which belong to the same part in the induced partition. We may conclude that the limit differential is in $(P)\Gamma_{hm}$. The last term on the right side of (1) is a limit of elements from $\Gamma_{h0}(\Omega)$. By Theorem V.14C of [3], we see that it is in Γ_{h0} . Its conjugate has vanishing periods along any cycle which is dividing relative to the partition P . Hence it is in $(P)\Gamma_{hse}^* \cap \Gamma_{h0}$. Thus (1) represents the reproducing kernel for Γ_h , except for a multiplicative constant, as the sum of the kernels for Γ_{he}^* , $(P)\Gamma_{hm}$, and $(P)\Gamma_{hse}^* \cap \Gamma_{h0}$. The reasoning is also valid for the kernels corresponding to a 1-chain γ . This proves the following two corollaries.

COROLLARY 1. *The differentials*

$$\frac{(n-1)!}{2\pi} (dp_{P_s} - dp_{I_s})$$

and

$$\frac{1}{2\pi} (dp_{I_\sigma} - dp_{P_\sigma})$$

are the reproducing kernels for the space $(P)\Gamma_{hm}$.

COROLLARY 2. *The differentials*

$$\frac{-(n-1)!}{2\pi} (dp_{P_s} + dp_{0_s}^*)$$

and

$$\frac{1}{2\pi} (dp_{P_\sigma} + dp_{0_\sigma}^*)$$

are the reproducing kernels for $(P)\Gamma_{hse}^* \cap \Gamma_{h0}$.

Since $(I)\Gamma_{hse}^* \cap \Gamma_{h0} = \Gamma_{h0}$, Corollary 2 contains the following special case.

COROLLARY 3. *The differentials*

$$\frac{-(n-1)!}{2\pi} (dp_{I_s} + dp_{0_s}^*)$$

and

$$\frac{1}{2\pi} (d\phi_{I\sigma} + d\phi_{0\tau}^*)$$

are the reproducing kernels for Γ_{h_0} .

REMARKS. In order to indicate explicitly the relation between the differentials in (1) and the classical domain functions we denote by $N(z, \zeta)$ and $G(z, \zeta)$ the Neumanns' and Green's functions for W . In case W is parabolic let $G(z, \zeta) = G(z, z_0; \zeta, \zeta_0)$ be the fundamental potential [4]. In terms of the complex notation

$$d' = \frac{\partial}{\partial z} dz, \quad d'' = \frac{\partial}{\partial \bar{z}} d\bar{z}, \quad d^c = i(d'' - d')$$

(so that $d^c f = df^*$), we have

$$\begin{aligned} d\phi_{I\sigma} + d\phi_{I\tau}^* &= \frac{4}{(n-1)!} \operatorname{Re} d'' \frac{\partial^n G}{\partial \zeta^n}, \\ (d\phi_{0\sigma} - d\phi_{I\tau})^* &= \frac{2}{(n-1)!} \operatorname{Im} d^c \frac{\partial^n}{\partial \zeta^n} (N - G), \\ d\phi_{I\sigma} + d\phi_{I\tau}^* &= 4 \operatorname{Re} \int_{\zeta \in \gamma} d_z'' d_\zeta' G. \end{aligned}$$

Theorems 1 and 2 can also be derived from the fact that $d''(\partial G/\partial \zeta)$ reproduces for Γ_a . If W is the interior of a compact bordered surface then by classical theory there is a potential $G_P(z, z_0; \zeta, \zeta_0)$ which as a function of z is constant and has zero flux along each P -boundary component. In this case we have

$$\begin{aligned} d\phi_{P\sigma} - d\phi_{I\sigma} &= \frac{2}{(n-1)!} \operatorname{Re} d \frac{\partial^n}{\partial \zeta^n} (G_P - G), \\ d\phi_{0\sigma}^* + d\phi_{P\sigma} &= \frac{2}{(n-1)!} \operatorname{Re} (dG_P - id^c N). \end{aligned}$$

6. The space Γ_h has the orthogonal decomposition

$$\Gamma_h = (P)\Gamma_{hm} \dot{+} \Gamma_{h_0}^* \dot{+} (P)\Gamma_{h\sigma\epsilon}^* \cap \Gamma_{h\epsilon}.$$

Since we are in possession of reproducing kernels for $(P)\Gamma_{hm}$ and $\Gamma_{h_0}^*$ (see the proof of Corollary 1) we obtain the next result immediately.

COROLLARY 4. *The differentials*

$$\frac{(n-1)!}{2\pi} (d\phi_{0\sigma} - d\phi_{P\sigma})$$

and

$$\frac{1}{2\pi} (dp_{P\sigma} - dp_{0\sigma})$$

are the reproducing kernels for $(P)\Gamma_{h\sigma}^* \cap \Gamma_{h\sigma}$.

The reproducing kernel for $\Gamma_{h\sigma}$ may be derived from Corollary 4.

COROLLARY 5. *The differentials*

$$\frac{(n-1)!}{2\pi} (dp_{0\sigma} - dp_{I\sigma})$$

and

$$\frac{1}{2\pi} (dp_{I\sigma} - dp_{0\sigma})$$

are the reproducing kernels for $\Gamma_{h\sigma}$.

7. The kernel for $(P)\Gamma_{h\sigma}$ can be found from the identity $-(dp_{I\sigma} + dp_{I\sigma}^*) = (dp_{P\tau}^* - dp_{I\sigma}^*) - (dp_{I\sigma} + dp_{P\tau}^*)$ and the orthogonal decomposition $\Gamma_h = (P)\Gamma_{hm}^* \dot{+} (P)\Gamma_{h\sigma}$.

COROLLARY 6. *The differentials*

$$\frac{-(n-1)!}{2\pi} (dp_{I\sigma} + dp_{P\tau}^*)$$

and

$$\frac{1}{2\pi} (dp_{I\sigma} + dp_{P\tau}^*)$$

are the reproducing kernels for $(P)\Gamma_{h\sigma}$.

8. From the orthogonal direct sum,

$$\Gamma_h = (P)\Gamma_{hm} \dot{+} (P)\Gamma_{hm}^* + (P)\Gamma_{a\sigma\epsilon} + (P)\bar{\Gamma}_{a\sigma\epsilon},$$

we can project the Γ_h -kernels into $(P)\Gamma_{a\sigma\epsilon}$.

COROLLARY 7. *The differentials*

$$\frac{-(n-1)!}{4\pi} (dp_{P\sigma} + dp_{P\tau}^* + i(dp_{P\sigma}^* - dp_{P\tau}))$$

and

$$\frac{1}{4\pi} (dp_{P\sigma} + dp_{P\tau}^* + i(dp_{P\sigma}^* - dp_{P\tau}))$$

are the reproducing kernels for $(P)\Gamma_{ase}$.

The kernels for Γ_a and Γ_{ase} may be found immediately from Corollary 7.

REMARKS. It is an open question whether results analogous to those above can be obtained for the subspaces Γ_S and Γ_{aS} . Investigations in this area might lead to new normal operators.

9. The results of §§1-8 indicate a connection between principal functions and Ahlfors' generalization of Abel's theorem [2, 3]. Recall that a differential ω on W , harmonic except for harmonic poles, is called *distinguished* if

- (i) ω^* is semiexact outside of some compact subset of W ,
- (ii) there exist differentials $\omega_{hm} \in \Gamma_{hm}$ and $\omega_{e0} \in \Gamma_{e0}^1$ such that $\omega = \omega_{hm} + \omega_{e0}$ in a boundary neighborhood of W .

LEMMA. Let γ be a 1-chain on W and δ be a 1-cycle. Let $dp_{P\sigma}$ and $dp_{P\tau}$ be differentials associated with γ (see §4). Then

$$(2) \quad \int_{\delta} (dp_{P\sigma} + idp_{P\tau}) = 2\pi i(\delta \times \gamma).$$

PROOF. Because (2) is linear in γ we may assume that γ is a simplex contained in a parametric disk $\Delta: \{|z| < 1\}$. Since $dp_{P\sigma} + idp_{P\tau}$ has no periods along cycles in $W - \Delta$ we may even assume $\delta \subset \bar{\Delta}$. Let $\partial\gamma = \zeta_2 - \zeta_1$. Then $dp_{P\sigma} + idp_{P\tau} = d \log (z - \zeta_2)/(z - \zeta_1) + du$ in Δ , where u is harmonic. The proof reduces to establishing

$$(3) \quad \int_{\delta} d \log \frac{z - \zeta_2}{z - \zeta_1} = 2\pi i(\delta \times \gamma).$$

Since $\delta \times \gamma$ is the number of times γ crosses δ from left to right, (3) follows from the argument principle.

With the help of the above lemma it can be seen that to each distinguished differential ω there corresponds a differential $\lambda(P, \omega)$ with the singularities and periods of ω and which, in a boundary neighborhood of W , is the differential of a function whose real and imaginary parts have $(P)L_1$ -behavior.

THEOREM 3. Let ω be a distinguished differential. Then $\lambda(Q, \omega) = \omega$.

PROOF. Because of the uniqueness theorem for distinguished differentials it suffices to prove merely that $\lambda(Q, \omega)$ satisfies conditions (i) and (ii) above. The first condition is obvious.

The differential $\omega_{hm} = \lambda(Q, \omega) - \lambda(I, \omega)$ is in Γ_{hm} . Let Ω be a canonical subregion of W such that ω is regular and exact in $W - \Omega$. Choose another canonical region $\Omega' \supset \bar{\Omega}$. By standard techniques one can construct an exact differential ω_{e0} in Γ^1 with the property,

$$\omega_{e0} = \begin{cases} 0 & \text{in } \Omega, \\ \lambda(I, \omega) & \text{in } W - \Omega'. \end{cases}$$

That $\omega_{e0} \in \Gamma_{e0}$ follows from the fact that an $(I)L_1$ principal function for W is the limit, uniformly on compacta, of $(I)L_1$ principal functions for bordered exhausting canonical subregions. The equality $\lambda(Q, \omega) = \omega_{hm} + \omega_{e0}$ is valid in $W - \Omega'$ and hence $\lambda(Q, \omega)$ is distinguished.

From the above considerations one sees that *a meromorphic function f is quasi-rational if and only if $\log f$ has a single-valued branch outside of a compact subset of W , the real and imaginary parts of which are $(Q)L_1$ principal functions.*

REMARK. There is a natural interpretation for "quasi-rational with respect to a regular partition P ." The corresponding generalized Abel's theorem is obtained by replacing Γ_{ase} by $(P)\Gamma_{ase}$.

10. Riemann-Roch type theorems. The classical theorem of Riemann-Roch has been extended to surfaces of class O_{KD} by Royden [4]. By the methods of this paper we obtain a similar theorem for an arbitrary Riemann surface.

Let D be a divisor on a Riemann surface W and let $D = B - A$ where A and B are disjoint integral divisors. Suppose that $A = \sum m_j A_j$ and $B = \sum n_k B_k$, where the A_j and B_k are points of W and the m_j and n_k are positive integers. For each j and k let Δ_j and Δ'_k be parametric disks with centers at A_j and B_k respectively. We assume that these disks are mutually disjoint. Let $\Delta = \cup \Delta_j$ and $\Delta' = \cup \Delta'_k$.

Let $S = s + it$ be a meromorphic function in Δ' and a multiple of the divisor $-B$. Define $S \equiv 0$ in a boundary neighborhood of W and form the differential $dF_S = dp_{P_s} + idp_{P_t}$. Let \mathfrak{B} denote the complex vector space consisting of all such dF_S . The dimension of \mathfrak{B} is equal to the degree of B , i.e., $\dim \mathfrak{B} = \deg B = \sum n_k$.

In the decomposition $dF_S = \phi_S + \psi_S$, where $\psi_S = \frac{1}{2}(dF_S - idF_S^*)$, the differential ϕ_S is meromorphic and ψ_S is in $(P)\Gamma_{ase}$. According to Corollary 7, ψ_S has certain reproducing properties. The following lemma is a generalization of that fact. For a divisor E let $(P)\Gamma_{ase}[E]$ denote the vector space of meromorphic differentials which are mul-

tuples of E , square integrable near the ideal boundary of W , and which have vanishing periods along each P -dividing cycle outside of some compact set.

LEMMA. Let $\alpha \in (P)\Gamma_{ase}[-A]$. Then

$$(4) \quad (\alpha, \psi_S)_{W-\Delta} = i \int_{\partial\Delta'} S\alpha + i \int_{\partial\Delta} F_S\alpha.$$

The inner product is to be understood as a Cauchy limit. That is, it is the limit of inner products taken over $W-\Delta-\Delta'$ as the radii of the parametric disks of Δ' tend to zero. Note that

$$(\alpha, \psi_S)_{W-\Delta-\Delta'} = (dF_S, \bar{\alpha})_{W-\Delta-\Delta'} = -i \lim_{\Omega \rightarrow W} \int_{\partial(\Omega-\Delta-\Delta')} F_S\alpha.$$

A modification of the proof of the lemma in §3 shows that $\int_{\partial\Omega} F_S\alpha \rightarrow 0$ as $\Omega \rightarrow W$. In Δ' we have $F_S = S + u$ where S is meromorphic and u is harmonic. When the disks about the B_k shrink to points we obtain (4).

We shall make use of the following algebraic facts [3, p. 325]. Let U and V be vector spaces over a field K . A bilinear mapping $T: U \times V \rightarrow K$ is called a pairing of U and V . The left kernel U_0 is the space $\{u \in U: T(u, V) = 0\}$, and the right kernel V_0 is the space $\{v \in V: T(U, v) = 0\}$. If one of the quotient spaces U/U_0 or V/V_0 is finite dimensional then there is an isomorphism $U/U_0 \cong V/V_0$.

Consider the pairing $T: \mathfrak{B} \times (P)\Gamma_{ase}[-A] \rightarrow \mathfrak{C}$ defined by $T(dF_S, \alpha) = \int_{\partial\Delta'} S\alpha$.

Suppose dF_S is in the left kernel of this pairing. From (4) we have $(\alpha, \psi_S)_{W-\Delta} = i \int_{\partial\Delta} F_S\alpha$ for all $\alpha \in (P)\Gamma_{ase}[-A]$. We may replace α by ψ_S and conclude that $\|\psi_S\|_W = 0$. Hence $\psi_S = 0$ and dF_S is meromorphic. We also find that

$$(5) \quad \int_{\partial\Delta} F_S\alpha = 0.$$

For appropriate choices of α we conclude from (5) that the additive constant in F_S can be chosen so that F_S is a multiple of the divisor $-D$. Conversely, if dF_S is meromorphic then $\psi_S = 0$ and $T(dF_S, \alpha) = \int_{\partial\Delta'} S\alpha = -\int_{\partial\Delta} F_S\alpha$. The differential α has a pole of order at most m_j at A_j , and $F_S^{(k)}(A_j) = 0$ ($k = 1, \dots, m_j - 1$) if F_S is a multiple of A . Hence $\int_{\partial\Delta} F_S\alpha = 0$ by Cauchy's integral formula. Thus dF_S is in the left kernel \mathfrak{B}_0 if and only if the function F_S is meromorphic and can be normalized so as to be a multiple of $-D$.

A differential $\alpha \in (P)\Gamma_{ase}[-A]$ is in the right kernel if and only if

$\int_{\partial\Delta} S\alpha = 0$ for all S which are multiples of $-B$. Convenient choices for S show that α must be a multiple of B , i.e., $\alpha \in (P)\Gamma_{ase}[D]$.

Since \mathfrak{B} is finite dimensional we have

$$\mathfrak{B}/\mathfrak{B}_0 \cong (P)\Gamma_{ase}[-A]/(P)\Gamma_{ase}[D],$$

or

$$(6) \quad \dim \mathfrak{B}_0 = \dim \mathfrak{B} - \dim (P)\Gamma_{ase}[-A]/(P)\Gamma_{ase}[D].$$

Let $(P)M$ be the complex vector space of meromorphic functions on W whose real and imaginary parts have $(P)L_1$ behavior near the ideal boundary. Denote by $(P)M[E]$, where E is a divisor, the subspace of $(P)M$ consisting of functions which are multiples of E . The homomorphism $d: (P)M[-D] \rightarrow \mathfrak{B}_0: F_S \rightarrow dF_S$ is surjective. The kernel is \mathbf{C} if $\deg A = 0$, and it is zero if $\deg A \neq 0$. From (6) we obtain

$$(7) \quad \begin{aligned} \dim (P)M[-D] &= \deg B + 1 - \min(1, \deg A) \\ &\quad - \dim (P)\Gamma_{ase}[-A]/(P)\Gamma_{ase}[D]. \end{aligned}$$

THEOREM 4. *Let A and B be disjoint integral divisors on a Riemann surface W and let $D = B - A$. Then (7) is valid for any regular partition P of the ideal boundary of W .*

BIBLIOGRAPHY

1. L. V. Ahlfors, *The method of orthogonal decomposition for differentials on open Riemann surfaces*, Ann. Acad. Sci. Fenn. Ser. A. I. No. 249/7 (1958).
2. ———, *Abel's theorem for open Riemann surfaces*, Seminars on analytic functions. II, 7–19. Institute for Advanced Study, Princeton, N. J., 1958.
3. L. V. Ahlfors and L. Sario, *Riemann surfaces*, Princeton Univ. Press, Princeton, N. J., 1960.
4. H. L. Royden, *The Riemann-Roch theorem*, Comment. Math. Helv. 34 (1960), 52–66.
5. L. Sario, *A linear operator method on arbitrary Riemann surfaces*, Trans. Amer. Math. Soc. 72 (1952), 281–295.
6. G. Weill, *Reproducing kernels and orthogonal kernels for analytic differentials on Riemann surfaces*, Pacific J. Math. (to appear).

HARVARD UNIVERSITY