

# SOME APPLICATIONS OF THE INEQUALITY OF ARITHMETIC AND GEOMETRIC MEANS TO POLYNOMIAL EQUATIONS

HERBERT S. WILF

The purpose of this note is to point out a simple generalization of the inequality

$$(z_1 z_2 \cdots z_n)^{1/n} \leq \frac{1}{n} (z_1 + \cdots + z_n)$$

of arithmetic and geometric means, which will hold when the arguments of the complex numbers  $z_1, \cdots, z_n$  are suitably restricted. We shall apply the resulting inequality to the roots of polynomial equations, obtaining first a quantitative form of the Gauss-Lucas theorem, and then some relationships between the coefficients of a polynomial and the size of a sector containing its roots.

**1. The inequality.** The basic result is

**THEOREM 1.** *Suppose*

$$|\arg z_i| \leq \psi < \frac{\pi}{2}, \quad i = 1, 2, \cdots, n.$$

*Then*

$$(1) \quad |z_1 z_2 \cdots z_n|^{1/n} < (\sec \psi) \frac{1}{n} |z_1 + z_2 + \cdots + z_n|$$

*unless  $n$  is even and  $z_1 = \cdots = z_{n/2} = \bar{z}_{(n/2)+1} = \cdots = \bar{z}_n = r e^{i\psi}$ , in which case equality holds.*

**PROOF.** We have

$$\begin{aligned} |z_1 + z_2 + \cdots + z_n| &\geq |\operatorname{Re}(z_1 + \cdots + z_n)| \\ &= (|z_1| \cos \phi_1 + |z_2| \cos \phi_2 + \cdots \\ &\quad + |z_n| \cos \phi_n) \\ (2) \quad &\geq (\cos \psi)(|z_1| + \cdots + |z_n|) \\ &\geq n \cos \psi (|z_1| |z_2| \cdots |z_n|)^{1/n} \end{aligned}$$

---

Received by the editors March 2, 1962.

as claimed. All signs of equality hold only when

$$(3) \quad \begin{aligned} (a) \quad & \operatorname{Im} (z_1 + \cdots + z_n) = 0 \\ (b) \quad & \cos \phi_i = \cos \psi \quad (i = 1, 2, \cdots, n) \\ (c) \quad & |z_1| = |z_2| = \cdots = |z_n| \end{aligned}$$

which imply the configuration stated in the theorem. For odd  $n$  the constant  $\sec \psi$  is only asymptotically best possible.

## 2. Application to polynomials. Let

$$(4) \quad P(z) = a_0 + a_1 z + \cdots + a_n z^n = a_n (z - z_1) \cdots (z - z_n)$$

be given and let  $K$  denote the convex hull of the zeros  $z_1, \cdots, z_n$  of  $P(z)$ . Let  $z$  be outside  $K$ , and suppose that, from  $z$ ,  $K$  subtends an angle  $2\psi$ . Then the spread in the arguments of the numbers

$$\frac{1}{z - z_1}, \cdots, \frac{1}{z - z_n}$$

is at most  $2\psi$ , and from Theorem 1,

$$\left| \frac{1}{(z - z_1)} \cdots \frac{1}{(z - z_n)} \right|^{1/n} \leq (\sec \psi) \frac{1}{n} \left| \sum_{v=1}^n \frac{1}{z - z_v} \right|.$$

But this is just the assertion that

$$\left| \frac{a_n}{P(z)} \right|^{1/n} \leq \frac{\sec \psi}{n} \left| \frac{P'(z)}{P(z)} \right|,$$

and we have proved

**THEOREM 2.** *If  $z$  is a point from which the convex hull of the zeros of the polynomial  $P(z)$  of degree  $n$  subtends an angle  $2\psi < \pi$ , then*

$$(5) \quad |P'(z)| \geq n |a_n|^{1/n} (\cos \psi) |P(z)|^{1-(1/n)}.$$

**COROLLARY 1.** *The zeros of  $P'(z)$  lie in the convex hull of the zeros of  $P(z)$  (Gauss-Lucas).*

**COROLLARY 2.** *If the zeros of  $P(z)$  lie in the unit circle, then we have for  $|z| > 1$ ,*

$$(6) \quad |P'(z)| \geq \frac{n |a_n|^{1/n}}{\sqrt{\left(1 - \frac{1}{|z|^2}\right)}} |P(z)|^{1-(1/n)}.$$

THEOREM 3. *The zeros of the polynomial*

$$P(z) = a_0 + a_1z + \cdots + a_nz^n,$$

*are not contained in any sector of central angle less than*

$$2 \cos^{-1} \left\{ \left| \frac{a_{n-1}}{na_n} \right| \min_{0 \leq k \leq n-1} \left| \frac{a_n}{a_k} \binom{n}{k} \right|^{1/n-k} \right\}.$$

PROOF. Suppose the zeros of  $P(z)$  lie in a sector of angle  $2\psi < \pi$ . From Theorem 1,

$$\left| \frac{a_0}{a_n} \right|^{1/n} \cong \frac{\sec \psi}{n} \left| \frac{a_{n-1}}{a_n} \right|,$$

or

$$\sec \psi \cong n \left| a_n \right|^{1-(1/n)} \left| a_0 \right|^{1/n} \left| a_{n-1} \right|^{-1}.$$

Applying this result to

$$P^{(k)}(z) = \sum_{\nu=0}^{n-k} \frac{(\nu+k)!}{\nu!} a_{\nu+k} z^\nu,$$

which, by Corollary 1 also satisfies the hypotheses, we find

$$\sec \psi \cong \left| \frac{na_n}{a_{n-1}} \right| \left| \frac{a_n}{a_k} \binom{n}{k} \right|^{-1/(n-k)} \quad (k = 0, 1, \dots, n-1),$$

and the result follows.

THEOREM 4. *Under the hypotheses of Theorem 2, let  $\rho$  denote the distance from  $z$  to the center of gravity of the zeros of  $P(z)$ . Then*

$$(7) \quad |P(z)| \leq |a_n| (\rho \sec \psi)^n.$$

PROOF. Apply Theorem 1 to the numbers  $z - z_1, \dots, z - z_n$ .

UNIVERSITY OF ILLINOIS