

Recall that, by Theorem 3.1, $E(\mathfrak{Q}) \supset \Delta(A)$. From our definitions $\Delta(A)$ is dense in \mathfrak{B} so that also $E(\mathfrak{Q}) \supset \mathfrak{B}$. Since the reverse inequality is clear, the proof is complete.

REFERENCES

1. I. M. Gel'fand, D. A. Raikov and G. E. Šilov, *Commutative normed rings*, Uspehi Mat. Nauk **1** (1946), 48–146; Amer. Math. Soc. Transl. (2) **5** (1957), 115–220.
2. C. E. Rickart, *General theory of Banach algebras*, Van Nostrand, New York, 1960.
3. G. E. Šilov, *On the extension of maximal ideals*, Dokl. Akad. Nauk SSSR **29** (1940), 83–84.

UNIVERSITY OF OREGON

SEMI-HOMOGENEOUS FUNCTIONS

LOUIS V. QUINTAS AND FRED SUPNICK

1. Introduction and statement of results. A function f is called *homogeneous of degree n with respect to the set A* , or briefly *semi-homogeneous* if

$$(1.0) \quad f(ax) = a^n f(x)$$

is satisfied for all x in the domain of f and all a in A .

With each admissible A there is associated a class of solutions of (1.0). E.g., let R denote the set of all real numbers and let f be a function on R to R . If A consists only of the irrationals, then $f(x) = cx^n$ ($c = f(1)$) is the unique solution of (1.0). On the other hand, if A consists only of the rationals, then in addition to $f(x) = cx^n$, (1.0) has other solutions (e.g., if n is any nonzero integer and $f(x) = x^n$ or 0 accordingly as x is rational or irrational).

We are interested in studying decompositions of the set of admissible A 's into classes and in characterizing the solutions of (1.0) corresponding to these classes. In this paper we show how this can be done in a natural way for semi-homogeneous functions of a real variable. We note that in this case our methods apply to

$$(1.1) \quad f(ax) = p(a)f(x) \quad (a \in A \subset R),$$

where p is a product-preserving function on R to R (cf. [1]). We shall therefore confine our attention to (1.1).

Received by the editors March 8, 1962.

Let $C(p, A)$ denote the set of all functions on R to R satisfying (1.1). Let $R^* = R - \{0\}$ and A^* be the multiplicative subgroup of R^* generated by $A - \{0\}$. It is easy to see that if A contains a set of generators for R^* , then $f(x) = f(1)p(x)$. Thus, the problem of determining which subsets of R lead to the solution $f(x) = f(1)p(x)$ of (1.1) is equivalent to classifying those subsets of R which generate R^* (cf. §5). With respect to the general case we first establish the following lemma.

LEMMA. *If $p \neq 0$ and $p \neq 1$ on A^* , then $C(p, A) = C(p, A^*)$ (cf. §2).*

Thus we are led to consider the problem: *For an arbitrary subgroup A^* of R^* , how can the members of $C(p, A^*)$ be characterized?*

The general answer to the above question is given in Theorem 1. Let R^*/A^* denote the multiplicative quotient group of R^* modulo A^* , B a set of representatives for R^*/A^* , and p a product-preserving function on R to R such that $p \neq 0$ and $p \neq 1$ on A^* (cf. [2]).

THEOREM 1. (i) $f \in C(p, A^*)$ if and only if

$$(1.2) \quad f(x) = \begin{cases} F(b)p(x) & x = ab, a \in A^*, b \in B, \\ 0 & x = 0, \end{cases}$$

where F is a function on B to R .

(ii) *If $f \in C(p, A^*)$, then f is uniquely determined when it is known on B (cf. §2).*

An explicit classification and description of the solutions of (1.1) depends on a consideration of the different possibilities for A^* and p . We note that A^* is either dense in the positive reals, dense in R , or is discrete and countable. Also, if p is continuous on R^* ($p \neq 0$), then p is of the form $p(x) = |x|^k$, or $p(x) \in \{x^k (0 < x), -|x|^k (x < 0)\}$ where k is a real constant.

THEOREM 2. *Let $A^* \subset R^*$, $p \neq 0, 1$ on A^* , and $f \in C(p, A^*)$. If p is continuous on R and A^* is dense in R , then f is of type*

$T_1: f(x) = cp(x)$, " c " a constant,

$T_2: f$ is continuous only at $x=0$, and the closure $\text{Cl}(G(f))$ of its (cartesian) graph $G(f)$ (in E_2) is a set of curves $\{y = cp(x)\}$ with $\{c\}$ bounded, or

$T_3: f$ is totally discontinuous and $\text{Cl}(G(f))$ is a set of curves $\{y = cp(x)\}$ with $\{c\}$ unbounded. $\text{Cl}(G(f))$ may be the entire plane if f is of type T_3 (cf. §3).

REMARK. It is of interest to note that if f is a function which satis-

fies $f(x+y)=f(x)+f(y)$, then the classic description of f given by G. Hamil (namely, that f is of the form $f(x)=cx$, or is totally discontinuous and the graph of f is dense in the entire plane) is contained within our Theorem 2 (cf. [3]). This follows from the fact that $f(x+y)=f(x)+f(y)$ implies f is homogeneous of degree 1 with respect to the rationals, i.e., $f(ax)=af(x)$ for all rational numbers a .

Let A^* be an infinite discrete subgroup of R^* . Let g denote the smallest member of A^* such that $1 < g$, $G_0 \equiv \{(x, f(x)): 1 \leq x < g\}$ where f is a function on R to R , D_a the transformation of E_2 into E_2 defined by $D_a(x, y) = (ax, p(a)y)$ ($a \in R$), and $D_a G_0$ the image of G_0 under D_a .

THEOREM 3. *If A^* is an infinite discrete subgroup of R^* containing at least one negative member and $f \in C(p, A^*)$, then $\{D_a G_0: a \in A^* \cup \{0\}\}$ is a decomposition of $G(f)$ (cf. [4] and §4).*

In Theorems 2 and 3 we have considered the cases (i) p continuous and A^* dense, and (ii) p arbitrary and A^* discrete. We note that if p is discontinuous on R^* , then p is totally discontinuous and the graph of p is dense in the first and second or the first and third quadrants depending on whether $p(-1) = 1$ or -1 , respectively. The problem of obtaining an esthetic classification of the solutions of (1.1) for the case p discontinuous and A^* dense in R remains open.

2. Proof of the Lemma and Theorem 1.

PROOF OF THE LEMMA. Since $A - \{0\} \subset A^*$, $C(p, A^*) \subset C(p, A - \{0\})$. If $0 \notin A$, then $C(p, A^*) \subset C(p, A)$. If $0 \in A$, then since $p(0) = 0$ every member of $C(p, A)$ must assign 0 to 0. But $p \neq 1$ on A^* implies there exist $a, a' \in A^*$ such that $p(a) \neq p(a')$. Thus, $f(0)(p(a) - p(a')) = 0$ yields $f(0) = 0$ whenever $f \in C(p, A^*)$. Therefore, $C(p, A^*) \subset C(p, A)$.

Let $f \in C(p, A)$. To prove $C(p, A) \subset C(p, A^*)$ we use the fact that every member of A^* is a finite product of members of $A - \{0\}$ or their reciprocals. Thus, if $s \in A^*$, then $s = at$ or $s = a^{-1}t$ with $a \in A - \{0\}$ and $t \in A^*$. If $s = at$, then

$$(2.1) \quad f(sx) = f(atx) = p(a)f(tx).$$

If $s = a^{-1}t$, then

$$(2.2) \quad f(sx) = p(a^{-1}a)f(a^{-1}tx) = p(a^{-1})f(tx).$$

Applying operations of the form (2.1) or (2.2) successively we obtain $f(sx) = p(s)f(x)$ ($s \in A^*$). Thus, $C(p, A) = C(p, A^*)$.

PROOF OF THEOREM 1 (i). Since R^*/A^* defines a partition of R^* ,

each nonzero real number x has a unique representation in terms of A^* and B . Namely, $x = ab$ ($a \in A^*$, $b \in B$). Thus (1.2) defines a function on R to R .

Let $f \in C(p, A^*)$. Since $p \neq 0$, then $p(b) \neq 0$ ($b \in R^*$). Thus $f(x) = p(x)(p(b))^{-1}f(b)$ whenever $x = ab$ ($a \in A^*$, $b \in B$). Since $p \neq 1$ on A^* , then $f(0) = 0$. Consequently, f can be expressed in the form (1.2) with F the function on B to R defined by $F(b) = (p(b))^{-1}f(b)$.

Conversely, let f be defined by (1.2). If $s \in A^*$ and $x \in R^*$, then $x = ab$ and $sx = sab$ ($a \in A^*$, $sa \in A^*$, $b \in B$). Thus,

$$f(sx) = F(b)p(sx) = p(s)F(b)p(x) = p(s)f(x).$$

If $x = 0$, then $f(s0) = 0 = p(s)f(0)$. Therefore, $f \in C(p, A^*)$.

PROOF OF THEOREM 1 (ii). If $f, f' \in C(p, A^*)$ and $f(b) = f'(b)$ ($b \in B$), then

$$f(x) = p(a)f(b) = p(a)f'(b) = f'(x)$$

whenever $x = ab$ ($a \in A^*$, $b \in B$), and $f(0) = 0 = f'(0)$.

3. Proof of Theorem 2. By Theorem 1 the image under f of each coset of R^*/A^* lies on a curve, $y = cp(x)$. Specifically, if $x \in A^*b$, then $(x, f(x))$ lies on $y = F(b)p(x)$. Since each coset of R^*/A^* is dense in R , the image I of each coset is dense in the curve containing I . Each $y = cp(x)$ is continuous, hence the closure of the graph of f is a set of curves $K \equiv \{y = cp(x)\}$ through the origin. If all the cosets of R^*/A^* are mapped onto the same curve, then f is a function of type T_1 . If this is not the case, then f is a function of type T_2 or T_3 accordingly as the set of coefficients $C \equiv \{c\}$ appearing in K is bounded or unbounded. If K contains at least two curves, it is clear that f cannot be continuous at nonzero values of x . Suppose C is bounded. If $c_1 = \text{lub } C$, $c_2 = \text{glb } C$, and $c_0 = \max\{|c_1|, |c_2|\}$, then $f(x)$ lies between $-c_0p(x)$ and $c_0p(x)$ for all x . If $\{x_i\}$ is any sequence of real numbers converging to 0, then $\lim_{x_i \rightarrow 0} \{c_0p(x_i)\} = 0$ implies $\lim_{x_i \rightarrow 0} \{f(x_i)\} = 0$. Since $f(0) = 0$, we have f is continuous at $x = 0$. Hence, f is of type T_2 . If C is unbounded, then f is unbounded in every neighborhood of the origin. Since $x = 0$ is the only possible point of continuity and $f(0) = 0$, f is totally discontinuous and is thus of type T_3 .

If the index of A^* in R^* is 2^{\aleph_0} and B is a set of representatives for R^*/A^* , then there exists a 1-1 function F on B to R . The function f defined by $f(x) = F(b)p(x)$ ($x = ab$, $a \in A^*$, $b \in B$) and $f(0) = 0$ is a type T_3 member of $C(p, A^*)$ and $G(f)$ is dense in $K \equiv \{y = F(b)p(x) : b \in B\}$. But $K \equiv \{y = rp(x) : r \in R\}$ is the entire plane with the exception of the points on the y -axis having nonzero ordinates. Hence $\text{Cl}(G(f))$ is the entire plane. This completes the proof of Theorem 2.

4. Proof of Theorem 3. Since A^* is discrete and not finite there exists a smallest member $g \in A^*$ such that $1 < g$ and $B = \{x \in R: 1 \leq x < g\}$ is a set of representatives for R^*/A^* .

A^* induces the following decomposition of R^* . The set of positive reals is decomposed into intervals $g^n \leq x < g^{n+1}$ where n ranges over the integers. A^* also contains a largest member h such that $h \leq -1$. We now decompose the set of negative reals into the sets $hg^{n+1} < x \leq hg^n$. In this way R^* is decomposed into intervals each of which is the image of the set B under multiplication by a suitable member of A^* . This decomposition of R^* induces a decomposition of $G(f)$ into the following sets:

$G_n = \{(x, f(x)): g^n \leq x < g^{n+1}\}$, $G'_n = \{(x, f(x)): hg^{n+1} < x \leq hg^n\}$, and $\{(0, 0)\}$, where g and h are as described above and n ranges over the integers.

We note that G_0 is the graph of f in E_2 which corresponds to B . Since f is completely determined when we know its values on B , we seek a relation between the set G_0 and the other sets of this decomposition of $G(f)$. Applying D_{g^n} to G_0 we obtain

$$\begin{aligned} \{(g^n x, p(g^n)f(x)): 1 \leq x < g\} &= \{(g^n x, f(g^n x)): 1 \leq x < g\} \\ &= \{(y, f(y)): g^n \leq y < g^{n+1}\}. \end{aligned}$$

Therefore, $D_{g^n}G_0 = G_n$. Similarly, $D_{hg^n}G_0 = G'_n$ and $D_0G_0 = \{(0, 0)\}$.

$G(f)$ thus decomposes into sets each of which is the image of the set G_0 under a transformation D_a . Therefore, $\{D_a G_0: a \in A^* \cup \{0\}\}$ is a decomposition of $G(f)$. This completes the proof of Theorem 3.

Theorems analogous to Theorems 2 and 3 may be obtained when A^* is a subgroup of the positive reals or when p is continuous only on R^* .

5. Remarks on the solution $f(1)p(x)$. Let $A \subset R$ and μ denote the Lebesgue measure on R .

1. If $\mu(A) > 0$ and A contains at least one negative member, then $f(1)p(x)$ is the unique solution of (1.1).

PROOF. Let R_p^* denote the positive reals. If $A_p = A^* \cap R_p^*$, then $\mu(A_p) > 0$. Since $\mu(\ln[A_p]) > 0$, we have

$$D(\ln[A_p]) \equiv \{x - y: x = \ln(a), y = \ln(b), a, b \in A_p\}$$

contains an open interval containing the origin (cf. [5]). Thus,

$$e^{D(\ln[A_p])} = \{e^{\ln(a) - \ln(b)} = a/b: a, b \in A_p\}$$

contains an open interval containing 1. Since every positive real

number has an n th root in this interval, $e^{D(\ln[A_p])}$ generates R_p^* . Hence, $R_p^* \subset A^*$. Since A contains a negative number $-1 \in A^*$. Therefore, $A^* = R^*$ and the assertion follows from the Lemma.

2. There exist sets $A \subset R$ such that $\mu(A) = 0$ and $f(1)p(x)$ is the unique solution of (1.1)

PROOF. Let C denote the Cantor set and $A = \{e^c: c \in C\} \cup \{-1\}$. It is a theorem of Steinhaus that for every number d between 0 and 1 there are two numbers of C which differ by d (cf. [6]). Thus, $\{e^c: c \in C\}$ generates R_p^* . Hence, $A^* = R^*$.

3. It is clear that if A is countable then $A^* \neq R^*$. We also note that R^* does not contain any minimal set of generators.

NOTES AND REFERENCES

1. The special case $p(a) = a$ ($a \in A \subset R$) was first considered by the authors and the results announced in Abstract 577-8, Notices Amer. Math. Soc. 8 (1961), 51.

2. If $0 \in A$ and $p \equiv 1$, then $C(1, A)$ is the set of all constant functions. If $0 \notin A$, A is not null, and $p \equiv 1$ on A^* , then $C(1, A)$ is the set of all functions which are constant on the cosets of R^*/A^* and $f(0)$ is an arbitrary constant.

3. G. Hamil, *Eine Basis aller Zahlen und die unstetige Lösungen der Funktionalgleichung $f(x+y) = f(x) + f(y)$* , Math. Ann. 60 (1905), 459-462.

4. By a *decomposition* of a set X we mean a disjoint family of subsets of X whose union is X .

5. P. R. Halmos, *Measure theory*, Van Nostrand, Princeton, N. J., 1950, Theorem B, p. 68.

6. H. Steinhaus, *A new property of G. Cantor's set*, Wektor 7 (1917). (Polish) See also, J. F. Randolph, *Distances between points of the Cantor set*, Amer. Math. Monthly 47 (1940), 549.

CITY COLLEGE