

## WEAKLY COMPACT $B^\#$ -ALGEBRAS

A. OLUBUMMO

1. A complex Banach algebra  $A$  is a compact (weakly compact) algebra if its left and right regular representations consist of compact (weakly compact) operators. Let  $E$  be any subset of  $A$  and denote by  $E_l$  and  $E_r$  the left and right annihilators of  $E$ .  $A$  is an annihilator algebra if  $A_l = (0) = A_r$ ,  $I_r \neq (0)$  for each proper closed left ideal  $I$  and  $J_l \neq (0)$  for each proper closed right ideal  $J$ .

In [6, Theorem 1], it was shown that a semi-simple compact algebra is an annihilator algebra. The first main result of the present paper (Theorem 2.1) is that a semi-simple annihilator algebra is a weakly compact algebra. Thus if  $\mathfrak{C}$ ,  $\mathfrak{A}$ ,  $\mathfrak{W}$  denote respectively the class of all semi-simple compact algebras, all semi-simple annihilator algebras and all weakly compact algebras, we have  $\mathfrak{C} \subset \mathfrak{A} \subset \mathfrak{W}$ .

§3 is devoted to the structure theory of weakly compact  $B^\#$ -algebras begun in [7]. A Banach algebra  $A$  is a  $B^\#$ -algebra if, given  $a \in A$ , there exists  $a^\# \neq 0$  in  $A$  such that

$$\|a^\#\| \|a\| = \|(a^\#a)^n\|^{1/n}, \quad n = 1, 2, 3, \dots$$

In their study of weakly compact  $B^*$ -algebras Ogasawara and Yoshinaga [4] obtained the following structure theorem:

**THEOREM.** *The following statements are equivalent for a  $B^*$ -algebra  $A$ :*

- (1)  $A$  is a weakly compact algebra;
- (2)  $A$  is the  $B^*(\infty)$ -sum of  $C^*$ -algebras, each of which consists of the set of all compact operators on a Hilbert space.

The following result was obtained in [7, Theorem 3.1]:

**THEOREM.** *A Banach algebra  $A$  is the algebra  $F(X)$  of all uniform limits of operators of finite rank on a reflexive Banach space  $X$  if and only if  $A$  is a simple, weakly compact  $B^\#$ -algebra with minimal ideals.*

Making use of this result and our present Theorem 2.1, we now obtain the following more general result:

**THEOREM 3.4.** *The following statements are equivalent:*

- (1)  $A$  is a weakly compact  $B^\#$ -algebra with a dense socle;
- (2)  $A$  is the  $B(\infty)$ -sum of  $B^\#$ -algebras, each of which is the algebra of all uniform limits of operators of finite rank on a reflexive Banach space.

---

Received by the editors September 4, 1962.

We note that every  $B^*$ -algebra is a  $B^\sharp$ -algebra and that a weakly compact  $B^*$ -algebra automatically has a dense socle [4, p. 15], so that Theorem 3.4 includes the result of Ogasawara and Yoshinaga.

2. **THEOREM 2.1.** *A semi-simple annihilator algebra  $A$  is a weakly compact algebra.*

**PROOF.** Let  $Ae$ ,  $e^2=e$ , be a minimal left ideal of  $A$ . Then  $Ae$  is a reflexive Banach space since it is also a minimal left ideal of the simple annihilator algebra  $(AeA)^-$  [1, Theorem 13]. Let  $a \in A$ ; then [3, p. 483, Corollary 3] right multiplication by  $ae$  is a weakly compact mapping of  $A$  into  $Ae$ , and a fortiori, of  $A$  into  $A$ . Then by [3, p. 484, Theorem 5], right multiplication by a socle element is weakly compact. Since the socle is dense [1, Theorem 4], it follows [3, p. 483, Corollary 4] that any  $x \in A$  is a right (and similarly left) weakly compact element.

3. In this section we prove a structure theorem for weakly compact  $B^\sharp$ -algebras.

**LEMMA 3.1.** *Let  $A$  be a semi-simple Banach algebra with a dense socle. Then every maximal regular left ideal  $M$  of  $A$  has a nonzero right annihilator.*

**PROOF.** If  $\{Ae_\alpha\}_{\alpha \in \Omega}$  denotes the set of all the minimal left ideals of  $A$ , then there exists  $\alpha_0 \in \Omega$  such that  $Ae_{\alpha_0} \not\subset M$ . Further,  $M \cap Ae_{\alpha_0} = (0)$  and  $M \oplus Ae_{\alpha_0} = A$ . Since  $M$  is a regular left ideal of  $A$ , there exists  $j \in A$  such that  $xj - x \in M$  for every  $x \in A$ . For some  $a_0 \in A$  and  $m_0 \in M$ , we have  $j = m_0 + a_0e_{\alpha_0}$ ,  $a_0e_{\alpha_0} \neq 0$ . Let  $m$  be an arbitrary element of  $M$ ; then  $mj - m \in M$  and  $mj - ma_0e_{\alpha_0} = mm_0 \in M$ , from which it follows that  $m - ma_0 \cdot e_{\alpha_0} \in M$ , and therefore  $ma_0 \cdot e_{\alpha_0} \in M$ . However,  $ma_0 \cdot e_{\alpha_0} \in Ae_{\alpha_0}$  since  $Ae_{\alpha_0}$  is a left ideal; thus  $ma_0 \cdot e_{\alpha_0} \in M \cap Ae_{\alpha_0} = (0)$ , and since  $m$  is arbitrary in  $M$ , the lemma is proved.

**LEMMA 3.2.** *Let  $A$  be a  $B^\sharp$ -algebra with a dense socle. If  $|\cdot|$  is any norm in  $A$  with  $|a| \leq \|a\|$  for each  $a \in A$ , then  $|\cdot| = \|\cdot\|$ .*

**PROOF.** Suppose that  $j \in A$  and  $j$  has no left reverse. We show that there exists  $a \neq 0$  such that  $ja = a$ . In fact, let  $J = [yj - y : y \in A]$ ; then  $J$  is a regular left ideal of  $A$  which is proper since  $j \notin J$ . Now  $J$  is contained in a maximal regular left ideal  $M$  and by Lemma 3.1, there exists  $a \in A$ ,  $a \neq 0$  such that  $Ja = (0)$ , i.e. such that  $yja - ya = 0$  for all  $y \in A$ ; i.e.,  $A(ja - a) = (0)$ . Since  $A$ , being a  $B^\sharp$ -algebra is semi-simple,  $A_r = (0)$  from which it follows that  $ja = a$ . The conclusion now follows exactly as in [2, Theorems 3 and 4].

**LEMMA 3.3.** *A semi-simple Banach algebra  $A$  with a dense socle (or with the annihilator property) is the completion of the direct join of all its minimal closed two-sided ideals.*

This is essentially Theorem 6 of Bonsall and Goldie [1], under the hypothesis that  $A$  be an annihilator algebra. The annihilator property implies that  $A$  has a dense socle and this, together with the semi-simplicity of  $A$ , is all that is required to prove the theorem.

**DEFINITION.** Let  $\{A_\alpha\}_{\alpha \in \Omega}$  denote a set of Banach algebras. The  $B(\infty)$ -sum of the  $A_\alpha$  is the Banach algebra  $A$  consisting of all the functions  $f(\cdot)$  defined on  $\Omega$  with  $f(\alpha) \in A_\alpha$  for each  $\alpha \in \Omega$  and such that, given  $\epsilon > 0$ , there is a finite subset  $\alpha_1, \alpha_2, \dots, \alpha_n$  of  $\Omega$  such that  $\|f(\alpha)\| < \epsilon$  for  $\alpha \neq \alpha_1, \alpha_2, \dots, \alpha_n$ . We define the algebraic operations in  $A$  in the obvious way, e.g.  $(f+g)(\alpha) = f(\alpha) + g(\alpha)$ , etc. and define the norm by  $\|f(\cdot)\| = \sup_{\alpha \in \Omega} \|f(\alpha)\|$ .

We now state our second main result:

**THEOREM 3.4.** *The following statements are equivalent:*

- (1)  *$A$  is a weakly compact  $B^\#$ -algebra with a dense socle.*
- (2)  *$A$  is the  $B(\infty)$ -sum of  $B^\#$ -algebras  $A_\alpha$ ,  $\alpha \in \Omega$ , each of which is the algebra of all uniform limits of operators of finite rank on a reflexive Banach space.*

**PROOF.** (1)  $\Rightarrow$  (2). Bonsall [2, Theorem 6] has shown that if  $A$  is a  $B^\#$ -annihilator algebra, then  $A$  is isomorphic and isometric to the  $B(\infty)$ -sum of its minimal closed two-sided ideals  $A_\alpha$ . For the case of a  $B^\#$ -algebra with a dense socle, Bonsall's proof applies almost word for word if Lemmas 3.2 and 3.3 are borne in mind.

That each  $A_\alpha$  is a weakly compact algebra in its own right is clear and that  $A_\alpha$  is simple follows readily from a routine argument which depends essentially on the fact that  $A$  is semi-simple. Thus each  $A_\alpha$  is a simple, weakly compact,  $B^\#$ -algebra with minimal ideals. (That  $A_\alpha$  contains a minimal left ideal of its own follows from the fact that  $A_\alpha$  contains a minimal left ideal of  $A$  which is also a minimal left ideal of  $A_\alpha$ .) Hence by [7, Theorem 3.1], each  $A_\alpha$  is the algebra of all uniform limits of operators of finite rank on a reflexive Banach space.

(2)  $\Rightarrow$  (1). Each  $A_\alpha$ , being the algebra of all uniform limits of operators of finite rank on a reflexive Banach space, is a  $B^\#$  annihilator algebra [2, Theorem 2]. Since a  $B^\#$ -algebra is semi-simple, the  $B(\infty)$ -sum of the  $A_\alpha$  is, by a result of Rickart's [8, p. 107], an annihilator algebra. That the  $B(\infty)$ -sum of an arbitrary class of  $B^\#$ -algebras is a  $B^\#$ -algebra is proved in [5, Lemma 4.7]. Thus the  $B(\infty)$ -sum  $A$  of the  $A_\alpha$  is a semi-simple annihilator algebra. From this it follows that

$A$  has a dense socle and by Theorem 2.1,  $A$  is weakly compact. This concludes the proof.

**COROLLARY.** *A  $B^{\#}$ -algebra is an annihilator algebra if and only if it has a dense socle and is a weakly compact algebra.*

It is to be noted that one-handed weak complete continuity is enough to prove (1)  $\Rightarrow$  (2). In fact, a very slight modification of the proofs of [7, Lemma 3.1 and Theorem 3.1] shows that a simple, left weakly compact  $B^{\#}$ -algebra with minimal ideals is isomorphic and isometric to the algebra  $F(X)$  of all uniform limits of operators of finite rank on a reflexive Banach space  $X$ . Since the  $B(\infty)$ -sum of the  $A_{\alpha}$  in Theorem 3.4 is an annihilator algebra and weakly compact, we obtain the following:

**THEOREM 3.5.** *A right weakly compact  $B^{\#}$ -algebra with a dense socle is a weakly compact algebra.*

#### REFERENCES

1. F. F. Bonsall and A. W. Goldie, *Annihilator algebras*, Proc. London Math. Soc. **3** (1954), 154–167.
2. F. F. Bonsall, *A minimal property of the norm in some Banach algebras*, J. London Math. Soc. **29** (1954), 156–164.
3. N. Dunford and J. Schwartz, *Linear operators*. Part I, Interscience, New York, 1958.
4. T. Ogasawara and K. Yoshinaga, *Weakly completely continuous Banach\*-Algebras*, J. Sci. Hiroshima Univ. Ser. A **18** (1954), 15–36.
5. A. Olubummo, *Left completely continuous  $B^{\#}$ -algebras*, J. London Math. Soc. **32** (1957), 270–276.
6. ———,  *$B^{\#}$ -algebras with a certain set of left completely continuous elements*, J. London Math. Soc. **34** (1959), 367–369.
7. ———, *Operators of finite rank in a reflexive Banach space*, Pacific J. Math. **12** (1962), 1023–1027.
8. C. E. Rickart, *General theory of Banach algebras*, Van Nostrand, New York, 1960.

UNIVERSITY COLLEGE, IBADAN, NIGERIA