

PROXIMAL RELATIONS IN TOPOLOGICAL DYNAMICS¹

T. S. WU

In this note we shall prove that when the proximal relation of a transformation group (X, T, π) with compact phase space X is transitive (i.e., it is an equivalence relation), then it is equivalent with the syndetically proximal relation. This would answer two questions in [4, Remark 5].

Standing notations. Let (X, T, π) be a transformation group with compact phase space. The proximal relation of (X, T, π) is denoted by $P(X)$ and the syndetically proximal relation by $L(X)$. The product transformation group induced by (X, T, π) will be denoted by $(X \times X, T, \rho)$, which is defined by $(x, y)\rho^t = (x\pi^t, y\pi^t)$ for $(x, y) \in X \times X$ and $t \in T$. For simplicity we shall write xt for $x\pi^t$ and $(xt, yt) = (x, y)t$ for $(x, y)\rho^t$.

Reference. The proximal relation was studied in [1], [2], [3], [4]. The syndetically proximal relation was defined and studied in [4].

PROPOSITION. *If $P(X)$ is transitive, then $P(X) = L(X)$.*

PROOF. Since $P(X)$ is transitive, so is $P(X \times X)$ [1]. Then each orbit closure $\text{Cl}(x, y)T$ in $(X \times X, T, \rho)$ contains a unique minimal set. Let $(x, y) \in P(X)$. If $\text{Cl}(x, y)T - P(X) \neq \emptyset$, then there is $(a, b) \in \text{Cl}(x, y)T - P(X)$. Let M be the (unique) minimal set contained in $\text{Cl}(a, b)T$. There are two cases.

Case 1. $M \cap P(X) = \emptyset$. By Lemma 2 of [1], there is a point $(u, v) \in M$ such that $((x, y), (u, v)) \in P(X \times X)$. This shows that $(x, u) \in P(X)$, $(y, v) \in P(X)$, a fortiori, $(u, v) \in P(X)$ by the transitivity of $P(X)$. We have the contradiction.

Case 2. $M \cap P(X) \neq \emptyset$. By the definition of $P(X)$, if $(x', y') \in P(X)$ and N is the minimal set contained in $\text{Cl}(x', y')T$, then $N \subset \Delta(X)$, the diagonal of $X \times X$. This shows that $M \subset \Delta(X)$, a fortiori, $(a, b) \in P(X)$. We have the contradiction also. Hence, $\text{Cl}(x, y)T \subset P(X)$ when $(x, y) \in P(X)$. By Lemma 5 of [4], $P(X) \subset L(X)$, $P(X) = L(X)$.

COROLLARY. *$P(X)$ is an equivalence relation if and only if $P(X \times X) = P(X) \times P(X)$.*

Received by the editors February 10, 1964.

¹ This work was supported by NSF Contract No. G-20301

REFERENCES

1. J. Auslander, *On the proximal relation in topological dynamics*, Proc. Amer. Math. Soc. **11** (1960), 890-895.
2. R. Ellis, *A semigroup associated with a transformation group*, Trans. Amer. Math. Soc. **94** (1960), 272-281.
3. R. Ellis and W. Gottschalk, *Homomorphisms of transformation groups*, *ibid.*, **94** (1960), 258-271.
4. J. P. Clay, *Proximity relations in transformation groups*, Trans. Amer. Math. Soc. **108** (1963), 88-96.

STANFORD UNIVERSITY

A TOTALLY BOUNDED, COMPLETE UNIFORM SPACE IS COMPACT

D. L. FRANK

Let X be a set and U a uniformity on X . We will show that if (X, U) is totally bounded, every net in X has a Cauchy subnet. For each $d \in U$, let S_d^1, \dots, S_d^n be a finite covering of X by d -spheres. Let T_d be the topology on X having S_d^1, \dots, S_d^n as its subbasis. Clearly the space (X, T_d) is compact. Therefore, $Y = \prod_{d \in U} (X, T_d)$ is compact.

Now, let (p_i) be a net in X . Then $\Delta \circ (p_i)$ is a net in Y , where $\Delta: X \rightarrow Y$ is the diagonal. By compactness, there exists a convergent subnet, (q_j) , of $\Delta \circ (p_i)$. Then $\Delta^{-1} \circ (q_j)$ is a subnet of (p_i) which is clearly Cauchy.

Thus, if (X, U) is also complete, every net in X has a convergent subnet, so (X, U) is compact.

COLUMBIA UNIVERSITY

Received by the editors March 12, 1964.