

NONSINGULAR PATH FIELDS ON COMPACT TOPOLOGICAL MANIFOLDS

ROBERT F. BROWN¹ AND EDWARD FADELL²

1. **Introduction.** If X is a topological space, a path field on X is a map $s: X \rightarrow X^I$ such that $s(x)$ is either a path beginning at x and never hitting its initial point again or $s(x)$ is the constant path at x . The path field s is nonsingular if $s(x)$ is never constant. This concept generalizes the notion of vector field on a differentiable manifold. In [4] it was shown that a compact, connected, *orientable*, topological manifold (without boundary) admits a nonsingular path field if, and only if, its Euler characteristic is zero. Our objective here is to prove this theorem without the assumption of orientability and, in addition, for compact manifolds with boundary.

2. **The Lefschetz formula.** Let X denote an ANR (compact metric) and $f: X \rightarrow X$ a given map. Then, F. Browder [2] and also Leray and Deleanu [12], [6] have developed a local index theory which assigns to every open set $O \subset X$, such that f has no fixed points on its boundary ∂O , an integer $i(f, O)$ called the index of f on O . This index function has very nice properties listed explicitly in [2, p. 256]. Incidentally, the theory only requires f to be defined on \bar{O} , the closure of O , i.e., $f: \bar{O} \rightarrow X$.

Suppose we add the assumption that f has an isolated fixed point x_0 and X is locally Euclidean at x_0 , i.e., there is an open set U , containing x_0 , homeomorphic to Euclidean n -space R^n . Then, there is an index $I_f(x_0)$ for f at x_0 defined in terms of local degree ([13], [15]).

PROPOSITION (2.1). *If O is any open set in X containing x_0 , then for O sufficiently small,*

$$i(f, O) = (-1)^n I_f(x_0).$$

PROOF. Let $h: R^n \rightarrow U$ denote a homeomorphism such that $h(0) = x_0$. Let C denote the closed unit ball in R^n , C^0 its interior, and $V = h(C^0) \subset U$. We may assume that V contains no fixed points of f other than x_0 . We will show that if O is any open set containing x_0 such that $\bar{O} \cup f(\bar{O}) \subset V$, then $i(f, O) = (-1)^n I_f(x_0)$. Let $O_1 = h^{-1}(O) \subset C^0$. $h^{-1}f: \bar{O} \rightarrow C$ has the property that $h(h^{-1}f) = f$ has no fixed points on ∂O . Therefore, by the Commutative Property of i [2]

Presented to the Society, August 27, 1964; received by the editors April 16, 1964.

¹ Supported, in part, by the Air Force Office of Scientific Research.

² Supported by the National Science Foundation under Grant NSF G-23849 and the Wisconsin Alumni Research Foundation.

$$i((h^{-1}f)h, O_1) = i(h(h^{-1}f), O) = i(f, O).$$

Let $g = h^{-1}fh: \bar{O}_1 \rightarrow C$. If $G: C \rightarrow C$ is any extension of g , then $i(f, O) = i(G, O_1)$. Since C is a polyhedron, $i(G, O_1)$ is the local index restricted to the category of polyhedra and continuous maps. But, O'Neill [13] has shown that a local index function satisfying appropriate conditions (satisfied by the Browder-Leray-Deleanu index) is unique on this category of polyhedra. His uniqueness proof shows, in fact, that $i(G, O_1) = (-1)^n I_G(0)$. But by definition, $I_f(x_0) = I_G(0)$. Therefore, $i(f, O) = (-1)^n I_f(x_0)$.

PROPOSITION (2.2) (THE LEFSCHETZ FORMULA). *Let X denote an ANR (compact metric), $f: X \rightarrow X$ a given map with a finite number of fixed points, x_1, \dots, x_r . If each $x_j, 1 \leq j \leq r$, possesses an open neighborhood U_j homeomorphic to $R^{n(j)}$, then*

$$\Lambda(f) = \sum_{j=1}^r (-1)^{n(j)} I_f(x_j),$$

where $\Lambda(f)$ is the Lefschetz number of f .

PROOF. Choose sufficiently small, mutually disjoint open sets O_j such that $x_j \in O_j$ and $i(f, O_j) = (-1)^{n(j)} I_f(x_j)$. Then, the whole space X is an open set and $X - \bigcup_{j=1}^r O_j$ has no fixed points of f . Therefore, by the Additivity Property [2] of the index i ,

$$i(f, X) = \sum_{j=1}^r i(f, O_j).$$

But the Normalization Property [2] tells us that $i(f, X) = \Lambda(f)$, the Lefschetz number of f . Thus,

$$\Lambda(f) = i(f, X) = \sum_{j=1}^r i(f, O_j) = \sum_{j=1}^r (-1)^{n(j)} I_f(x_j).$$

COROLLARY (2.3). *Let M denote a compact topological n -manifold with a finite number of fixed points x_1, \dots, x_r . Then*

$$\Lambda(f) = (-1)^n \sum_{j=1}^r I_f(x_j).$$

If M is a compact topological manifold with boundary B , the result is still valid provided the fixed points x_1, \dots, x_r , do not belong to B .

REMARK (2.4). Proposition (2.2) generalizes the form of the Lefschetz formula proved in [5], as well as Theorem 3 in [15].

3. **A basic theorem.** We recall first [7] that if $\mathfrak{F}=(E, p, B)$, $\mathfrak{F}_0=(E_0, p_0, B)$ are regular fiber spaces, in the sense of Hurewicz [10] over the same base B , then $(\mathfrak{F}, \mathfrak{F}_0)=(E, E_0, p, B)$ is a fibered pair provided $E_0 \subset E, p_0=p|_{E_0}$ and \mathfrak{F} admits a lifting function λ such that λ lifts paths in B into E_0 if the preassigned initial point is in E_0 . $(F, F_0)=(p^{-1}(b), p_0^{-1}(b)), b \in B$, is the fiber of $(\mathfrak{F}, \mathfrak{F}_0)$. A generalized n -plane bundle (n -gpb) [7] is a fibered pair $(\mathfrak{F}, \mathfrak{F}_0)$ with fiber (F, F_0) such that

- (i) there exists a cross section $\sigma: B \rightarrow E$ such that $E_0 = E - \sigma(B)$,
- (ii) $(F, F_0) \sim (R^n, R^n - 0)$.

If $s: B \rightarrow E$ is a cross section, $b \in B$ is called a singular point of s if $s(b) \notin E_0$. Our objective here is the following.

THEOREM (3.1). *Let $(\mathfrak{F}, \mathfrak{F}_0)=(E, E_0, p, M)$ denote an n -gpb, $n \geq 2$, with fiber (F, F_0) and base M , where M is a compact connected topological n -manifold with boundary B . Suppose $s_0: B \rightarrow E_0$ is a given partial cross section (without singularities). Then, there is a cross section $s: M \rightarrow E$ with at most one singularity, which extends s_0 .*

PROOF. The simpler case $B = \emptyset$ is given in [4] and we therefore assume $B \neq \emptyset$.

I. First we observe that by [3] the boundary B is "collared," i.e., there is an imbedding $h: B \times [0, 1] \rightarrow M$ such that $h(b, 0) = b, b \in M$. Let $A = h(B \times [0, 1])$, $A^0 = h(B \times [0, 1])$. Choose an open cover χ (all coverings are considered finite) of M so fine that if $U \in \chi$ and $U \cap B \neq \emptyset$, then the star of U (rel χ) is contained in A^0 . Since M is an ANR (compact metric), and hence LC^* [11], there are open covers $\alpha_0 > \beta_0 > \chi$ (α_0 refines β_0, β_0 refines χ) such that if $\alpha > \alpha_0$ and N_α is the geometric nerve of α , there are maps

$$N_\alpha \begin{matrix} \xrightarrow{q_\alpha} \\ \xleftarrow{p_\alpha} \end{matrix} M$$

with the following properties.

- (a) $q_\alpha p_\alpha \sim_\chi 1$.
- (b) If $x \in M, p_\alpha(x) \in |V_{i_0}, \dots, V_{i_k}|$, where V_{i_0}, \dots, V_{i_k} are all the open sets in α containing x , and $|V_{i_0}, \dots, V_{i_k}|$ is the k -simplex determined by the "vertices" V_{i_j} .
- (c) If $V \in N_\alpha$, is a "vertex," $q_\alpha(V) \in V$ and for any simplex τ of $N_\alpha, q_\alpha(\tau)$ belongs to some member of β_0 .

Since M has dimension n , we may choose α so that $K = N_\alpha$ is an n -complex, $\alpha > \alpha_0$. Let L denote the following subcomplex of K . L is the union of all closed stars of vertices $V \in \alpha$ such that $V \cap B \neq \emptyset$.

Let $f = q_\alpha$, $g = p_\alpha$, where q_α , p_α are chosen to satisfy (a), (b), and (c). Then

- (1)
$$K \begin{matrix} \xrightarrow{f} \\ \xleftarrow{g} \end{matrix} M,$$
- (2)
$$fg \sim_x 1,$$
- (3)
$$f(L) \subset A^0,$$
- (4)
$$g(B) \subset L.$$

II. We now show that the given cross section $s_0: B \rightarrow E_0$ can be extended to $s_1: A \rightarrow E_0$. This is done by applying the following simple lemma.

LEMMA (3.2). *If $\mathfrak{F} = (E_0, p_0, A)$ is a regular fiber space and $B \subset A$ is a strong deformation retract of A , then a partial cross section $s_0: B \rightarrow E_0$ is always extendable to a cross section $s_1: A \rightarrow E_0$.*

III. The map $f: K \rightarrow M$ induces an n -gpb $(\mathfrak{F}^*, \mathfrak{F}_0^*) = (E^*, E_0^*, p^*, K)$ over K with fiber (F, F_0) using the given fibered pair $(\mathfrak{F}, \mathfrak{F}_0)$ over M . Since $f(L) \subset A$, the partial cross section $s_1: A \rightarrow E_0$ induces a cross section $s_1^*: L \rightarrow E_0^*$. Since $\pi_i(F_0) = 0$ for $i < n - 1$, s_1^* has an extension $s^*: K \setminus \{n-1\} \cup L \rightarrow E_0^*$. If σ^n is an n -simplex of $K - L$, a simple argument extends s^* to σ^n where s^* takes on values in E_0^* except at the barycenter of σ^n . Thus we obtain a cross section $s^*: K \rightarrow E$ such that s^* has only a finite number of singularities y_1, \dots, y_r , each the barycenter of a n -simplex of $K - L$.

IV. As is usual with induced fibrations, there is an associated commutative diagram

$$\begin{array}{ccc} E^* & \xrightarrow{\tilde{f}} & E \\ p^* \downarrow & & \downarrow p \\ K & \xrightarrow{f} & M \end{array}$$

where $\tilde{f}(E_0^*) \subset E_0$. Let us investigate the map $\phi = \tilde{f}s^*g$. First, we remark that ϕ is not a cross section, in fact, $p\phi = p\tilde{f}s^*g = fp^*s^*g = fg \sim_x 1$. Let $H: M \times I \rightarrow M$ denote a homotopy with the following properties. If $\tilde{H}: M \rightarrow M^I$ is the corresponding map into the space of paths then,

- (i) for $x \in M$, $\tilde{H}(x)$ begins at $fg(x)$ and ends at x ,
- (ii) for $x \in B$, $\tilde{H}(x) \in A^I$.

Now, if λ is a regular lifting function for $(\mathfrak{F}, \mathfrak{F}_0)$, define a cross section $\psi: M \rightarrow E$ by

$$\psi(x) = \lambda[\phi(x), \tilde{H}(x)](1), \quad x \in M.$$

Thus, ϕ can be “pulled over” to a cross section ψ with the property that if $\phi(x) \in E_0$, then $\psi(x) \in E_0$, $x \in M$. Let $\psi_0 = \psi|_B: B \rightarrow E_0$. We wish to show that s_0 and ψ_0 are homotopic as cross sections into E_0 . If $x \in B$, $\phi(x) = \tilde{f}s^*g(x) = \tilde{f}s_1^*g(x) = s_1fg(x)$. Thus, $s_1\tilde{H}(x)$ is a path in E_0 beginning at $\phi(x)$ and ending at $s_1(x) = s_0(x)$. $\lambda[\phi(x), \tilde{H}(x)]$ is a path in E_0 beginning at $\phi(x)$ and ending at $\psi_0(x)$. Both of these paths lie over $\tilde{H}(x)$. Applying Proposition 1 of [8], we obtain $s_0 \sim \psi_0: B \rightarrow E_0$ as cross sections. We note, at this point, the singularities of ψ . Let $Q_j = g^{-1}(y_j)$, where y_1, \dots, y_r are the singularities of s^* (III). The sets Q_j are closed, mutually disjoint, and none of them intersect the boundary B . Furthermore, the initial cover χ may be chosen so that each set Q_j is a compact subset of a Euclidean neighborhood which does not intersect the boundary. Thus, if $Q = \cup Q_j$, we see that the singularities of ψ lie in Q .

V. Now, using the techniques in the proof of Theorem 1.10 [4], there is a cross section $\zeta: M \rightarrow E$, with a finite number of singularities, which agrees with ψ on the boundary B . The proof of Lemma 1.9 [4] will yield a cross section $s': M \rightarrow E$ with one singularity such that s' agrees with ζ (and hence with ψ) on the boundary B . Thus, if $s'_0 = s'|_B$, $s'_0 \sim s_0$ as cross sections into E_0 .

VI. To complete the proof we proceed as follows. Let $x_0 \in M - B$ denote the singular point for s' and choose a small concentric n -cell neighborhood $\bar{V} \supset V \supset \bar{U} \supset U$ such that $x_0 \in U$. Then $(M - U, B)$ is an ANR pair. Applying the fiber homotopy extension theorem (FHET) in [1]. There is a homotopy

$$G: (M - U) \times I \rightarrow E_0$$

such that $G_0 = s'|_{M-U}$, G_t is a cross section over $M - U$ for each t , $0 \leq t \leq 1$, and $G_1|_B = s_0$. Now, $(M, M - U)$ is again an ANR pair and G_0 has s' as an extension to M . Again applying the FHET, this time in the fiber space (E, p, B) , G has an extension to a homotopy

$$G': M \times I \rightarrow E$$

such that $G'_1 = s''$ is a cross section over M which extends s_0 and has all its singularities in U . s'' can be redefined in U , using the techniques already referred to, to produce a cross section s which has $x_0 \in U$ as its only singular point and which extends s_0 .

4. The main result. Let X denote a topological space and

$$T_0(X) = \{ \omega \in X^I: \omega(t) = \omega(0) \text{ if, and only if, } t = 0, 0 \leq t \leq 1 \}.$$

Let $p_0: T_0(X) \rightarrow X$ be given by $p_0(\omega) = \omega(0)$. Let $T(X)$ denote $T_0(X)$ plus all the constant paths on X and $p: T(X) \rightarrow X$ be given by $p(\omega)$

$=\omega(0)$. A path field on X is a cross section for p , i.e., a map $s: X \rightarrow T(X)$ such that $ps=1$. A singularity for s is a point x such that $s(x)$ is the constant path at x . A nonsingular path field is a path field without singularities.

If $X=M$, a topological n -manifold (with empty boundary), then $(\mathfrak{J}, \mathfrak{J}_0) = (T(M), T_0(M), p, M)$ is an n -gpb [7], called the tangent fiber space of M .

Suppose that M is a compact topological n -manifold with boundary B . Let $h: B \times I \rightarrow M$ denote an imbedding which puts a "collar" on B and $h(x, 0) = x$, $x \in B$. Let $A = h(B \times I)$, $A^0 = h(B \times [0, 1))$, $A_1 = h(B \times \{1\})$. Then, h may be chosen so that if $M' = M - A^0$, $B' = A_1$, then (M', B') is a homeomorphic copy of (M, B) . Let $(\mathfrak{J}', \mathfrak{J}'_0) = (T, T_0, p, M - B)$ denote the tangent fiber space of $M - B$ and $(\mathfrak{J}', \mathfrak{J}'_0) = (T', T'_0, p', M')$ the restriction of $(\mathfrak{J}, \mathfrak{J}_0)$ to M' .

LEMMA (4.1). *Suppose $s_0: B' \rightarrow T_0(M')$ is a given nonsingular path field (in M') on B' . Since $T_0(M') \subset T'_0$, s_0 can be considered as a partial cross section over B' in the fibered pair $(\mathfrak{J}', \mathfrak{J}'_0)$. Then, s_0 admits an extension to a path field $s: M' \rightarrow T_0(M')$ if, and only if, $s_0: B' \rightarrow T'_0$ admits an extension to a cross section $s': M' \rightarrow T'_0$.*

PROOF. The necessity is trivial. Suppose, therefore, that $s_0: B' \rightarrow T'_0$ admits an extension, $s': M' \rightarrow T'_0$. s' would serve for s except for the fact that some of the paths might wander outside of M' . Such paths, however, start in $M' - B'$. Let $\rho: M - B \rightarrow M'$ denote a retraction such that ρ maps points outside of M' into B' . Then ρ induces $\rho': (M - B)^I \rightarrow M'^I$ and $s = \rho's': M' \rightarrow T_0(M')$, is the required extension of s_0 .

THEOREM (4.2). *Let M denote a compact connected n -manifold ($n \geq 2$) with boundary B and $s_0: B \rightarrow T_0(M)$ a given nonsingular path field (in M) on B . Then, s_0 can be extended to a nonsingular path field $s: M \rightarrow T_0(M)$ if, and only if, the Euler characteristic $\chi(M) = 0$.*

PROOF. The proof for an empty boundary B is simpler and is implicit in the argument for the case $B \neq \emptyset$. We, therefore, assume $B \neq \emptyset$.

The necessity is immediate since here we get a fixed point free map of M into itself homotopic to the identity.

To prove the sufficiency, identify (M, B) with (M', B') of the previous lemma. Using the notation in the previous lemma s_0 can be considered as a partial cross section $s_0: B' \rightarrow T'_0$, and it suffices to extend s_0 to $s: M' \rightarrow T'_0$. Now, $(\mathfrak{J}', \mathfrak{J}'_0)$ is an n -gpb and hence by our basic Theorem (3.1), there is a cross section $s': M' \rightarrow T'$ with at most

one singularity (in $M' - B'$) and $s'|_{B'} = s_0$. We may assume (see proof of (4.1)) that $s'(M') \subset T(M')$. s' gives rise to a map $f: M' \rightarrow M'$ by setting $f(x) = s'(x)(1)$, $x \in M'$, and $f \sim 1$. f has a single fixed point x_0 and by (1.3) $\Lambda_f = (-1)^n I_f(x_0)$. But $\chi(M) = \Lambda_f$ and hence the assumption that $\chi(M) = 0$ implies $I_f(x_0) = 0$. Now, using an argument identical with that in [4, 2.3], $I_f(x_0) = 0$ is a sufficient condition for redefining s' in a neighborhood of x_0 to produce an extension $s: M' \rightarrow T'_0$ of s_0 . Thus, our theorem follows.

Since a compact topological n -manifold M with boundary B always admits a nonsingular path field $s_0: B \rightarrow T_0(M)$ on its boundary we have the following.

THEOREM (4.3). *A compact connected topological n -manifold M , with or without boundary (not necessarily orientable), admits a nonsingular path field if, and only if, $\chi(M) = 0$. In the case of a manifold with boundary, the nonsingular path field may be preassigned on its boundary.*

If M is a compact topological manifold with boundary B , a path field $s: M \rightarrow T(M)$ is called *regular* if $x \in B$ implies $s(x)$ is a path in B .

THEOREM (4.4). *A compact connected topological n -manifold M with boundary components B_1, \dots, B_r , admits a regular nonsingular path field if, and only if, $\chi(M) = 0$ and $\chi(B_j) = 0$, $1 \leq j \leq r$.*

THEOREM (4.5). *A compact connected topological n -manifold M with connected boundary B admits a regular nonsingular path field if, and only if, $\chi(M) = 0$.*

PROOF. We prove only the sufficiency and it suffices to show $\chi(B) = 0$. If n is even, $n - 1$ is odd and $\chi(B) = 0$. Otherwise, if n is odd, let $2M$ denote the double of M . Then

$$\chi(2M) = 2\chi(M) - \chi(B).$$

Therefore, since $\chi(2M) = 0$ and $\chi(M) = 0$, we have $\chi(B) = 0$.

The following theorem bears on a question of M. Hirsch [9]. It is an immediate corollary of (4.3).

COROLLARY (4.6). *A compact connected topological n -manifold M , with or without boundary (not necessarily orientable) admits a map $f: M \rightarrow M$ which is fixed point free and homotopic to the identity if, and only if, $\chi(M) = 0$.*

REMARK (4.7). An example of Y. H. Clifton, namely $S^2 \vee S^1 \vee S^1$ shows that (4.6) is false in general for polyhedra. However, for a large class of polyhedra (zweidimensional zusammenhängende) Wecken [14] has proved (4.6). Our result (4.3) is much stronger than (4.6).

REMARK (4.8). The analogue of (4.3) for compact differentiable n -manifolds and nonsingular vector fields is well known, at least in the case of no boundary. The result for differentiable manifolds with boundary is also classical (H. Hopf, *Math. Ann.* **96** (1927), 225–250). Both of these results are obtainable as corollaries of our results for topological manifolds by showing that the tangent bundles in question are fiber homotopy equivalent to appropriate tangent fiber spaces. The no-boundary case is given in [4], [7] and the case of a differentiable manifold M with boundary B is handled as follows. First “shrink” (M, B) to another copy (M', B') via a tubular neighborhood of B . Then, the tangent bundle of M may be identified with the tangent bundle of M restricted to M' . This restricted tangent bundle is easily seen to be fiber homotopy equivalent to the tangent fiber space of $M - B$, restricted to M' . Lemma (4.1) completes the necessary paraphernalia.

BIBLIOGRAPHY

1. G. Allaud and E. Fadell, *A fiber homotopy extension theorem*, *Trans. Amer. Math. Soc.* **104** (1962), 239–251.
2. F. E. Browder, *On the fixed point index for continuous mappings of locally connected spaces*, *Summa Brasil. Math.* **4** (1960), 253–293.
3. M. Brown, *Locally flat embeddings of topological manifolds*, *Topology of 3-Manifolds*, M. Fort, Ed., Prentice-Hall, Englewood Cliffs, N. J., 1962; pp. 83–91.
4. R. Brown, *Path fields on manifolds*, *Trans. Amer. Math. Soc.* **118** (1965), 180–191.
5. ———, *On the Lefschetz fixed point formula*, *Amer. J. Math.* **87** (1965), 1–10.
6. A. Deleanu, *Théorie des points fixes sur les rétracts de voisinage des espace convexe*, *Bull. Soc. Math. France* **87** (1959), 235–243.
7. E. Fadell, *Generalized normal bundles for locally flat imbeddings*, *Trans. Amer. Math. Soc.* **114** (1965), 488–513.
8. ———, *On fiber spaces*, *Trans. Amer. Math. Soc.* **90** (1959), 1–14.
9. M. Hirsch, Problem 18, *Combinatorial and differentiable structures*, *Problems in Differential and Algebraic Topology* (mimeographed), Summer Topology Institute, 1962, Seattle, Wash.
10. W. Hurewicz, *On the concept of fiber space*, *Proc. Nat. Acad. Sci. U.S.A.* **41** (1955), 956–961.
11. S. Lefschetz, *Topics in topology*, *Annals of Mathematical Studies*, No. 10, Princeton Univ. Press, Princeton, N. J., 1942.
12. J. Leray, *Théorie des points fixes. Indice total et nombre de Lefschetz*, *Bull. Soc. Math. France* **87** (1959), 221–233.
13. B. O'Neill, *Essential sets and fixed points*, *Amer. J. Math.* **75** (1953), 497–509.
14. F. Wecken, *Fixpunktklassen*. III, *Math. Ann.* **118** (1941–1943), 544–577.
15. J. Weier, *Fixpunkttheorie in topologischen Mannigfaltigkeiten*, *Math. Z.* **59** (1953), 171–190.

UNIVERSITY OF CALIFORNIA, LOS ANGELES,
 INSTITUTE FOR ADVANCED STUDY AND
 UNIVERSITY OF WISCONSIN