

ON THE BOUNDARY BEHAVIOR OF BLASCHKE PRODUCTS IN THE UNIT DISK

PETER COLWELL¹

1. Introduction. Let A be any sequence of points in the unit disk $D: \{ |z| < 1 \}$ such that $\sum_A (1 - |a|) < \infty$. The Blaschke product with respect to A , given by $B(z; A) = \prod_A (|a|/a) (a-z)/(1-\bar{a}z)$, defines an analytic function in D such that $|B(z; A)| < 1$ for z in D . F. Riesz [7, p. 94] showed that for almost all points $e^{i\theta}$ of $C: \{ |z| = 1 \}$ the radial limit $B(e^{i\theta}) = \lim_{r \rightarrow 1} B(re^{i\theta}; A)$ exists and is of modulus one. For later reference we state the following result of O. Frostman [4, p. 170].

THEOREM 1. *A necessary and sufficient condition that $B(z; A)$ and all its subproducts have radial limits of modulus one at $e^{i\theta}$ is that*

$$(1) \quad \sum_A [(1 - |a|)/|e^{i\theta} - a|] < \infty.$$

It is our purpose to consider the boundary behavior of Blaschke products possessing radial limits of modulus one at every point of C .

2. Because of condition (1), it is only at accumulation points of A that a Blaschke product $B(z; A)$ can possibly fail to have a radial limit of modulus one. The following theorem gives the restrictions to be imposed on A' , the derived set of A , in order that $B(z; A)$ have radial limits of modulus one at every point of C .

THEOREM 2. *Let E be a set on C . A necessary and sufficient condition that there exist a Blaschke product $B(z; A)$ for which $B(e^{i\theta})$ is defined and of modulus one at every point of C and such that $A' = E$ is that E be closed and nowhere dense on C .*

Let E be a closed and nowhere dense set on C . We shall construct a sequence A in D with $A' = E$ for which $\sum_A (1 - |a|) < \infty$ and for which (1) is satisfied at every point of C . Then, by Theorem 1, the corresponding Blaschke product $B(z; A)$ will have radial limits of modulus one at every point of C .

Presented to the Society, April 10, 1965; received by the editors August 30, 1965.

¹ The work presented here was done while the author held a National Aeronautics and Space Administration Traineeship at the University of Minnesota, and it forms a part of the author's doctoral dissertation written under the direction of Professor D. A. Storvick.

The complement of E with respect to C is a countable union of disjoint open arcs on C . Let F be the set of end-points of these open arcs, where we write $F = \{c_m, d_m \mid m=1, 2, 3, \dots\}$, with $|c_m| = |d_m| = 1$, $\xi_m = \arg c_m < \arg d_m = \delta_m$ for each m . It is clear that F is a countable dense subset of E .

For each m , let $l_m = \delta_m - \xi_m$, and let a number t , $0 < t < 1$, be chosen. Define k_m for each m to be the minimum of t and $l_m/2\pi$. For each m and $n=1, 2, 3, \dots$, we define the sequences $\{g_{mn}\}$ and $\{h_{mn}\}$ by $g_{mn} = \xi_m + (k_m)^n/2$, $h_{mn} = \delta_m - (k_m)^n/2$. For each fixed m , $\{g_{mn}\}$ is non-increasing in n , $\{h_{mn}\}$ is nondecreasing in n , while $\lim_{n \rightarrow \infty} g_{mn} = \xi_m$ and $\lim_{n \rightarrow \infty} h_{mn} = \delta_m$. For each m and n we let $r_{mn} = 1 - (k_m)^n/2^{m+n}$, and we let $A = \{a_{mn}, b_{mn} \mid m, n=1, 2, 3, \dots\}$, where $a_{mn} = r_{mn} \exp(ig_{mn})$ and $b_{mn} = r_{mn} \exp(ih_{mn})$.

It may be seen that $A' = E$ and $\sum_A (1 - |a|) < \infty$. For any point $e^{i\theta}$ of $C - E$, clearly (1) is satisfied. If $e^{i\theta}$ is a point of E , we see that $\theta \neq \arg a$ for each a in A . If we denote by $|\alpha - \beta|$ the length along C of the shorter arc from $e^{i\alpha}$ to $e^{i\beta}$, then $|\theta - \arg a| \neq 0$ for $e^{i\theta}$ in E and a in A . By a lemma of G. T. Cargo [2, p. 10], to show (1) is satisfied for $e^{i\theta}$ in E , it suffices to show that

$$(2) \quad \sum_A [(1 - |a|)/|\theta - \arg a|] < \infty.$$

However, for $e^{i\theta}$ in E we have $|\theta - \arg a_{mn}| \geq (k_m)^n/2$, $|\theta - \arg b_{mn}| \geq (k_m)^n/2$ for $m=1, 2, 3, \dots$, $n=1, 2, 3, \dots$, so that (2) will be satisfied for $e^{i\theta}$ in E .

Let $B(z; A)$ be a Blaschke product with radial limits of modulus one at every point of C , and let $E = A'$.

Of course E is closed, but suppose E is not nowhere dense on C . Then there is some arc I on C in which E is dense. Write $I = \{e^{i\theta} \mid \alpha \leq \theta \leq \beta\}$, and let γ , $0 < \gamma < \pi$, be arbitrarily chosen. We denote by S_θ the Stolz angle in D at $e^{i\theta}$ with vertex angle γ symmetric about the radius to $e^{i\theta}$. In the region $\{re^{i\theta} \mid 0 < r < 1, \alpha < \theta < \beta\}$ we can select a point a_1 of A . If $\arg a_1 = \phi_1$, we may choose α_{11}, α_{12} such that $\alpha \leq \alpha_{11} < \phi_1 < \alpha_{12} \leq \beta$ and such that a_1 is in S_θ for $\alpha_{11} < \theta < \alpha_{12}$. Now choose β_{11}, β_{12} such that $\alpha_{11} < \beta_{11} < \phi_1 < \beta_{12} < \alpha_{12}$, and let $J_1 = \{e^{i\theta} \mid \beta_{11} \leq \theta \leq \beta_{12}\}$. Clearly a_1 is in S_θ for every $e^{i\theta}$ in J_1 .

In the region $\{re^{i\theta} \mid |a_1| < r < 1, \beta_{11} < \theta < \beta_{12}\}$ we can select a point a_2 of A . If $\arg a_2 = \phi_2$, we choose α_{21}, α_{22} such that $\beta_{11} \leq \alpha_{21} < \phi_2 < \alpha_{22} \leq \beta_{12}$ and such that a_2 is in S_θ for $\alpha_{21} < \theta < \alpha_{22}$. Now we choose β_{21}, β_{22} such that $\alpha_{21} < \beta_{21} < \phi_2 < \beta_{22} < \alpha_{22}$, and we let $J_2 = \{e^{i\theta} \mid \beta_{21} \leq \theta \leq \beta_{22}\}$. We see that $J_2 \subset J_1$, while a_1 and a_2 are both in S_θ for all $e^{i\theta}$ in J_2 .

Continuing in this fashion, we construct a sequence $\{J_j\}$ of closed

arcs on C such that $J_1 \supset J_2 \supset \cdots \supset J_j \supset \cdots$, and we select a sequence $\{a_j\}$ of points in A with $|a_1| < |a_2| < \cdots < |a_j| < \cdots < 1$ and such that for each value of j and for every $e^{i\theta}$ in J_j , a_k is in S_θ , $k=1, 2, \cdots, j$.

Now $\cap J_j$, where the intersection is taken over all values of j , is not empty, and we can find a point $e^{i\phi}$ of $\cap J_j$ which is an accumulation point of $\{a_j\}$. Also, for each value of j , a_j is in S_ϕ .

We connect the points of $\{a_j\}$ in order of increasing index by a polygonal path $P(z)$ to $e^{i\phi}$ lying in S_ϕ . The limit of $B(z; A)$ as z approaches $e^{i\phi}$ along $P(z)$, if it exists, cannot be of modulus one, for $B(a_j; A) = 0$ for $j=1, 2, 3, \cdots$. An application of a theorem of E. Lindelöf [5] shows that $B(z; A)$ cannot then have a radial limit of modulus one at $e^{i\phi}$. From this contradiction we conclude that E must be nowhere dense on C .

The proof that E is necessarily nowhere dense on C , while cumbersome, uses only elementary techniques. By appealing to cluster set theory, a far more elegant proof is possible. The author is indebted to Professor K. Noshiro for the following alternative proof of the fact that E is nowhere dense on C .

Each point $e^{i\theta}$ of E is an accumulation point of the zeros of $B(z; A)$ and thus is an essential singularity of $B(z; A)$. By a theorem of W. Seidel [9, p. 211], the interior cluster set of $B(z; A)$ at $e^{i\theta}$, $C(B, e^{i\theta})$, is the closed unit disk. However, by hypothesis $B(z; A)$ possesses a radial limit at each point of C , so that radial cluster set for $B(z; A)$ at $e^{i\theta}$, $C_r(B, e^{i\theta})$, is a single point.

Hence at each point $e^{i\theta}$ of E we have $C(B, e^{i\theta}) \neq C_r(B, e^{i\theta})$. By a theorem of E. F. Collingwood [3, p. 5], E must be a set of category I on C . Since E is closed, E is necessarily nowhere dense on C .

THEOREM 3. *Let $B(z; A)$ be a Blaschke product with $B(e^{i\theta})$ defined and of modulus one at every point of C . Then, as a function of θ , $B(e^{i\theta})$ is discontinuous at $\theta = \theta_0$ if, and only if, $e^{i\theta_0}$ is in A' .*

If $e^{i\theta_0}$ is not a point of A' , then a theorem of C. Tanaka [10, p. 410] states that $B(z; A)$ is analytic throughout a neighborhood of $e^{i\theta_0}$. Throughout this neighborhood, $B(e^{i\theta}) = \lim_{r \rightarrow 1} B(re^{i\theta}; A) = B(e^{i\theta}; A)$, so that $B(e^{i\theta})$ is evidently continuous at $\theta = \theta_0$.

If $e^{i\theta_0}$ is a point of A' , then $e^{i\theta_0}$ is a singularity of $B(z; A)$. Consequently, as was proved by W. Seidel [9, p. 208], in each arc on C containing $e^{i\theta_0}$, $B(e^{i\theta})$ assumes every value of modulus one infinitely often. Then $B(e^{i\theta})$ is discontinuous at $\theta = \theta_0$.

From Theorems 2 and 3 follows immediately a corollary which is a special case of a theorem of A. J. Lohwater and G. Piranian [6, p. 5].

COROLLARY. Let $B(z; A)$ be any Blaschke product for which $B(e^{i\theta})$ is defined and of modulus one at every point of C . In order that a set E on C be exactly the set of points where the radial limit function $B(e^{i\theta})$ is discontinuous, it is necessary and sufficient that E be closed and nowhere dense on C .

3. For a Blaschke product $B(z; A)$ in D , the radial variation of $B(z; A)$ at a point $e^{i\theta}$ of C is defined to be $V(B; \theta) = \int_0^1 |B'(re^{i\theta}; A)| dr$. The quantity $V(B; \theta)$ is the length of the image under $B(z; A)$ of the radius to $e^{i\theta}$. G. T. Cargo [1, p. 425] proved that (1) is a necessary and sufficient condition for the radial variations of $B(z; A)$ and all its subproducts at $e^{i\theta}$ to be uniformly bounded.

THEOREM 4. Let $B(z; A)$ be a Blaschke product for which (1) holds at every point of C . Then, as a function of θ , $V(B; \theta)$ is a discontinuous at $\theta = \theta_0$ if, and only if, $e^{i\theta_0}$ is in A' .

Suppose $e^{i\theta_0}$ is not in A' . Then there is an open neighborhood N of $e^{i\theta_0}$ throughout which $B(z; A)$ is analytic, [10, p. 410], and $B(z; A)$ is analytic in $D \cup N$. Consequently, $B'(z; A)$ is defined and continuous throughout $D \cup N$.

If d denotes the distance from $e^{i\theta_0}$ to the closest boundary point of N , let R be the closed region $\{re^{i\theta} | 0 \leq r \leq 1, |e^{i\theta} - e^{i\theta_0}| \leq d/2\}$. Now $R \subset D \cup N$, and $B'(z; A)$ is uniformly continuous on R . Given any $\epsilon > 0$ there exists $\delta > 0$ such that for all r , $0 \leq r \leq 1$, $|B'(re^{i\theta}; A) - B'(re^{i\theta_0}; A)| < \epsilon$ when $|\theta - \theta_0| < \delta$.

Then for all θ , $|\theta - \theta_0| < \delta$, we have

$$\begin{aligned} |V(B; \theta) - V(B; \theta_0)| &= \left| \int_0^1 |B'(re^{i\theta}; A)| dr - \int_0^1 |B'(re^{i\theta_0}; A)| dr \right| \\ &\leq \int_0^1 |B'(re^{i\theta}; A) - B'(re^{i\theta_0}; A)| dr < \epsilon, \end{aligned}$$

so that $V(B; \theta)$ is continuous at $\theta = \theta_0$.

Suppose $e^{i\theta_0}$ is in A' while $V(B; \theta)$ is continuous at $\theta = \theta_0$. We may select a subsequence $\{r_k \exp(i\phi_k) | k = 1, 2, 3, \dots\}$ from A such that $r_k \leq r_{k+1}$ and $|\phi_k - \theta_0| \geq |\phi_{k+1} - \theta_0|$ for $k = 1, 2, 3, \dots$, $\lim_{k \rightarrow \infty} r_k \exp(i\phi_k) = e^{i\theta_0}$, and $\lim_{k \rightarrow \infty} V(B; \phi_k) = V(B; \theta_0)$.

Let s , $0 < s < 1$, be arbitrarily chosen, and let $k(s)$ be a positive integer such that $r_k \geq s$ when $k \geq k(s)$. Since for each value of k , $B[r_k \exp(i\phi_k); A] = 0$ while $|B[\exp(i\phi_k)]| = 1$ by Theorem 1, we see that for $k \geq k(s)$, $\int_s^1 |B'[r \exp(i\phi_k); A]| dr \geq 1$.

Thus for $k \geq k(s)$ we have $V(B; \phi_k) = \int_0^1 |B'[r \exp(i\phi_k); A]| dr \geq \int_0^s |B'[r \exp(i\phi_k); A]| dr + 1$. Now $B'(z; A)$ is uniformly continuous

on the compact subset $\{|z| \leq s\}$ of D , while for any r , $0 \leq r \leq s$, we have $\lim_{k \rightarrow \infty} |B'[r \exp(i\phi_k); A]| = |B'(re^{i\theta_0}; A)|$. Consequently, $\lim_{k \rightarrow \infty} \int_0^s |B'[r \exp(i\phi_k); A]| dr = \int_0^s \{\lim_{k \rightarrow \infty} |B'[r \exp(i\phi_k); A]| \} dr = \int_0^s |B'(re^{i\theta_0}; A)| dr$, and $V(B; \theta_0) = \lim_{k \rightarrow \infty} V(B; \phi_k) \geq \int_0^s |B'(re^{i\theta_0}; A)| dr + 1$, where $0 < s < 1$.

Since (1) holds for $e^{i\theta_0}$, $V(B; \theta_0) < \infty$, and $\lim_{s \rightarrow 1} \int_0^s |B'(re^{i\theta_0}; A)| dr = V(B; \theta_0)$, so that $V(B; \theta_0) \geq V(B; \theta_0) + 1$. We conclude that if $e^{i\theta_0}$ is in A' , then $V(B; \theta)$ is discontinuous at $\theta = \theta_0$.

We remark here that since $V(B; \theta)$ is a lower semicontinuous function of θ on $[0, 2\pi]$ (cf. [8, p. 235], Theorem 3 implies that $V(B; \theta)$ cannot have a relative maximum at $\theta = \theta_0$ if $e^{i\theta_0}$ is in A' .

4. It is known, [4, p. 177], that if a Blaschke product $B(z; A)$ satisfies

$$(3) \quad \sum_A [(1 - |a|)/|e^{i\theta} - a|^2] < \infty$$

at a point $e^{i\theta}$ of C , then the derivative of $B(z; A)$ has a finite radial limit at $e^{i\theta}$ given by

$$B'(e^{i\theta}) = \lim_{r \rightarrow 1} B'(re^{i\theta}; A) = B(e^{i\theta})e^{-i\theta} \sum_A [(1 - |a|^2)/|e^{i\theta} - a|^2],$$

where $B(e^{i\theta}) = \lim_{r \rightarrow 1} B(re^{i\theta}; A)$.

At any point $e^{i\theta}$ of C where (3) holds, (1) also holds, and a slight modification of the proof of Theorem 2 justifies

THEOREM 5. *A necessary and sufficient condition that a set E on C be the set of accumulation points of the zeros of a Blaschke product $B(z; A)$ whose derivative has a finite radial limit at every point of C is that E be closed and nowhere dense on C .*

THEOREM 6. *Let $B(z; A)$ be a Blaschke product for which (3) holds at every point of C . As a function of θ , $B'(e^{i\theta})$ is discontinuous at $\theta = \theta_0$ if, and only if, $e^{i\theta_0}$ is in A' .*

Let $M(\theta) = \sum_A [(1 - |a|^2)/|e^{i\theta} - a|^2]$ for each θ ; we note that for each value of θ each summand of $M(\theta)$ is a continuous function of θ .

Suppose $e^{i\theta_0}$ is not a point of A' . Then for some $\delta > 0$, $|e^{i\theta_0} - a| \geq \delta$ for all points a of A , and for each point $e^{i\theta}$ of $C \cap \{z \mid |e^{i\theta_0} - z| \leq \delta/2\}$, we have $|e^{i\theta} - a| \geq \delta/2$ and $M(\theta) \leq (8/\delta^2) \sum_A (1 - |a|)$. Thus $M(\theta)$ converges uniformly to a continuous function of θ in $\{\theta \mid |e^{i\theta} - e^{i\theta_0}| \leq \delta/2\}$.

By Theorem 2, $B(e^{i\theta})$ is continuous at $\theta = \theta_0$, so $B'(e^{i\theta}) = B(e^{i\theta}) \cdot e^{-i\theta}$, $M(\theta)$ is continuous at $\theta = \theta_0$.

Suppose now that $e^{i\theta_0}$ is in A' and $B'(e^{i\theta})$ is continuous at $\theta = \theta_0$. We see that $M(\theta)$ is real-valued, and $M(\theta) \geq (1/4) \sum_A (1 - |a|)$. Further, since (3) holds, $|B(e^{i\theta})| = 1$ for all θ , and $|M(\theta) - M(\theta_0)| \leq |B'(e^{i\theta}) - B'(e^{i\theta_0})|$. The continuity of $B'(e^{i\theta})$ at $\theta = \theta_0$ implies that of $M(\theta)$ at $\theta = \theta_0$.

But then $B(e^{i\theta}) = [e^{i\theta} B'(e^{i\theta})] / M(\theta)$ is also continuous at $\theta = \theta_0$, and this contradicts Theorem 2. Thus $B'(e^{i\theta})$ is discontinuous at $\theta = \theta_0$ if $e^{i\theta_0}$ is in A' .

BIBLIOGRAPHY

1. G. T. Cargo, *The radial images of Blaschke products*, J. London Math. Soc. **36** (1961), 424-430.
2. ———, *The boundary behavior of Blaschke products*, J. Math. Anal. Appl. **5** (1962), 1-16.
3. E. F. Collingwood, *Cluster sets and prime ends*, Ann. Acad. Sci. Fenn. A. I. No. 250 (1958), 13 pp.
4. O. Frostman, *Sur les produits de Blaschke*, Kungl. Fysiogr. Sällsk. i Lund Forh. **12** (1942), 169-181.
5. E. Lindelöf, *Sur un principe général de l'analyse et ses applications à la théorie de la représentation conforme*, Acta Soc. Sci. Fenn. **46** (1915), 1-35.
6. A. J. Lohwater and G. Piranian, *The boundary behavior of functions analytic in a disk*, Ann. Acad. Sci. Fenn. A. I. No. 239 (1957), 17 pp.
7. F. Riesz, *Über die Randwerte einer analytischen Funktionen*, Math. Z. **18** (1923), 87-95.
8. W. Rudin, *The radial variation of analytic functions*, Duke Math. J. **22** (1955), 235-242.
9. W. Siedel, *On the distribution of values of bounded analytic functions*, Trans. Amer. Math. Soc. **36** (1934), 201-226.
10. C. Tanaka, *Boundary convergence of Blaschke products in the unit-circle*, Proc. Japan Acad. **39** (1963), 410-412.

UNIVERSITY OF MINNESOTA AND
IOWA STATE UNIVERSITY