

UNUSUAL POWER PRODUCTS AND THE IDEAL $[y^2]$

KATHLEEN B. O'KEEFE¹

A power product $P = y_{i(1)}y_{i(2)} \cdots y_{i(n)}$, where $y_{i(j)}$ is the $i(j)$ th derivative of y , has *weight* $w = \sum_{j=1}^n i(j)$ and *degree* $d = n$. The sequence of integers (e_1, e_2, \cdots, e_n) , where $e_k = \sum_{j=1}^k i(j) - k(k-1)$, is called the *weight sequence* of P . According to a result of H. Levi ([1], Theorem 1.2, p. 545), if some $e_k < 0$, then $P \equiv 0[y^2]$. The discovery by D. G. Mead [2] that $Q = y_1y_3y_4y_5 \equiv 0[y^2]$ shows that this criterion is not a necessary condition for membership in $[y^2]$. Q is an *unusual* power product; that is, Q is in $[y^2]$ and has a nonnegative weight sequence. By [3], there also exist unusual power products for $[y^p]$, $p > 2$; however, this note is concerned only with $[y^2]$. All unusual power products with weight sequences consisting of 0's, 1's, and 2's are described by Theorem IV, [2]:

A power product P with weight sequence (e_1, \cdots, e_n) , $0 \leq e_i \leq 2$, $i = 1, \cdots, n$, is unusual if and only if somewhere in the sequence at least one of the following patterns appears: 1, 2, 2, 1; 1, 2, 2, 2, 2, 0; 0, 2, 2, 2, 2, 1; 0, 2, 2, 2, 2, 2, 2, 0.

For particular arrangements of 0's, 1's, 2's, and 3's, similar results may be obtained; for example, using the notation of [2],

THEOREM A. *Let $g(a) = m(1, 3_1, 3_2, \cdots, 3_a, 2, 0)$, $[3_i = 3]$, then $g(a) = 0$ if and only if $a = 3$.*

The purpose of this note is to show that, in general, a finite list of patterns will not suffice to describe unusual power products. By Theorem B there are unusual power products of arbitrarily high degree with no proper factors in $[y^2]$. The following results from [2] are stated for easy reference:

- (a) $m(A, 0; B, 0) = m(A, 0)m(B, 0)$ for any sequences A and B
- (b) $m(1, 0) = -2$
- (c) $m(1, 1, C) = -m(1, C)$ for any sequence C
- (d) $m(1, 2, 1, D) = 2m(1, D)$ for any sequence D
- (e) $m(0, E) = m(E)$ for any sequence E
- (f) $m(1, 2, 2, F) = 2m(2, F) + m(1, 2, F)$ for any sequence F .

THEOREM B. *Let $f(a) = m(1, 2, 3_1, 3_2, \cdots, 3_a, 1)$, $[3_i = 3]$; then $f(a) = 0$ for all $a \geq 1$. Furthermore, if Q has the weight sequence $(1, 2, 3_1, \cdots, 3_a, 1)$, $a \geq 5$, then Q has no proper factor in $[y^2]$.*

Received by the editors May 17, 1965.

¹ This paper was written while the author held a fellowship from the American Association of University Women.

PROOF. The theorem rests on two easy lemmas.

LEMMA I. $m(1, 2, 3, 3, S) = -m(1, 2, 3, S)$, for any sequence S .

PROOF. $(1, 2, 3, 3, S)$ is the weight sequence of $Q = y_1y_3y_5y_6 \cdots$. Replacing y_5y_6 and using (a)–(d), (f), we get the equation

$$m(Q) = -2m(3, S) - 2m(1, 3, S) - 2m(2, 3, S) - m(1, 2, 3, S).$$

Use the equation $m(2, 3, S) = -m(3, S) - m(1, 3, S)$ to complete the proof.

In a similar fashion we prove

LEMMA II. $m(1, 2, 3, 2, S) = 2m(1, 2, S)$, for any sequence S .

Returning to the proof of Theorem B, by Lemma I, $f(a) = (-1)^{a-1}m(1, 2, 3, 1)$. But $(1, 2, 3, 1)$ represents the same power product as $(1, 2, 2, 1)$, namely $R = y_1y_3y_4y_6$; and by Theorem IV of [2], $R \equiv 0[y^2]$.

If P is a proper factor of Q , then P is either an α -term or a factor of

$$\begin{aligned} P_1(k) &= y_1y_3y_5y_6y_8 \cdots y_{2k}, \\ P_2(k) &= y_1y_3y_4y_6y_8 \cdots y_{2k}, \\ P_3(k) &= y_1y_3y_5^2y_8y_{10} \cdots y_{2k}, \\ P_4(k) &= y_0y_3y_5y_6y_8 \cdots y_{2k}, \\ P_5(k) &= y_1y_2y_5y_6y_8 \cdots y_{2k}. \end{aligned}$$

$P_1(k)$ is comparable to the α -term $y_1y_3y_5$ by Lemma I. $P_2(k)$ has the weight sequence $(1, 2, 2, \cdots, 2, 0)$; hence, is in $[y^2]$ if and only if $k=4$. By Lemma II, $m(P_3(k)) = 2m(P_2(k-2))$; and by (e) and (c) respectively, $m(P_4(k)) = m(P_2(k-1))$ and $m(P_5(k)) = -m(P_3(k-1))$. Thus Q has no proper factor in $[y^2]$.

BIBLIOGRAPHY

1. H. Levi, *On the structure of differential polynomials and on their theory of ideals*, Trans. Amer. Math. Soc. 51 (1942), 532–568.
2. D. G. Mead, *Differential ideals*, Proc. Amer. Math. Soc. 6 (1955), 420–432.
3. K. B. O'Keefe, *A property of the differential ideal $[y^p]$* , Trans. Amer. Math. Soc. 94 (1960), 483–497.

UNIVERSITY OF WASHINGTON