

# ON POLYNOMIALS CHARACTERIZED BY A CERTAIN MEAN VALUE PROPERTY

LEOPOLD FLATTO

Let  $\mathcal{V}$  denote the vector space of continuous real valued functions  $f(x)$  satisfying the mean value property

$$(1) \quad f(x) = \frac{1}{N} \sum_{i=1}^N f(x + ty_i)$$

for  $x \in R$ ,  $0 < t < \epsilon_x$  ( $R$  denotes an  $n$ -dimensional region;  $x$  and  $y_i$  are abbreviations for  $(x_1, \dots, x_n)$ ,  $(y_{i1}, \dots, y_{in})$ ). We assume that the  $y_i$ 's span  $E_n$  so that  $1 \leq n \leq N$ . We furthermore assume, without loss of generality, that  $y_1, \dots, y_n$  are linearly independent.

Friedman and Littman [5] have recently shown that  $\mathcal{V}$  consists of polynomials of degrees  $\leq N(N-1)/2$ . This bound is actually attained when the  $y_i$ 's form the  $N$  vertices of an  $(N-1)$ -dimensional regular simplex [see 4, p. 264]. On the other hand it is known that for  $n=2$ ,  $\deg f \leq N$  [see 4, Theorem 3.2]. The object of this paper is to obtain bounds on  $\deg \mathcal{V}$  and  $\dim \mathcal{V}$ , the bounds depending on  $N$  and  $n$  ( $1 \leq n \leq N$ ). We use the term  $\deg \mathcal{V}$ , to denote the maximum degree of the polynomials contained in  $\mathcal{V}$ . We also characterize for fixed  $N$  and varying  $n$  ( $1 \leq n \leq N$ ) those configurations for which  $\deg \mathcal{V}$  and  $\dim \mathcal{V}$  attain their maximum.

**THEOREM.** *We have*

$$(2) \quad \deg \mathcal{V} \leq \sum_{j=1}^n (N-j), \quad \dim \mathcal{V} \leq \prod_{j=0}^{n-1} (N-j)$$

so that for fixed  $N$  and varying  $n$  ( $1 \leq n \leq N$ )

$$(3) \quad \deg \mathcal{V} \leq \frac{N(N-1)}{2}, \quad \dim \mathcal{V} \leq N! \quad .$$

The latter bounds are obtained if and only if

$$n = N \quad \text{or} \quad n = N - 1 \quad \text{and} \quad \sum_{i=1}^N y_i = 0.$$

**REMARK.** The bounds in (2) are not best possible. For instance, we have stated above that for  $n=2$ ,  $\deg \mathcal{V} \leq N$  and this bound is best

---

Received by the editors September 7, 1965.

possible. For fixed  $n$  and  $N$  the problem of determining the maximum values of  $\deg \vee$ ,  $\dim \vee$  and the configurations for which these maximum values are attained remains open.

PROOF. We employ the following notation.

$$\frac{\partial}{\partial x} = \left( \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right), \quad x \cdot y = x_1 y_1 + \dots + x_n y_n,$$

$$P_k(x) = \sum_{i=1}^N (x \cdot y_i)^k \quad (1 \leq k < \infty).$$

It is shown in [5] that (1) is equivalent to the infinite system of homogeneous partial differential equations

$$(4) \quad P_k \left( \frac{\partial}{\partial x} \right) f = 0 \quad (1 \leq k < \infty)$$

and that  $\vee$ , which is thus the solution space of (4), is a finite dimensional space consisting of polynomials. Let  $R$  denote the ring of polynomials in  $x_1, \dots, x_n$  with real coefficients and let  $\mathfrak{P}$  denote the ideal generated by the  $P_k$ 's ( $1 \leq k < \infty$ ).  $R$ ,  $\mathfrak{P}$ , and  $\vee$  are vector spaces over the reals and it is known that  $R$  is the direct sum of  $\mathfrak{P}$  and  $\vee$ , i.e.  $R = \mathfrak{P} \oplus \vee$  [see 2, p. 53]. Thus the vector spaces  $R/\mathfrak{P}$  and  $\vee$  are isomorphic ( $R/\mathfrak{P} \cong \vee$ ).

$\deg \vee$  and  $\dim \vee$  will thus be determined if we know all the polynomials in  $\mathfrak{P}$ . We introduce the new variables  $\xi_i = x \cdot y_i$  ( $1 \leq i \leq N$ ). Since the  $y_i$ 's ( $1 \leq i \leq N$ ) are linearly independent we must have  $\xi_{n+k} = \sum_{i=1}^n a_{ki} \xi_i$  ( $1 \leq k \leq N-n$ ) for an appropriate choice of real  $a_{ki}$ 's. Let  $R'$  denote the ring of polynomials in  $\xi_1, \dots, \xi_n$  with real coefficients and let  $\mathfrak{P}'$  denote the ideal generated by the  $\eta_k$ 's where  $\eta_k = \sum_{i=1}^n \xi_i^k$  ( $1 \leq k \leq \infty$ ). We adopt the following notation:

$$\xi = (\xi_1, \dots, \xi_n), \quad i = (i_1, \dots, i_n), \quad \xi^i = \xi_1^{i_1}, \dots, \xi_n^{i_n}.$$

It is known [see 1, p. 41] that every polynomial  $Q(\xi)$  can be expressed as

$$(5) \quad Q(\xi) = \sum' R_i \xi^i,$$

where the summation in  $\sum'$  extends over those  $i$ 's for which  $0 \leq i_j \leq N-j$  ( $1 \leq j \leq n$ ) and  $R_i$  is a polynomial in  $\eta_1, \dots, \eta_N$ . This representation is unique for  $n=N$ . Let  $c_i$  denote the constant term in  $R_i$  and let  $S_i = R_i - c_i$ . Clearly  $S_i \in \mathfrak{P}'$ . It follows from (5) that  $Q(\xi) = \sum' C_i \xi^i + \sum' S_i \xi^i$  so that

$$(6) \quad Q(\xi) \equiv \sum' c_i \xi^i \pmod{\mathfrak{P}'}$$

As there are  $\prod_{j=0}^{n-1} (N-j)$  distinct  $\xi^i$ 's, (6) shows that  $\dim R'/\mathfrak{P}' \leq \prod_{j=0}^{n-1} (N-j)$ . Since  $\vee \cong R/\mathfrak{P} \cong R'/\mathfrak{P}'$  we have  $\dim \vee \leq \prod_{j=0}^{n-1} (N-j)$ . It follows furthermore from (6) that if  $Q$  is homogeneous and  $\deg Q > \sum_{j=1}^n (N-j)$ , then  $Q \in \mathfrak{P}'$ . This implies that if  $P(x)$  is homogeneous and  $\deg P > \sum_{j=1}^n (N-j)$  then  $P \in \mathfrak{P}$ . Thus  $\deg \vee \leq \sum_{j=1}^n (N-j)$ .

If  $n \leq N-2$ , then we conclude from (2) that  $\deg \vee < N(N-1)/2$ ,  $\dim \vee < N!$ . It remains to treat the two cases: (a)  $n = N$ , (b)  $n = N-1$ . In case (a) the  $\xi^i$ 's form a basis for  $R'/\mathfrak{P}'$ . For suppose that  $\sum' c_i \xi^i \equiv 0 \pmod{\mathfrak{P}'}$  for some choice of real  $c_i$ 's. Then  $\sum' c_i \xi^i = \sum_{j=1}^n T_j(\xi) \eta_j$  where the  $T_j$ 's are polynomials in  $\xi_1, \dots, \xi_n$ . But each  $T_j$  has a representation (5). I.e.  $T_j(\xi) = \sum' R_{ji}(\eta) \xi^i$  where the  $R_{ji}$ 's are polynomials in  $\eta_1, \dots, \eta_n$ . Thus  $\sum' c_i \xi^i = \sum_{j=1}^n \sum' R_{ji}(\eta) \xi^i = \sum' (\sum_{j=1}^n R_{ji}(\eta)) \xi^i$ . Since the representation (5) is unique for  $n = N$  we have

$$(7) \quad c_i = \sum_{j=1}^n R_{ji}(\eta_j)$$

The left side of (7) is void of  $\eta$ 's so that all  $R_{ji}$ 's and  $c_i$ 's equal 0. Thus  $\dim R'/\mathfrak{P}' = N!$  and since  $\vee \cong R/\mathfrak{P} \cong R'/\mathfrak{P}'$ ,  $\dim \vee = N!$ . Now  $\prod_{j=1}^{N-1} \xi_j^{N-j}$  has degree  $N(N-1)/2$  and  $\notin \mathfrak{P}'$ . This implies that there exists a homogeneous polynomial of degree  $N(N-1)/2 \notin \mathfrak{P}$ . Hence  $\deg \vee = N(N-1)/2$ .

If  $n = N-1$  then we distinguish two cases. If  $\sum_{i=1}^N y_i \neq 0$ , then it follows from [1, Theorem 2.2] that there exists an orthogonal transformation  $x = Tx'$  such that  $g(x') = f(Tx')$  is independent of  $x'_n$  and satisfies the equation

$$(8) \quad g(x'_p) = \frac{1}{N} \sum_{i=1}^N g(x'_p + ty_{p,i}),$$

where  $y_i = Ty'_i$ ,  $x'_p = (x'_1, \dots, x'_{n-1})$ ,  $y'_{pi} = (y'_{1i}, \dots, y'_{ni-1})$ . Let  $\vee'$  denote the solution space of (8). Clearly  $\deg \vee = \deg \vee'$ ,  $\dim \vee = \dim \vee'$ . It follows from (2) that  $\deg \vee = \deg \vee' < N(N-1)/2$ ,  $\dim \vee = \dim \vee' < N!$ . If  $\sum_{i=1}^N y_i = 0$ , then define

$$\bar{x} = (x_1, \dots, x_n, x_{n+1}), \quad y_i = (y_{i1}, \dots, y_{in}, 1) \quad (1 \leq i \leq N), \quad F(\bar{x}) = f(x).$$

We notice that  $\sum_{i=1}^N \bar{y}_i \neq 0$ . It therefore follows from [3, Theorem 2.2] that  $\vee$  is identical with the solution space  $\bar{\vee}$  of

$$(9) \quad F(\bar{x}) = \frac{1}{N} \sum_{i=1}^N F(\bar{x} + t\bar{y}_i).$$

Equation (9) is included in Case (a). It follows that  $\deg \nabla = \deg \overline{\nabla} = N(N-1)/2$ ,  $\dim \nabla = \dim \overline{\nabla} = N!$ .

#### REFERENCES

1. E. Artin, *Lectures on Galois theory*, Notre Dame Mathematical Lectures, Ann. Arbor, Michigan, 1944.
2. E. Fischer, *Über Algebraische Modulsysteme und lineare homogene partiell Differentialgleichungen mit konstanten Koeffizienten*, J. Reine Angew. Math. **140** (1911), 48–81.
3. L. Flatto, *Functions with a mean value property*, J. Math. Mech. **10** (1961), 11–18.
4. ———, *Functions with a mean value property*. II, Amer. J. Math. **85** (1963), 248–270.
5. A. Friedman and W. Littman, *Functions satisfying the mean value property*, Trans. Amer. Math. Soc. **102** (1962), 167–180.

BELFER GRADUATE SCHOOL OF SCIENCES, YESHIVA UNIVERSITY