

THE UNIFORMIZATION OF A HYPERBOLIC RIEMANN SURFACE

GEORGE INNIS

In this paper a hyperbolic Riemann surface is described, and the uniformizing function for this surface is constructed. The uniformizing function for this hyperbolic surface is an infinite product similar to the ones constructed for some parabolic surfaces in earlier studies [1], [2], [4].

Description of the surface. Let I denote the positive integers. For each $i \in I$, let C_i be the vertical line segment in the Z -plane of length 2 centered at the point $(1/i, 0)$. Let S_1 be the Z -sphere cut along C_1 and for each $i \in I$, $i > 1$, let S_i be the Z -sphere cut along C_{i-1} and along C_i . For each $i \in I$, join S_i to S_{i+1} along C_i in such a way as to form first-order branch points over $(1/i, 1)$ and $(1/i, -1)$. The resulting Riemann surface F is simply connected and is easily seen to be hyperbolic because the surface has a free edge along the vertical line segment from $-i$ to i .

Approximating surfaces. For each $i \in I$, let F_i be the Riemann surface formed from the first i sheets of F with the points of the curve C_i on S_i deleted. Let F_i^* be the closed surface formed from the first i sheets of F with C_i on S_i healed. There exists a unique function g_i mapping F_i^* onto the Z -sphere such that $f_i = g_i^{-1}$, $f_i(0) = 0 \in S_1$, $f_i'(0) = 1$, and $f_i(\infty) = \infty \in S_i$. Let g be the unique function mapping F onto the disk $\{|z| < R \leq \infty\}$ such that if $f = g^{-1}$, then $f(0) = 0 \in S_1$ and $f'(0) = 1$.

LEMMA 1. f_i is real for real z .

PROOF. The proof of this lemma is similar to the arguments of earlier authors [1], [4].

$g_i(F_i)$ is a simply connected domain of the Z -sphere with at least two boundary points (all points of $g_i(C_i)$ are boundary points). On removing the real axis from $g_i(F_i)$, the portion of $g_i(F_i)$ in the upper half plane is simply connected and open and, hence, can be mapped one-one and conformally onto a half disk $\{z: |z| < 1, \text{Im } z > 0\}$ in such a way that the real axis corresponds to a portion of the real axis and the points of $g_i(C_i)$ correspond to points of the semicircle. The symmetry principle applies to assure the existence of a function h_i such that $h_i(g_i(F_i))$ is a disk with center at 0 and points of the real axis correspond to points of the real diagonal of the disk. Furthermore, we

Received by the editors June 7, 1965.

can normalize h_i (call it h_i still) so that $h_i(0) = 0$ and $h'_i(0) = 1$. Then $h_i(g_i(F_i))$ is a disk of radius $r_i > 0$ about 0.

Notation. Let the origin on S_i be denoted by 0_i and the point at ∞ on S_i by ∞_i . Also let

$$\begin{aligned} \delta_{k,i} &= g_i(0_k), & \delta_k &= g(0_k), \\ \gamma_{k,i} &= g_i(\infty_k), & \gamma_k &= g(\infty_k), \\ \delta_{k,i}^* &= 1 - \frac{z}{\delta_{k,i}}, & \delta_k^* &= 1 - \frac{z}{\delta_k}, \\ \gamma_{k,i}^* &= 1 - \frac{z}{\gamma_{k,i}}, & \gamma_k^* &= 1 - \frac{z}{\gamma_k}. \end{aligned}$$

LEMMA 2.

$$R_i(z) = \frac{z}{\gamma_{1,i}^*} \prod_{k=2}^i \frac{\delta_{k,i}^*}{\gamma_{k,i}^*}$$

maps $\{|z| < r_i\}$ one-one and conformally onto F_i .

PROOF. F_i^* is a closed simply connected Riemann surface and is, therefore, the surface of the inverse of a unique rational function Q_i such that $Q_i(0) = 0_1$, $Q'_i(0) = 1$, and $Q_i(\infty) = \infty_i$. Composing Q_i with h_i^{-1} yields a unique rational function R_i mapping $\{|z| < r_i\}$ one-one and conformally onto F_i such that $R_i(0) = 0$, and $R'_i(0) = 1$.

LEMMA 3. $R_i(z) \rightarrow f(z)$ uniformly on compact subsets of $\{|z| < r\}$ for some $r > 0$.

PROOF. Let $D_i = \{|z| < r_i\}$. This result then follows immediately by applying MacLane's theorem [3, p. 33] with Corollary α to the two sequences of surfaces $\{F_i, 0\}$ and $\{D_i, 0\}$. We also get that $r_i \rightarrow r > 0$.

LEMMA 4. $\delta_{k,i} \rightarrow \delta_k$ and $\gamma_{k,i} \rightarrow \gamma_k$ as $i \rightarrow \infty$.

PROOF. This is a consequence of Hurewitz theorem or follows as one of the conclusions of the theorem of MacLane cited in the proof of Lemma 3.

LEMMA 5. *The infinite product*

$$\pi(z) = \frac{z}{\gamma_1^*} \prod_{k=2}^{\infty} \frac{\delta_k^*}{\gamma_k^*}$$

converges subuniformly on $\{|z| < r\}$.

PROOF. Let $r_0 \leq r$. Only a finite number of the δ_k or γ_k lie in $\{|z| < r_0\}$ and thus there exists $n_0 > 0$ such that for $k > n_0$, δ_k and γ_k lie in the ring $r_0 < |z| < r$. Since for each $k \in I$, $\delta_k > \gamma_k$, there exists $\epsilon_k > 0$ such that $\gamma_k + \epsilon_k = \delta_k$. Because of the location of γ_k and δ_k , $\sum_{k=1}^{\infty} \epsilon_k$ converges to some number less than $2r$. To prove the subuniform convergence of π , it is sufficient to prove the uniform convergence of

$$\sum_2^{\infty} \log \frac{\delta_k^*}{\gamma_k^*}$$

for $z \in \{|z| \leq r_0\}$. By the Cauchy criterion, this last series converges uniformly in $\{|z| \leq r_0\}$ provided that for each $\epsilon > 0$ and for all z with $|z| \leq r_0$, there exists a number $N(\epsilon) > 0$ such that for $n > N(\epsilon)$ and for all $p > 0$,

$$\left| \sum_{k=n_0+n}^{n_0+n+p} \log \frac{\delta_k^*}{\gamma_k^*} \right| < \epsilon.$$

Now

$$\begin{aligned} \left| \sum_{k=n_0+n}^{n_0+n+p} \log \frac{\delta_k^*}{\gamma_k^*} \right| &= \left| \sum_{m=1}^{\infty} \frac{z^m}{m} \sum_{k=n_0+n}^{n_0+n+p} \left(\frac{1}{\gamma_k^m} - \frac{1}{\delta_k^m} \right) \right| \\ &= \left| \sum_{m=1}^{\infty} \frac{z^m}{m} \sum_{k=n_0+n}^{n_0+n+p} \frac{\delta_k^m - \gamma_k^m}{\gamma_k^m \delta_k^m} \right| \\ &\leq \left| \sum_{m=1}^{\infty} \frac{z^m}{m \gamma_{n_0+n}^m} \sum_{k=n_0+n}^{n_0+n+p} 1 - \left(\frac{\gamma_k}{\delta_k} \right)^m \right|. \end{aligned}$$

If we sum this last expression first with respect to m and use the fact that $|z| < \gamma_{n_0+n}$, we get

$$\begin{aligned} \sum_{k=n_0+n}^{n_0+n+p} \frac{|z|}{\gamma_{n_0+n} - |z|} - \frac{|z| \gamma_k}{\gamma_{n_0+n} \delta_k - |z| \gamma_k} \\ = \sum_{k=n_0+n}^{n_0+n+p} \frac{|z| \gamma_{n_0+n} \left(\frac{\epsilon_k}{\gamma_k} \right)}{(\gamma_{n_0+n} - |z|) \left(\gamma_{n_0+n} \left(1 + \frac{\epsilon_k}{\gamma_k} \right) - |z| \right)} \\ \leq \frac{|z|}{(\gamma_{n_0+n} - |z|)^2} \sum_{k=n_0+n}^{n_0+n+p} \epsilon_k. \end{aligned}$$

Then if we choose $N(\epsilon)$ so large that

$$\sum_{k=N(\epsilon)}^{\infty} \epsilon_k < \frac{(\gamma_{n_0+1} - r_0)^2}{r_0} \epsilon,$$

the uniform convergence of the original series in $\{|z| \leq r_0\}$ is established.

LEMMA 6. $\pi(z) = f(z)$ for $z \in \{|z| < r\}$.

PROOF. Because of Lemma 4, there exists $\rho > 0$ such that $R_n(z)/z \neq 0$ and $\pi(z)/z \neq 0$ for $|z| < \rho$. For $m \geq 1$,

$$\begin{aligned} 0 &< \sum_{k=i}^{\infty} \left(\frac{1}{\gamma_k^m} - \frac{1}{\delta_k^m} \right) = \sum_{k=i}^{\infty} \frac{\delta_k^m - \gamma_k^m}{\gamma_k^m \delta_k^m} \\ &\leq \frac{1}{\gamma_i^m} \sum_{k=i}^{\infty} \left(1 - \left(\frac{\gamma_k}{\delta_k} \right)^m \right) \\ &= \frac{1}{\gamma_i^m} \sum_{k=i}^{\infty} \left[m \frac{\epsilon_k}{\delta_k} - \frac{m(m-1)}{2} \left(\frac{\epsilon_k}{\delta_k} \right)^2 + \dots \pm \left(\frac{\epsilon_k}{\delta_k} \right)^m \right] \\ &= \frac{1}{\gamma_i^m} \Phi(i) \end{aligned}$$

where $\lim_{i \rightarrow \infty} \Phi(i) = 0$. Similarly,

$$0 < \sum_{k=i}^n \left(\frac{1}{\gamma_{k,i}^m} - \frac{1}{\delta_{k,i}^m} \right) \leq \frac{1}{\gamma_{i,n}^m} \psi_n(i)$$

where $\lim_{i \rightarrow \infty} (\lim_{n \rightarrow \infty} \psi_n(i)) = \lim_{i \rightarrow \infty} \psi(i) = 0$. Therefore,

$$\begin{aligned} 0 &\leq \limsup_{n \rightarrow \infty} \left| \sum_{k=2}^n \left[\frac{1}{\gamma_{k,n}^m} - \frac{1}{\delta_{k,n}^m} \right] - \sum_{k=2}^{\infty} \left[\frac{1}{\gamma_k^m} - \frac{1}{\delta_k^m} \right] \right| \\ &\leq \limsup_{n \rightarrow \infty} \left| \sum_{k=n_0}^n \left[\frac{1}{\gamma_{k,n}^m} - \frac{1}{\delta_{k,n}^m} \right] - \sum_{k=n_0}^{\infty} \left[\frac{1}{\gamma_k^m} - \frac{1}{\delta_k^m} \right] \right| \\ &\leq \lim_{n \rightarrow \infty} \left(\frac{1}{\gamma_{n_0,n}^m} \psi_n(n_0) + \frac{1}{\gamma_{n_0}^m} \Phi(n_0) \right) \\ &= \frac{1}{\gamma_{n_0}^m} (\psi(n_0) + \Phi(n_0)). \end{aligned}$$

Since $\lim_{n_0 \rightarrow \infty} \psi(n_0) + \Phi(n_0) = 0$ and $\lim_{n_0 \rightarrow \infty} \gamma_{n_0}$ is $r > 0$,

$$\limsup_{n \rightarrow \infty} \left| \sum_{k=2}^n \left(\frac{1}{\gamma_{k,n}^m} - \frac{1}{\delta_{k,n}^m} \right) - \sum_{k=2}^{\infty} \left(\frac{1}{\gamma_k^m} - \frac{1}{\delta_k^m} \right) \right| = 0.$$

Thus all the coefficients of the Taylor series expansion of $\log (R_n(z)/\pi(z))$ about $z=0$ can be shown to have a limit zero as $n \rightarrow \infty$, and this $\lim_{n \rightarrow \infty} \log \{R_n(z)/\pi(z)\} = 0$ which implies that $\lim_{n \rightarrow \infty} R_n(z) = \pi(z) = f(z)$.

Remarks. It is clear that the specific form of the cuts was not essential in the construction of the uniformizing function. For this reason it is clear that the conditions on the cuts could be relaxed to permit variation in length and shape provided that they are symmetric, the resulting surface is hyperbolic, and the zeros and poles lie (except for a finite number of cases) on one side of the origin.

BIBLIOGRAPHY

1. H. B. Curtis, Jr., *Some properties of functions which uniformize a class of simply connected Riemann surfaces*, Proc. Amer. Math. Soc. **10** (1959), 525-530.
2. ———, *The uniformization of a class of simply connected Riemann surfaces*, Proc. Amer. Math. Soc. **11** (1960), 511-516.
3. G. R. MacLane, *Riemann surfaces and asymptotic values associated with real entire functions*, The Rice Institute Pamphlet, Houston, Texas, 1952.
4. H. E. Taylor, *Determination of the type and properties of the mapping functions of a class of doubly connected Riemann surfaces*, Proc. Amer. Math. Soc. **4** (1953), 52-68.

DEFENSE RESEARCH LABORATORY, THE UNIVERSITY OF TEXAS