

## IDEALS GENERATED BY PRODUCTS

A. P. HILLMAN, D. G. MEAD, K. B. O'KEEFE<sup>1</sup> AND E. S. O'KEEFE

In [1] Levi obtained results on the structure of the differential ideals  $[y^n]$  and  $[uv]$  and applied these results to the component theory of differential polynomials. The present paper uses Levi's methods to extend his main results on  $[uv]$  to  $[y_1 y_2 \cdots y_n]$ . Since the  $y_i$  are independent indeterminates,  $[y_1 \cdots y_n]$  is related to  $[y^n]$  but is not quite a generalization. The results are motivated by and apply to differential algebra; however, we follow Levi's suggestion at the close of [1] in stating them in the more general form in which  $y_{ij}$  is not necessarily a derivative of  $y_i$ .

Let  $y_{ij}$  ( $i=1, \cdots, n$  and  $j=0, 1, \cdots$ ) be a set of independent indeterminates over a field  $F$  and let  $R$  be the polynomial ring in the  $y_{ij}$  over  $F$ . The signature of a monomial  $M$  in  $R$  is  $D=(d_1, \cdots, d_n)$  if  $M$  has degree  $d_h$  in the  $y_{ij}$  with  $i=h$ . The weight of  $M$  is the sum of the  $j$ 's for the factors  $y_{ij}$  of  $M$ . A polynomial of  $R$  is homogeneous with signature  $D$  if each of its terms has this signature; it is isobaric of weight  $w$  if each term has weight  $w$ .

For  $j=0, 1, \cdots$  let  $x_j$  be a linear combination, with nonzero coefficients in  $F$ , of all the products  $y_{1j_1} y_{2j_2} \cdots y_{nj_n}$  of weight  $j$ . Let  $I_t$  be the ideal  $(x_0, x_1, \cdots, x_t)$  in  $R$ . Let  $I=(x_0, x_1, \cdots)$  and let  $Q$  be the quotient ring of  $R$  modulo  $I$ . Below we describe functions  $f(D)$  and  $g(D)$  such that a monomial with signature  $D$  and weight  $w$  is in  $I_t$  if  $w < f(D)$  and  $t \geq g(D)$  and such that for every  $D$  and  $w \geq f(D)$  there is a monomial with signature  $D$  and weight  $w$  that is not in  $I$ .

2. **Levi bases.** In the case  $n=2$ , Levi obtained the following bases for  $Q$  and  $R$  as vector spaces over  $F$ . Let  $u_j = y_{1j}$  and  $v_j = y_{2j}$ . A product

$$(1) \quad P = u_{i_1} \cdots u_{i_r} v_{j_1} \cdots v_{j_s}$$

$$(i_1 \leq i_2 \leq \cdots \leq i_r, j_1 \leq j_2 \leq \cdots \leq j_s)$$

of signature  $(r, s)$  is an  $\alpha$  term if  $s=0$  or  $j_1 \geq r$  and a  $\beta$  term otherwise. A  $\lambda$  term is of the form  $AX$  with  $A$  an  $\alpha$  term and  $X$  a power product in the  $x_j$ . ( $X$  may be 1; thus the  $\lambda$  terms include Levi's  $\alpha$  and  $\gamma$  terms.) Levi showed that the  $\alpha$  terms are a basis for  $Q$  and the  $\lambda$  terms for  $R$ .

Let  $P'$  be a factor of  $P$  in (1) selected as follows. If  $P$  is an  $\alpha$  term,

---

Received by the editors April 3, 1965.

<sup>1</sup> The work on this paper by one of the authors was supported by a grant from the American Association of University Women.

$P' = P$ . If  $P$  is a  $\beta$  term, let  $e = j_1 + 1$ ,  $S = u_i v_{j_1}$ , and  $P' = P/S$ . Let  $P_i$  be defined by  $P_0 = P$  and  $P_{i+1} = P'_i$ . Let  $\theta(P)$  be the expression for  $P$  in the form  $AS_1 \cdots S_a$  where  $a$  is the first  $i$  for which  $P_i$  is an  $\alpha$  term,  $A = P_a$ , and  $S_i = P_i/P_{i-1}$ . The weight  $h_i$  of  $S_i$  satisfies  $h_1 \leq h_2 \leq \cdots \leq h_a$ . Let  $\lambda(P) = Ax_{h_1} \cdots x_{h_a}$ . Levi showed that  $P \rightarrow \lambda(P)$  is a 1-1 mapping of the set of all power products in the  $u_j$  and  $v_j$  onto the set of all  $\lambda$  terms.

**3. Extension to general  $n$ .** Let  $P = Y_1 Y_2 \cdots Y_n$  where  $Y_k$  is a power product in the  $y_{ij}$  with  $i = k$ . For  $2 \leq k \leq n$  we inductively define  $\alpha_k$  terms in the  $y_{1j}, \cdots, y_{kj}$  and an expression  $\theta_k$  for  $Y_1 \cdots Y_k$ . When  $k = 2$ ,  $\alpha_2$  terms and  $\theta_2 = \theta(Y_1 Y_2)$  are defined as in the previous section with  $y_{1j}$  in the role of  $u_j$  and  $y_{2j}$  in that of  $v_j$ . We assume the definition of  $\alpha_m$  terms and of  $Y_1 \cdots Y_m$  in the form  $\theta_m = A_m S_{m1} \cdots S_{mb}$  where  $A_m$  is an  $\alpha_m$  term and  $S_{mj}$  has signature  $(1, \cdots, 1, 0, \cdots, 0)$ , with  $m$  ones, and weight  $h_{mj}$  satisfying  $h_{m1} \leq h_{m2} \leq \cdots \leq h_{mb}$ . Thinking of  $S_{mj}$  as  $u_j$  and  $y_{m+1,j}$  as  $v_j$ , let  $\theta(S_{m1} \cdots S_{mb} Y_{m+1}) = A^* S_{m+1,1} \cdots S_{m+1,c}$ . We then define  $\theta_{m+1}$  to be

$$(2) \quad A_{m+1} S_{m+1,1} \cdots S_{m+1,c}$$

where  $A_{m+1} = A_m A^*$ . The power product  $Y_1 \cdots Y_{m+1}$  is defined to be an  $\alpha_{m+1}$  term if  $c$  in (2) is zero, i.e., if  $Y_1 \cdots Y_{m+1} = A_{m+1}$ .

As before a  $\lambda$  term is of the form  $AX$  with  $A$  an  $\alpha$  (i.e.,  $\alpha_n$ ) term and  $X$  a power product in the  $x_j$ . Let  $\theta(P) = \theta_n = AS_1 \cdots S_e$  with  $A$  an  $\alpha$  term and  $S_j$  of weight  $h_j$  satisfying  $h_1 \leq \cdots \leq h_e$  and let  $\lambda(P) = Ax_{h_1} \cdots x_{h_e}$ . The following outline of the process of showing that the  $\alpha$  terms are a basis for  $Q$  and the  $\lambda$  terms for  $R$  is essentially as in Levi's work:

Since  $x_{h_1}$  is a linear combination of power products one of which is  $S_1$ ,  $(P/S_1)x_{h_1} \equiv 0 \pmod{I}$  can be solved as

$$(3) \quad P \equiv f_1 N_1 + \cdots + f_s N_s \pmod{I}$$

where the  $f_i$  are in  $F$  and the  $N_i$  are power products in the  $y_{ij}$ . Any  $N$  in (3) which is not an  $\alpha$  term can be replaced by an expression (3) in which  $N$  plays the role of  $P$ . Continuing the process, one ends in a finite number of steps with  $P$  congruent to a linear combination of  $\alpha$  terms of the same signature and weight as  $P$ . Thus the  $\alpha$  terms generate  $Q$ . It follows from the process of obtaining the  $\theta_k$  and the 1-1 character of  $P \rightarrow \lambda(P)$  in the case  $n = 2$  that, in the case of general  $n$ ,  $P \rightarrow \lambda(P)$  is a 1-1 mapping of the set of power products in the  $y_{ij}$  onto the set of  $\lambda$  terms. This establishes that the  $\lambda$  terms are a basis for  $R$ . Then the  $\alpha$  terms are linearly independent modulo  $I$  and form a basis for  $Q$ .

The function  $f(D)$  is the minimum of the weights of monomials with signature  $D$  that are not in  $I$ ; hence  $f(D)$  is the minimum of the weights of the  $\alpha$  terms with signature  $D$ . An  $\alpha$  term  $A$  is of the form  $A_{n-1}A^*$  where  $A_{n-1}$  is an  $\alpha_{n-1}$  term and  $A^*$  is an  $\alpha_2$  term in the  $S_{n-1,j}$  and the  $y_{nj}$ . If  $D = (d_1, \dots, d_n)$  and  $A^*$  is of degree  $t$  in the  $S$ 's, the weight of  $A$  is at least

$$(4) \quad f(d_1 - t, \dots, d_{n-1} - t) + td_n$$

and is (4) if  $A_{n-1}$  is an  $\alpha_{n-1}$  term of minimal weight for its signature and  $Y_n = (y_{ni})^{d_n}$ . Hence  $f(D)$  is the minimum of (4) for  $0 \leq t \leq \min(d_1, \dots, d_{n-1})$ .

There are other ways of defining a basis for  $Q$ . Although the basis may be different, the function  $f(D)$  does not depend on the basis. One such alternative shows that

$$(5) \quad f(D) = \min[f(d_1 - r, \dots, d_a - r) + f(d_{a+1} - s, \dots, d_b - s) + \dots + f(d_{c+1} - t, \dots, d_n - t) + f(r, s, \dots, t)]$$

where  $r, s, \dots, t$  range over all nonnegative integers such that the arguments in (5) are nonnegative.

It will be shown in another paper that  $f(D)$  may be evaluated explicitly as follows. We assume without loss of generality that  $d_1 \leq d_2 \leq \dots \leq d_n$ . For  $2 \leq i \leq n$  let  $q_i = (d_1 + \dots + d_i)/(i - 1)$  and let  $k$  be the smallest  $i$  for which  $q_i$  assumes its minimum. Let  $q$  and  $r$  be integers defined by  $d_1 + \dots + d_k = (k - 1)q + r$  and  $0 \leq r < k - 1$ . Let  $c_i = q - d_i$  for  $i = 1, \dots, k$ . Let  $\sigma_1 = c_1 + \dots + c_k$  and  $\sigma_2 = \sum_{i < j} c_i c_j$ . Then  $f(D) = \sigma_2 + r\sigma_1 + [(r + 1)r/2]$ .

It is easily seen from Levi's process that  $g(d_1, d_2)$  may be chosen as  $d_1 + d_2 - 2$ . This and the process described above show that  $g(d_1, \dots, d_n)$  may be chosen as

$$d_n + \min(d_1, \dots, d_{n-1}) - 2.$$

Using the symmetry of the ideal  $I_t$  in the subscripts  $i$  of the  $y_{ij}$ , this may be improved to  $g(D) = \min_{i \neq j} (d_i + d_j) - 2$ .

REFERENCES

1. Howard Levi, *On the structure of differential polynomials and on their theory of ideals*, Trans. Amer. Math. Soc. 51 (1942), 532-568.
2. A. P. Hillman, D. G. Mead, K. B. O'Keefe and E. S. O'Keefe, *A dynamic programming generalization of xy to n variables*, Proc. Amer. Math. Soc. 17 (1966), 718-721.

UNIVERSITY OF SANTA CLARA AND  
UNIVERSITY OF WASHINGTON