

A COMPARISON THEOREM FOR ELLIPTIC DIFFERENTIAL EQUATIONS

C. A. SWANSON¹

Clark and the author [2] recently obtained a generalization of the Hartman-Wintner comparison theorem [4] for a pair of self-adjoint second order linear elliptic differential equations. The purpose of this note is to extend this generalization to general second order linear elliptic equations. As in [2], the usual pointwise inequalities for the coefficients are replaced by a more general integral inequality. The result is new even in the one-dimensional case, and extends Leighton's result for self-adjoint ordinary equations [5].

Protter [6] obtained pointwise inequalities in the nonself-adjoint case in two dimensions by the method of Hartman and Wintner [4]. We obtain an alternative to Protter's result as a corollary of our main theorem.

Let R be a bounded domain in n -dimensional Euclidean space with boundary B having a piecewise continuous unit normal. The linear elliptic differential operator L defined by

$$(1) \quad Lu = \sum_{i,j=1}^n D_i(a_{ij}D_ju) + 2 \sum_{i=1}^n b_i D_iu + cu, \quad a_{ij} = a_{ji}$$

will be considered in R , where D_i denotes partial differentiation with respect to x^i , $i=1, 2, \dots, n$. We assume that the coefficients a_{ij} , b_i , and c are real and continuous on \bar{R} , the b_i are differentiable in R , and that the symmetric matrix (a_{ij}) is positive definite in R . A "solution" u of $Lu=0$ is supposed to be continuous on \bar{R} and have uniformly continuous first partial derivatives in R , and all partial derivatives involved in (1) are supposed to exist, be continuous, and satisfy $Lu=0$ in R .

Let $Q[z]$ be the quadratic form in $(n+1)$ variables z_1, z_2, \dots, z_{n+1} defined by

$$(2) \quad Q[z] = \sum_{i,j=1}^n a_{ij}z_i z_j - 2z_{n+1} \sum_{i=1}^n b_i z_i + g z_{n+1}^2,$$

where the continuous function g is to be determined so that this form is positive semidefinite. The matrix Q associated with $Q[z]$ has the block form

Received by the editors October 21, 1965.

¹ This research was supported by the United States Air Force Office of Scientific Research, under grant AF-AFOSR-379-65.

$$Q = \begin{pmatrix} A & -b \\ -b^T & g \end{pmatrix}, \quad A = (a_{ij}),$$

where b^T is the n -vector (b_1, b_2, \dots, b_n) . Let B_i denote the cofactor of $-b_i$ in Q . Since A is positive definite, a necessary and sufficient condition for Q to be positive semidefinite is $\det Q \geq 0$, or

$$(3) \quad g \det (a_{ij}) \geq - \sum_{i=1}^n b_i B_i.$$

The proof is a slight modification of the well-known proof for positive definite matrices [3, p. 306].

Let J be the quadratic functional defined by

$$(4) \quad J[u] = \int_R F[u] dx$$

where

$$F[u] = \sum_{i,j} a_{ij} D_i u D_j u - 2u \sum_i b_i D_i u + (g - c)u^2,$$

with domain \mathfrak{D} consisting of all real-valued continuous functions on \bar{R} which have uniformly continuous first partial derivatives in R and vanish on B .

LEMMA. *Suppose g satisfies (3). If there exists $u \in \mathfrak{D}$ not identically zero such that $J[u] < 0$, then every solution v of $Lv = 0$ vanishes at some point of \bar{R} .*

PROOF. Suppose to the contrary that there exists a solution $v \neq 0$ in \bar{R} . For $u \in \mathfrak{D}$ define

$$X^i = v D_i (u/v);$$

$$Y^i = v^{-1} \sum_j a_{ij} D_j v, \quad i = 1, 2, \dots, n;$$

$$E[u, v] = \sum_{i,j} a_{ij} X^i X^j - 2u \sum_i b_i X^i + gu^2 + \sum_i D_i (u^2 Y^i).$$

A routine calculation yields the identity

$$E[u, v] = F[u] + u^2 v^{-1} Lv.$$

Since $Lv = 0$ in R ,

$$\begin{aligned}
 (5) \quad J[u] &= \int_R \left[\sum_{i,j} a_{ij} X^i X^j - 2u \sum_i b_i X^i + gu^2 \right] dx \\
 &+ \int_R \sum_i D_i(u^2 Y^i) dx.
 \end{aligned}$$

Since $u=0$ on B , the second integral is zero by Green's formula. The first integrand is a positive semidefinite form by hypothesis (3). The contradiction $J[u] \geq 0$ establishes the lemma.

Consider in addition to (1) a second differential operator L^* of the same form,

$$L^*u = \sum_{i,j=1}^n D_i(a_{ij}^* D_j u) + 2 \sum_i b_i^* D_i u + c^* u, \quad a_{ij}^* = a_{ji}^*$$

in which the coefficients satisfy the same conditions as the coefficients in (1). L^* is the Euler-Jacobi operator associated with the quadratic functional J^* defined by

$$(6) \quad J^*[u] = \int_R \left[\sum_{i,j} a_{ij}^* D_i u D_j u - 2u \sum_i b_i^* D_i u - c^* u^2 \right] dx.$$

Define $V[u] = J^*[u] - J[u]$, $u \in \mathfrak{D}$. Since $u=0$ on B , it follows from partial integration that

$$\begin{aligned}
 (7) \quad V[u] &= \int_R \left[\sum (a_{ij}^* - a_{ij}) D_i u D_j u \right. \\
 &\quad \left. + \left\{ \sum D_i(b_i^* - b_i) + c - c^* - g \right\} u^2 \right] dx.
 \end{aligned}$$

THEOREM 1. *Suppose g satisfies (3). If there exists a nontrivial solution u of $L^*u=0$ in R such that $u=0$ on B and $V[u]>0$, then every solution of $Lv=0$ vanishes at some point of \bar{R} .*

PROOF. The hypothesis $V[u]>0$ is equivalent to $J[u]<J^*[u]$. Since $u=0$ on B , it follows from Green's formula that $J^*[u]=0$. Hence the hypothesis $J[u]<0$ of the lemma is fulfilled.

THEOREM 2. *Suppose $g \det(a_{ij}) > -\sum b_i B_i$. If there exists a nontrivial solution of $L^*u=0$ in R such that $u=0$ on B and $V[u] \geq 0$, then every solution of $Lv=0$ vanishes at some point of \bar{R} .*

Since Q is positive definite, the lemma is valid when the hypothesis $J[u]<0$ is replaced by $J[u] \leq 0$. The proof of Theorem 2 is then analogous to that of Theorem 1.

In the case that equality holds in (3), that is

$$(8) \quad g = - \sum_i b_i B_i / \det (a_{ij}),$$

define

$$\delta = \sum D_i(b_i^* - b_i) + c - c^* - g.$$

L is called a “strict Sturmian majorant” of L^* by Hartman and Wintner [4] when the following conditions hold: (i) $(a_{ij}^* - a_{ij})$ is positive semidefinite and $\delta \geq 0$ in \bar{R} ; (ii) either $\delta > 0$ at some point or $(a_{ij}^* - a_{ij})$ is positive definite and $c^* \neq 0$ at some point. The corollary below follows immediately from Theorem 1.

COROLLARY. *Suppose that L is a strict Sturmian majorant of L^* . If there exists a solution u of $L^*u = 0$ in R such that $u = 0$ on B and u does not vanish in any open set contained in R , then every solution of $Lv = 0$ vanishes at some point of \bar{R} .*

If the coefficients a_{ij}^* are of class $C^{2,1}(R)$ (i.e. all second derivatives exist and are Lipschitzian), the hypothesis that u does not vanish in any open set of R can be replaced by the condition that u does not vanish identically in R because of Aronszajn’s unique continuation theorem [1].

In the case $n = 2$ considered by Protter [6], the condition $\delta \geq 0$ reduces to

$$(9) \quad (a_{11}a_{22} - a_{12}^2) \left(\sum_{i=1}^2 D_i(b_i^* - b_i) + c - c^* \right) \geq a_{11}b_2^2 - 2a_{12}b_1b_2 + a_{22}b_1^2,$$

which is considerably simpler than Protter’s condition. It reduces to Protter’s condition

$$\sum_{i=1}^2 D_i b_i^* + c - c^* \geq 0$$

in the case that $b_1 = b_2 = 0$, and also in the case that $a_{12} = a_{12}^* = 0$, $a_{11} = a_{11}^*$, $a_{22} = a_{22}^*$. (Two incorrect signs appear in [6]).

The following example in the case $n = 2$ illustrates that Theorem 1 is more general than the pointwise condition (9). Let R be the square $0 < x^1, x^2 < \pi$. Let L^* , L be the elliptic operators defined by

$$L^*u = D_1^2u + D_2^2u + 2u,$$

$$Lv = D_1^2v + D_2^2v + D_1v + cv,$$

where

$$c(x^1, x^2) = f(x^1)f(x^2) + 5/4,$$

and $f \in C[0, \pi]$ is not identically zero. The function $u = \sin x^1 \sin x^2$ is zero on B and satisfies $L^*u = 0$. The condition $V[u] > 0$ of Theorem 1 reduces to

$$\int_0^\pi \int_0^\pi f(x^1)f(x^2) \sin^2 x^1 \sin^2 x^2 dx^1 dx^2 > 0.$$

Since this is fulfilled, every solution of $Lv = 0$ vanishes at some point of \bar{R} . This cannot be concluded from (9) or from Protter's result [6] unless f has constant sign in R .

In the case $n = 1$, L is an ordinary differential operator of the form

$$Lu = (au')' + 2bu' + cu,$$

and R is an interval (x_1, x_2) . We assert that \bar{R} can be replaced by R in the lemma and theorems; for v can have at most a simple zero at the boundary points x_1 and x_2 , and hence the first integral on the right side of (5) exists and is nonnegative provided only that $v \neq 0$ in R .

THEOREM 3. *If there exists a nontrivial solution u of $L^*u = 0$ in R such that $u = 0$ on B and*

$$(10) \quad \int_{x_1}^{x_2} \left[(a^* - a)u'^2 + \left(b^* - b' + c - c^* - \frac{b^2}{a} \right) u^2 \right] dx > 0,$$

then every solution of $Lv = 0$ has a zero in (x_1, x_2) .

In the self-adjoint case $b = b^* = 0$ it was shown by Clark and the author [2] that the strict inequality in the hypothesis $V[u] > 0$ of Theorem 1, and therefore also in (10), can be replaced by \geq . Indeed, this is transparent when the proof of the above lemma is specialized to the self-adjoint case. With $>$ replaced by \geq , (10) reduces to Leighton's condition in the self-adjoint case [5].

REFERENCES

1. N. Aronszajn, *A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order*, J. Math. Pures Appl. **36** (1957), 235-249.

2. Colin Clark and C. A. Swanson, *Comparison theorems for elliptic differential equations*, Proc. Amer. Math. Soc. **16** (1965), 886–890.
3. F. R. Gantmacher, *The theory of matrices*, Vol. I, Chelsea, New York, 1959.
4. Philip Hartman and Aurel Wintner, *On a comparison theorem for self-adjoint partial differential equations of elliptic type*, Proc. Amer. Math. Soc. **6** (1955), 862–865.
5. Walter Leighton, *Comparison theorems for linear differential equations of second order*, Proc. Amer. Math. Soc. **13** (1962), 603–610.
6. M. H. Protter, *A comparison theorem for elliptic equations*, Proc. Amer. Math. Soc. **10** (1959), 296–299.

THE UNIVERSITY OF BRITISH COLUMBIA

q -ANALOGUES OF CAUCHY'S FORMULAS

WALEED A. AL-SALAM

1. Let q be a given number and let α be real or complex. The α th “basic number” is defined by means of $[\alpha] = (1 - q^\alpha)/(1 - q)$. This has served as a basis for an extensive amount of literature in mathematics under such titles as Heine, basic, or q -series and functions. The basic numbers also occur naturally in many theta identities. The works of Jackson (for bibliography see [2]) and Hahn [3] have stimulated much interest in this field.

One important operation that is intimately connected with basic series as well as with difference and other functional equations is the q -derivative of a function f . This is defined by

$$(1.1) \quad Df(x) = \frac{f(qx) - f(x)}{x(q - 1)}.$$

Jackson defined the operations, which he called q -integration,

$$(1.2) \quad \int_0^x f(t) d(q, t) = x(1 - q) \sum_{k=0}^{\infty} q^k f(xq^k)$$

and

$$(1.3) \quad \int_x^{\infty} f(t) d(q, t) = x(1 - q) \sum_{k=0}^{\infty} q^{-k} f(xq^{-k})$$

Received by the editors October 28, 1965 and, in revised form, November 30, 1965.