

# ON TOTALLY PARACOMPACT METRIC SPACES<sup>1</sup>

A. LELEK

A topological space is said to be *totally paracompact* [4] if every open basis contains a locally finite cover. It is known [2] that no reflexive infinite-dimensional Banach space is totally paracompact. It is also known [1], [3] that the space of irrationals with usual topology is not totally paracompact. In the present paper we prove a theorem which gives a necessary condition in order that a subset of a complete metric space be totally paracompact. This allows us to exhibit some pathology (Corollary 2) among the separable metric spaces which are not totally paracompact.

Let  $\gamma$  be a collection of sets. We denote by  $|\gamma|$  the union of elements of  $\gamma$ . The collection  $\gamma$  is called *point-finite* if no point belongs to infinitely many elements of  $\gamma$ . By a *Cantor set* we mean a set homeomorphic with the Cantor ternary set.

**LEMMA.** *Let  $X$  be a complete metric space and  $Y \subset X \setminus Z$  where  $Z$  is a nonempty subset of  $X$  such that no point of  $Z$  is isolated in  $Z$ . There exists an open basis  $\beta$  of  $Y$  such that if  $\gamma \subset \beta$  is point-finite, then  $X \setminus |\gamma|$  contains a Cantor set.*

**PROOF.** We take an increasing infinite sequence  $A_1 \subset A_2 \subset \dots$  of nonempty countable compact subsets  $A_n$  of  $Z$  such that no point of  $A_n$  is isolated in  $A_{n+1}$  for  $n = 1, 2, \dots$ . Let  $U_{n1} \supset U_{n2} \supset \dots$  be open subsets of  $X$  such that

$$A_n = U_{n1} \cap U_{n2} \cap \dots = \text{Cl } U_{n1} \cap \text{Cl } U_{n2} \cap \dots$$

and let  $G_n(1), \dots, G_n(k_n)$  be open subsets of  $X$  such that

$$A_n \subset G_n(1) \cup \dots \cup G_n(k_n), \quad \text{diam } G_n(i) < n^{-1}$$

for  $i = 1, \dots, k_n$ . Let  $V_n$  be an open subset of  $X$  such that

$$A_n \subset V_n, \quad \text{Cl } V_n \subset G_n(1) \cup \dots \cup G_n(k_n)$$

and let  $\beta'_n$  denote the collection of all open subsets of  $Y \setminus \text{Cl } V_n$ .

We put

$$\beta_n = \beta'_n \cup \{ [G_n(i) \setminus \text{Cl } U_{nj}] \cap Y \mid i = 1, \dots, k_n; j = 1, 2, \dots \}$$

and we define  $\beta = \beta_1 \cup \beta_2 \cup \dots$ . Since  $Y \subset X \setminus A_n$ , each point from

Received by the editors September 23, 1966.

<sup>1</sup> This research had partially been done when the author was visiting the Louisiana State University and the University of Wisconsin.

$Y \cap \text{Cl} V_n$  belongs to a set  $G_n(i) \setminus \text{Cl} U_{nj}$  whose diameter is less than  $n^{-1}$ . Therefore  $\beta$  is an open basis of  $Y$ .

Consider an arbitrary point-finite collection  $\gamma \subset \beta$ . The collection  $\beta_n \setminus \beta'_n$  splits into  $k_n$  increasing infinite sequences. Setting  $\gamma_n = \beta_n \cap \gamma$  we conclude that  $A_n \subset X \setminus \text{Cl} |\gamma_n|$  for  $n = 1, 2, \dots$ . No point of  $A_1$  is isolated in  $A_2$ , and so there exist two points  $p_0, p_1$  of  $A_2$  belonging to  $X \setminus \text{Cl} |\gamma_1|$ . We take an open neighborhood  $W(m)$  of  $p_m$  ( $m = 0, 1$ ) such that

$$W(m) \subset X \setminus \text{Cl} |\gamma_1|, \quad \text{Cl} W(0) \cap \text{Cl} W(1) = \emptyset,$$

and  $\text{diam } W(m) < 1$ . Let us assume  $n > 1$  is an integer, and an open neighborhood  $W = W(m_1, \dots, m_{n-1})$  of a point from  $A_n$  is defined, where the indices  $m_1, \dots, m_{n-1}$  are 0 or 1. Then there exist two points  $q_0, q_1$  of  $A_{n+1}$  belonging to  $W \setminus \text{Cl} |\gamma_n|$ . We take an open neighborhood  $W(m_1, \dots, m_{n-1}, m)$  of  $q_m$  ( $m = 0, 1$ ) such that

$$\begin{aligned} \text{Cl } W(m_1, \dots, m_{n-1}, m) &\subset W(m_1, \dots, m_{n-1}) \setminus \text{Cl} |\gamma_n|, \\ \text{Cl } W(m_1, \dots, m_{n-1}, 0) \cap \text{Cl } W(m_1, \dots, m_{n-1}, 1) &= \emptyset, \end{aligned}$$

and  $\text{diam } W(m_1, \dots, m_{n-1}, m) < n^{-1}$ . The intersection

$$C = \bigcap_{n=1}^{\infty} \bigcup_{(m_1, \dots, m_n)} \text{Cl } W(m_1, \dots, m_n)$$

is a Cantor set, and  $C \subset X \setminus (|\gamma_1| \cup |\gamma_2| \cup \dots) = X \setminus |\gamma|$ .

**THEOREM.** *Let  $X$  be a complete metric space and  $Y \subset X$ . If  $Y$  is totally paracompact and  $p \in X \setminus Y$ , then either  $X \setminus Y$  contains a Cantor set containing  $p$  or each subset of  $X \setminus Y$  containing  $p$  has an isolated point.*

**PROOF.** Suppose there exists a set  $Z \subset X \setminus Y$  such that  $p \in Z$  and no point of  $Z$  is isolated in  $Z$ . Applying the lemma to the sets

$$\begin{aligned} X_n &= \{x \mid \text{dist}(p, x) \leq n^{-1}\}, \\ Y_n &= X_n \cap Y, \quad Z_n = \text{Int } X_n \cap Z, \end{aligned}$$

we get a Cantor set  $C_n \subset X_n \setminus Y_n$  because  $Y_n$  is totally paracompact ( $n = 1, 2, \dots$ ). The union  $\{p\} \cup C_1 \cup C_2 \cup \dots$  is a Cantor set contained in  $X \setminus Y$ .

**COROLLARY 1.** *Let  $X$  be a complete separable metric space and  $Y \subset X$ . If  $Y$  is uncountable and  $Y$  contains no Cantor set, then  $X \setminus Y$  is not totally paracompact.*

Each closed subset of a totally paracompact space is totally paracompact. Thus if  $X$  is totally paracompact, no closed subset of  $X$  is

homeomorphic with the irrationals. Answering a question raised by E. Michael we prove the inverse implication is not true. Namely, as a result of Bernstein's construction [5, p. 422] which uses the axiom of choice one can obtain a subset  $X$  of the Euclidean  $(n+1)$ -space  $E^{n+1}$  such that both  $X$  and  $E^{n+1} \setminus X$  have the cardinality of the continuum and contain no Cantor sets.

**COROLLARY 2.** *If the axiom of choice is true, then there exists, for every  $n \geq 0$ , an  $n$ -dimensional subset  $X$  of the Euclidean  $(n+1)$ -space such that  $X$  is not totally paracompact and  $X$  contains no Cantor set.*

Observe that an analogous singularity exists among totally paracompact spaces. In fact, as a result of Lusin's construction [5, pp. 432–433] using the axiom of choice and the continuum hypothesis one gets an uncountable subset  $X$  of the irrationals such that  $X$  is totally paracompact and  $X$  contains no Cantor set.

#### BIBLIOGRAPHY

1. A. V. Arhangel'skiĭ, *On the metrization of topological spaces*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **8** (1960), 589–595.
2. H. H. Corson, *Collections of convex sets which cover a Banach space*, Fund. Math. **49** (1961), 143–145.
3. H. H. Corson, T. J. McMinn, E. A. Michael, and J.-I. Nagata, *Bases and local finiteness*, Notices Amer. Math. Soc. **6** (1959), 814.
4. R. M. Ford, *Basis properties in dimension theory*, Doctoral Dissertation, Auburn University, Auburn, Ala., 1963.
5. C. Kuratowski, *Topologie*. I, Państwowe Wydawnictwo Naukowe, Warsaw, 1958.

INSTITUTE OF MATHEMATICS OF THE POLISH ACADEMY OF SCIENCES