

AN EQUIVALENCE THEOREM FOR EMBEDDINGS OF COMPACT ABSOLUTE NEIGHBORHOOD RETRACTS

J. L. BRYANT AND C. L. SEEBECK III¹

In this paper we wish to prove the following theorem.

THEOREM 1. *Suppose that each of A_0 and A_1 is a compact absolute neighborhood retract (ANR) of dimension k in Euclidean n -space E^n ($2k+2 \leq n$, $n \geq 5$) such that $E^n - A_i$ is 1-ULC (uniformly locally simply connected) for $i=0, 1$, and $f: A_0 \rightarrow A_1$ is a homeomorphism such that $d(a, f(a)) < \epsilon$ for each $a \in A_0$. Then there exists an ϵ -push h of (E^n, A^0) such that $h|_{A_0} = f$.*

In [2] the authors showed that if A is a k -dimensional polyhedron topologically embedded in E^n ($2k+2 \leq n$, $n \geq 5$) such that $E^n - A$ is 1-ULC, then for each $\epsilon > 0$, there is an ϵ -push h of (E^n, A) such that $h|_A: A \rightarrow E^n$ is piecewise linear. Hence, a well-known theorem of Bing and Kister [1, Theorem 5.5] applies to prove Theorem 1 when A_0 is a polyhedron. In fact, Theorem 5.5 of [1], together with the techniques of Homma [4] and Gluck [3] and the following engulfing theorem proved in [2], make our result possible.

THEOREM 2. *Suppose that A is a k -dimensional compact ANR in E^n ($n-k \geq 3$, $n \geq 5$) such that $E^n - A$ is 1-ULC and $\epsilon > 0$. Then there exists $\delta > 0$ such that if $f: A \rightarrow E^n$ is a δ -homeomorphism and U is an open subset of E^n containing $f(A)$, then there exists an ϵ -push h of (E^n, A) such that $h(U) \supset A$.*

Following Gluck [3], we define an ϵ -push h of the pair (X, A) , where X is a metric space and A is a subset of X such that \bar{A} is compact, to be a homeomorphism of X onto itself that is ϵ -isotopic to the identity by an isotopy h_t ($t \in [0, 1]$) of X such that for each $t \in [0, 1]$, $h_t|_{X - N_\epsilon(A)} = \text{identity}$. Other terminology used here is standard, and we shall assume that it is familiar to the reader.

Actually, the proof of Theorem 1 follows from known results, once we prove a

LEMMA. *Suppose that A is a compact ANR of dimension k in E^n ($2k+2 \leq n$, $n \geq 5$) such that $E^n - A$ is 1-ULC and $f: A \rightarrow E^n$ is an embedding such that $d(a, f(a)) < \epsilon$ for each $a \in A$. Then for each $\delta > 0$ there exists an ϵ -push h of (E^n, A) such that $d(h(a), f(a)) < \delta$ for each $a \in A$.*

Received by the editors September 12, 1967.

¹ This research was supported in part by National Science Foundation Grant GP 5458.

Theorem 1 is then proved exactly as is Theorem 4.4 of [3], and we shall not repeat the details of the constructions involved.

PROOF OF THE LEMMA. Given $\delta > 0$, there exists $\eta > 0$ such that if $a, b \in A$ with $d(a, b) < \eta$, then $d(f(a), f(b)) < \frac{1}{2}\delta$. Let N be a polyhedral neighborhood of A in E^n that retracts onto A by a retraction $r: N \rightarrow A$ such that $d(x, r(x)) < \frac{1}{2}\eta$ and $d(x, fr(x)) < \epsilon$ for each $x \in N$. Let T be a triangulation of N with mesh less than $\frac{1}{4}\eta$ and let T^k denote the k -skeleton of T , with $N^k = |T^k|$, the polyhedron of T^k .

The mapping $fr: N^k \rightarrow E^n$ has the property that if $x \in N^k$, $a \in A$, and $d(x, a) < \frac{1}{2}\eta$, then $d(fr(x), f(a)) < \frac{1}{2}\delta$, since $d(r(x), a) < \eta$.

Let $g': N^k \rightarrow E^n$ be a piecewise linear embedding such that $d(g'(x), fr(x)) < \frac{1}{2}\delta$ and $d(x, g'(x)) < \epsilon$ for each $x \in N^k$. Since $2k+2 \leq n$, we may apply Theorem 5.5 of [1] to obtain an ϵ -push g of (E^n, N^k) such that $g|N^k = g'$. Notice that if $x \in N^k$, $a \in A$, and $d(x, a) < \frac{1}{2}\eta$, then $d(g(x), f(a)) \leq d(g(x), fr(x)) + d(fr(x), f(a)) < \delta$. Thus there is an open set U in E^n containing N^k such that the above implication is true for $x \in U$; that is, if $x \in U$, $a \in A$, and $d(x, a) < \frac{1}{2}\eta$, then $d(g(x), f(a)) < \delta$. We need one additional fact concerning the open set U .

SUBLEMMA. *There exists a $\frac{1}{2}\eta$ -push ϕ of (E^n, A) such that $\phi(A) \subset U$.*

PROOF. Let \tilde{T}^{n-k-1} be the dual $(n-k-1)$ -skeleton of T with $\tilde{N}^{n-k-1} = |\tilde{T}^{n-k-1}|$. Choose $\eta' > 0$ corresponding to $\frac{1}{4}\eta$ as in Theorem 2. From the construction in the proof of Theorem V5 of [5], we can obtain an embedding ψ of A into E^n such that $d(a, \psi(a)) < \eta'$ for each $a \in A$ and $\psi(A) \cap \tilde{N}^{n-k-1} = \emptyset$. Let V be an open subset of E^n containing $\psi(A)$ such that $V \cap \tilde{N}^{n-k-1} = \emptyset$.

By Theorem 2, there exists a $\frac{1}{4}\eta$ -push ϕ_1 of (E^n, A) such that $\phi_1(V) \supset A$. Then ϕ_1^{-1} is a $\frac{1}{4}\eta$ -push of (E^n, A) and $\phi_1^{-1}(A) \cap \tilde{N}^{n-k-1} = \emptyset$. Since the mesh of T is less than $\frac{1}{4}\eta$, the technique of Stallings [7] may be used to obtain a $\frac{1}{4}\eta$ -push ϕ_2 of $(E^n, \phi_1^{-1}(A))$ such that $\phi_2\phi_1^{-1}(A) \subset U$. Then $\phi = \phi_2\phi_1^{-1}$ is the desired $\frac{1}{2}\eta$ -push of (E^n, A) .

We complete the proof of the Lemma by setting $h = g\phi$. We may assume that η is chosen sufficiently small so that the composition $g\phi$ is an ϵ -push of (E^n, A) . Given $a \in A$, we have $d(\phi(a), a) < \frac{1}{2}\eta$ and $\phi(a) \in U$, so that $d(g\phi(A), f(a)) = d(h(a), f(a)) < \delta$.

The question as to whether Theorem 1 is true when $k=1$ and $n=4$ seems very hard to answer. The method used to prove Theorem 1 involves engulfing techniques that are valid only for $n \geq 5$. It might be possible, however, to improve the codimension restriction by one if certain other conditions are satisfied. For example, Price has shown [6] that any two piecewise linear embeddings of a k -complex K into E^n ($n = 2k+1$) are equivalent by an isotopy of E^n that is the identity outside a compact set if $H^k(K, Z) = 0$. A natural question then is

QUESTION 1. Is Theorem 1 true with $n = 2k + 1$ if $H^k(A_0, Z) = 0$? In particular, one might consider a special case.

QUESTION 2. Is Theorem 1 true with $n = 2k + 1$ if A_0 is an absolute retract?

REFERENCES

1. R. H. Bing and J. M. Kister, *Taming complexes in hyperplanes*, Duke Math. J. **31** (1964), 491–511.
2. J. L. Bryant and C. L. Seebeck III, *Locally nice embeddings of polyhedra*, Quart. J. Math. Oxford Ser. (2) **19** (1968), 257–274.
3. H. Gluck, *Embeddings in the trivial range*, Ann. of Math. (2) **81** (1965), 195–210.
4. T. Homma, *On the imbedding of polyhedra in manifolds*, Yokohama Math. J. **10** (1962), 5–10.
5. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Univ. Press, Princeton, N. J., 1948.
6. T. M. Price, *Equivalence of embeddings of k -complexes in E^n for $n \leq 2k + 1$* , Michigan Math. J. **13** (1966), 65–69.
7. J. Stallings, *On topologically unknotted spheres*, Ann. of Math. **77** (1963), 490–502.

FLORIDA STATE UNIVERSITY