

## A NEW PROOF THAT METRIC SPACES ARE PARACOMPACT

MARY ELLEN RUDIN

By using a well-ordered open cover, there is a simple proof of the nice theorem [1] that every metric space is paracompact.

Assume that  $X$  is a metric space and that  $\{C_\alpha\}$  is an open cover of  $X$  indexed by ordinals. Let  $\rho$  be a metric on  $X$  and let  $S(x, r)$  be the open sphere with center  $x$  and radius  $r$ . For each positive integer  $n$  define  $D_{an}$  (by induction on  $n$ ) to be the union of all spheres  $S(x, 2^{-n})$  such that:

- (1)  $\alpha$  is the smallest ordinal with  $x \in C_\alpha$ ,
- (2)  $x \notin D_{\beta j}$  if  $j < n$ ,
- (3)  $S(x, 3 \cdot 2^{-n}) \subset C_\alpha$ .

Then  $\{D_{an}\}$  is a locally finite refinement of  $\{C_\alpha\}$  which covers  $X$ ; hence  $X$  is paracompact.

Certainly  $\{D_{an}\}$  refines  $\{C_\alpha\}$ . To see that  $\{D_{an}\}$  covers  $X$ , observe that, for  $x \in X$ , there is a smallest ordinal  $\alpha$  such that  $x \in C_\alpha$ , and an  $n$  so large that (3) holds. Then, by (2),  $x \in D_{\beta j}$  for some  $j \leq n$ .

To prove that  $\{D_{an}\}$  is locally finite, assume an  $x \in X$  and let  $\alpha$  be the smallest ordinal such that  $x \in D_{an}$  for some  $n$ , and choose  $j$  so that  $S(x, 2^{-j}) \subset D_{an}$ . The proof consists of showing that:

- (a) if  $i \geq n + j$ ,  $S(x, 2^{-n-j})$  intersects no  $D_{\beta i}$ ,
- (b) if  $i < n + j$ ,  $S(x, 2^{-n-j})$  intersects  $D_{\beta i}$  for at most one  $\beta$ .

PROOF OF (a). Since  $i > n$ , by (2), every one of the spheres of radius  $2^{-i}$  used in the definition of  $D_{\beta i}$  has its center  $y$  outside of  $D_{an}$ . And since  $S(x, 2^{-j}) \subset D_{an}$ ,  $\rho(x, y) \geq 2^{-j}$ . But  $i \geq j + 1$  and  $n + j \geq j + 1$ , so  $S(x, 2^{-n-j}) \cap S(y, 2^{-i}) = \emptyset$ .

PROOF OF (b). Suppose  $p \in D_{\beta i}$ ,  $q \in D_{\gamma i}$ , and  $\beta < \gamma$ ; we want to show that  $\rho(p, q) > 2^{-n-j+1}$ . There are points  $y$  and  $z$  such that  $p \in S(y, 2^{-i}) \subset D_{\beta i}$ ,  $q \in S(z, 2^{-i}) \subset D_{\gamma i}$ ; and, by (3),  $S(y, 3 \cdot 2^{-i}) \subset C_\beta$  but, by (2),  $z \notin C_\beta$ . So  $\rho(y, z) \geq 3 \cdot 2^{-i}$  and  $\rho(p, q) > 2^{-i} \geq 2^{-n-j+1}$ .

### BIBLIOGRAPHY

1. A. H. Stone, *Paracompactness and product spaces*, Bull. Amer. Math. Soc. 54 (1948), 977-982.

UNIVERSITY OF WISCONSIN

Received by the editors January 30, 1968.