

AN IDENTITY FOR THE SCHUR COMPLEMENT OF A MATRIX

DOUGLAS E. CRABTREE AND EMILIE V. HAYNSWORTH¹

1. Introduction. Let $A = (a_{ij})$ be an $n \times n$ complex matrix, and suppose that B is a nonsingular principal submatrix of A . We define the *Schur complement of B in A* , denoted by (A/B) , as follows: Let \hat{A} be the matrix obtained from A by a simultaneous permutation of rows and columns which puts B into the upper left corner of \hat{A} .

$$A = \begin{bmatrix} B & E \\ D & G \end{bmatrix}.$$

Then $(A/B) = G - DB^{-1}E$.

Since

$$\begin{aligned} \det A &= \det \hat{A} = \det \begin{bmatrix} I & 0 \\ -DB^{-1} & I \end{bmatrix} \det \hat{A} = \det \begin{bmatrix} I & 0 \\ -DB^{-1} & I \end{bmatrix} \begin{bmatrix} B & E \\ D & G \end{bmatrix} \\ &= \det \begin{bmatrix} B & E \\ 0 & G - DB^{-1}E \end{bmatrix} = \det B \det(G - DB^{-1}E), \end{aligned}$$

we see that

$$\det A = \det B \det(A/B).$$

This result is known as *Schur's formula*.

In case A is Hermitian, Haynsworth [5] has shown that the inertia of A can be determined from the inertia of any nonsingular principal submatrix of A together with that of its Schur complement. Other applications and properties of the Schur complement will appear in a later paper.

In §2 of this note, we prove that the Schur complement can also be constructed using quotients of minors of A . Details on this method of construction and its relation to partitioned matrices and M -matrices can be found in [1], [2], [3].

In §3, this construction is used to prove a quotient identity for the Schur complement: $(A/B) = ((A/C)/(B/C))$.

2. Elements of the Schur complement. The notation $A(i_1, \dots, i_p; j_1, \dots, j_p)$ denotes the submatrix of A formed using rows i_1, \dots, i_p

Received by the editors October 2, 1968.

¹ The work of the second author was done under Contract DA-91-591-EUC-3686 of the U. S. Army with the Institute of Mathematics, University of Basel. The author wishes to thank Professor A. M. Ostrowski for very helpful discussions.

and columns j_1, \dots, j_p . For principal submatrices we abbreviate this notation to $A(i_1, \dots, i_p)$.

LEMMA. *Let $C = A(1, \dots, k)$ be a nonsingular leading principal submatrix of A . Let $F = (f_{ij})$ be the matrix with elements*

$$f_{ij} = \det A(1, \dots, k, i; 1, \dots, k, j) / \det C \quad (i, j = k + 1, \dots, n).$$

Then $F = (A/C)$, the Schur complement of C in A .

PROOF. Let b_{ij} denote the bordered minor

$$b_{ij} = \det A(1, \dots, k, i; 1, \dots, k, j) = f_{ij} \det C.$$

With A partitioned in the form

$$A = \begin{bmatrix} C & E \\ D & G \end{bmatrix},$$

let $D^{(i)}$ denote the $(i - k)$ th row of D , and let $E_{(j)}$ denote the $(j - k)$ th column of E . Thus

$$D^{(i)} = [a_{i1}, \dots, a_{ik}] \quad (i = k + 1, \dots, n)$$

and

$$E_{(j)} = \begin{bmatrix} a_{1j} \\ \vdots \\ a_{kj} \end{bmatrix} \quad (j = k + 1, \dots, n).$$

Then for $i, j = k + 1, \dots, n$,

$$b_{ij} = \det \begin{bmatrix} C & E_{(j)} \\ D^{(i)} & a_{ij} \end{bmatrix}.$$

By Schur's formula, $b_{ij} = (a_{ij} - D^{(i)}C^{-1}E_{(j)}) (\det C)$. Thus

$$f_{ij} = (a_{ij} - D^{(i)}C^{-1}E_{(j)}).$$

But these are precisely the elements of the matrix $(A/C) = G - DC^{-1}E$, so the lemma is proved.

We remark that the lemma allows us to restate a result contained in [1, Lemma 1]: *The Schur complement of an M-matrix is an M-matrix.*

3. The quotient property of the Schur complement.

THEOREM. *If B is a nonsingular principal submatrix of A , and C*

is a nonsingular principal submatrix of B , then (B/C) is a nonsingular principal submatrix of (A/C) , and $(A/B) = ((A/C)/(B/C))$.

PROOF. We assume without loss of generality that $C = A(1, \dots, k)$ and $B = A(1, \dots, p)$, with $k < p$. Let $V = (A/C)$, of order $n - k$, and let $W = (B/C)$, of order $p - k$. (We label the rows and columns of V from $k + 1$ to n . Similarly, the indices for W are $i, j = k + 1, \dots, p$, while for (V/W) we use $i, j = p + 1, \dots, n$.) It follows from the lemma that W is a principal submatrix of V . Moreover, W is nonsingular, since by Schur's formula,

$$\det W = (\det B)/(\det C).$$

Now let $\bar{V} = (\det C)V$. For $i, j = p + 1, \dots, n$ we have

$$\begin{aligned} (V/W)_{i,j} &= \det V(k + 1, \dots, p, i; k + 1, \dots, p, j)/\det W \\ &= \det C \det V(k + 1, \dots, p, i; k + 1, \dots, p, j)/\det B \\ &= \det \bar{V}(k + 1, \dots, p, i; k + 1, \dots, p, j)/\det B(\det C)^{p-k}. \end{aligned}$$

Since the elements of the matrix \bar{V} are bordered minors from A , Sylvester's determinant identity [4] enables us to express the determinant of any square submatrix of \bar{V} in terms of the corresponding submatrix of A . In particular,

$$\begin{aligned} \det \bar{V}(k + 1, \dots, p, i; k + 1, \dots, p, j) \\ = (\det C)^{p-k} \det A(1, \dots, p, i; 1, \dots, p, j). \end{aligned}$$

Thus

$$(V/W)_{i,j} = \det A(1, \dots, p, i; 1, \dots, p, j)/\det B,$$

which, by the lemma, equals $(A/B)_{i,j}$.

REFERENCES

1. Douglas E. Crabtree, *Applications of M-matrices to nonnegative matrices*, Duke Math. J. **33** (1966), 197–208.
2. ———, *Characteristic roots of M-matrices*, Proc. Amer. Math. Soc. **17** (1966), 1435–1439.
3. ———, *A matrix identity*, Amer. Math. Monthly **75** (1968), 648–649.
4. F. R. Gantmacher, *The theory of matrices*, Vol. 1, Chelsea, New York, 1959, p. 33.
5. Emilie V. Haynsworth, *Determination of the inertia of a partitioned Hermitian matrix*, Linear Algebra and Appl. **1** (1968), 73–81.

AMHERST COLLEGE AND
AUBURN UNIVERSITY