

## FUNCTIONS POSITIVE DEFINITE IN $C[0, 1]$ <sup>1</sup>

SHELDON J. EINHORN

A series of papers by I. J. Schoenberg discusses positive definite functions and isometries between metric spaces. In particular, we have the following definition from [3]. Let  $S$  be a metric space with the distance between points  $P$  and  $Q$  denoted by  $PQ$ . A real function  $\phi$ , defined on the range of values of the distances, is said to be positive definite in  $S$  if

$$(1) \quad \sum_{i,j=1}^n \phi(P_i P_j) \rho_i \rho_j \geq 0$$

for arbitrary real  $\rho_i$ ,  $n=2, 3, \dots$ , and any  $n$  points  $P_i$  of  $S$ . If, in particular,  $S$  is the space  $C[0, 1]$ , we have the following

**THEOREM.** *Let  $f(x)$  be positive definite in  $C[0, 1]$ . Then*

$$(2) \quad f(x) = k \geq 0 \quad \text{if } x > 0, \quad \text{and } f(0) \geq k.$$

*Conversely, every  $f(x)$  defined by (2) is positive definite in  $C[0, 1]$ .*

The proof depends on a

**LEMMA.** *Let  $0 < x \leq y \leq 2x$ . If the set of all pairs  $(i, j)$  of positive integers such that  $1 \leq i < j \leq n$  is partitioned into two sets  $A$  and  $B$ , then it is possible to find functions  $P_j$  in  $C[0, 1]$  such that*

$$\|P_i - P_j\| = x, \quad (i, j) \in A, \quad \|P_i - P_j\| = y, \quad (i, j) \in B.$$

**PROOF.** The lemma follows directly from a theorem of Banach and Mazur [1, p. 187], which states that every separable metric space may be isometrically imbedded in  $C[0, 1]$ .

**PROOF OF THEOREM.** Let  $\phi$  be positive definite in  $C[0, 1]$ , with  $\phi(x) = a$ ,  $\phi(y) = b$ , and choose  $P$ 's as in the lemma. From (1) we get

$$(3) \quad \{\phi(0) - b\} \sum \rho_j^2 + b \left\{ \sum \rho_j \right\}^2 + 2(a - b) \sum_A \rho_i \rho_j \geq 0,$$

and similarly with interchanges of  $a$  with  $b$  and  $A$  with  $B$ .

Suitable choices of  $\rho_j$ ,  $A$ , and  $n$  in (3) permit the following deductions:

---

Received by the editors January 15, 1969.

<sup>1</sup> This result was previously reported in [2]. The present version responds to a request for publication of a brief proof. A more leisurely discussion with more references is available from the author.

- (i) Let  $\sum \rho_j = 0$  and  $A$  be empty. Then  $\phi(0) \geq b$ .
- (ii) Let  $n = 2m$  with  $m$  of the  $\rho_j$ 's =  $-1$ , the rest =  $+1$ . Let  $A = \{(i, j) \mid \rho_i \rho_j = -1\}$  and let  $m$  become large. Then  $a - b \leq 0$ . Similarly,  $b - a \leq 0$ , so  $a = b$ .
- (iii) Let all  $\rho_j = 1$  and let  $n$  become large. Then  $a \geq 0$ .

Steps (ii) and (iii) show that  $\phi$  is constant on any interval  $[x, 2x]$  for positive  $x$ , whence it follows that it is constant for all positive  $x$ . The theorem is thus proved except for the converse, which is trivial.

#### REFERENCES

1. S. Banach, *Théorie des opérations linéaires*, Chelsea, New York, 1955.
2. S. J. Einhorn, *Functions positive definite in the space C*, Amer. Math. Monthly (4) **75** (1968), 393.
3. I. J. Schoenberg, *Metric spaces and positive definite functions*, Trans. Amer. Math. Soc. **44**(1938), 522-536.

KEYSTONE COMPUTER ASSOCIATES, INC., FORT WASHINGTON, PENNSYLVANIA