

## ON THE SUBSTITUTION PROBLEM FOR FREE GROUPS

PAUL E. SCHUPP

By a substitution problem for a group  $G$  we mean the following. Let  $F = \langle \alpha_1, \dots, \alpha_n \rangle$  be a free group. Let  $G$  be a group with presentation  $\langle a_1, \dots, a_m; R_1, \dots, R_l \rangle$ . Suppose we are given a reduced word  $W(\alpha_1, \dots, \alpha_n)$ . We wish to decide, for  $x \in G$  (where  $x$  is expressed in the generators  $a_1, \dots, a_m$  of  $G$ ), whether or not  $x$  is of the form  $W$ , that is, to decide if there exist  $u_1, \dots, u_n$  in  $G$  such that  $x = W(u_1, \dots, u_n)$ .

There exist groups in which it is even impossible to decide if an element is a commutator [1]. The question of solving substitution problems for free groups seems to arise in several contexts and we henceforth restrict ourselves to the case where  $G$  is free. Even here, the only results we know are that one can decide if an element of a free group is a  $k$ th power and the elegant method of Wicks [4] for deciding if an element of a free group is a commutator. This note presents a proof that the substitution problem is solvable for  $G$  a free group and  $W(\alpha, \beta)$  any element of the free group on  $\alpha$  and  $\beta$ .

**LEMMA.** *Let  $F = \langle \alpha_1, \dots, \alpha_n \rangle$  and  $G = \langle a_1, \dots, a_m \rangle$  be free groups. Let  $W \in F$ , and let  $x \in G$ . Suppose that  $W(u_1, \dots, u_n) = x$  for some  $u_1, \dots, u_n$  in  $G$ . Then there exist  $W^* \in F$  and  $v_1, \dots, v_n$  in  $G$  such that:*

- (i) *The set  $\{v_1, \dots, v_n\}$  is Nielsen reduced.*
- (ii)  *$W^*(v_1, \dots, v_n) = x$  and  $W^*$  is the image of  $W$  under an automorphism of  $F$ .*

**PROOF.** We regard Nielsen transformations as acting on  $n$ -tuples (cf. [2]). Now there is a sequence  $T_1, \dots, T_k$  of Nielsen transformations and an  $n$ -tuple  $(v_1, \dots, v_n)$  of Nielsen reduced elements such that  $(u_1, \dots, u_n) \rightarrow (v_1, \dots, v_n)$  by successively applying the transformations  $T_i$ . We use the same letters to denote both the corresponding Nielsen transformations applied to  $n$ -tuples of generators of  $F$  and the automorphisms of  $F$  which the transformations induce. Thus if  $T_1$  is the Nielsen transformation sending  $(a_1 a_2^{-1}, a_2) \rightarrow (a_1, a_2)$ , then  $(\alpha_1, \alpha_2) \rightarrow (\alpha_1 \alpha_2^{-1}, \alpha_2)$  under  $T_1^{-1}$  and the word  $\alpha_1 \alpha_2^2 \alpha_1 \rightarrow \alpha_1 \alpha_2 \alpha_1 \alpha_2^{-1}$  under the automorphism  $T_1^{-1}$  induced by the Nielsen transformation. Suppose that  $W \rightarrow W^*$  by the successive application of the auto-

---

Received by the editors December 9, 1968.

morphisms  $T_1^{-1}, \dots, T_k^{-1}$  of  $F$ . Then  $W^*(v_1, \dots, v_n) = x$  and  $W^*$  is the image of  $W$  under an automorphism of  $F$ .

For  $y \in G$ , let  $|y|$  denote the length of  $y$  with respect to the generators  $a_1, \dots, a_m$ . Similarly, for  $Z \in F$ , let  $|Z|$  denote the length of  $Z$  with respect to  $\alpha_1, \dots, \alpha_n$ . We will show that one can assume that  $|v_i| \leq |x|$ ,  $i = 1, \dots, n$ , and, if for those  $\alpha_i$  which actually occur in  $W^*$ ,  $v_i \neq 1$ , then  $|W^*| \leq |x|$ . If  $\alpha_j$  does not occur in  $W^*$ , then  $v_j$  can be replaced by 1 without altering the fact that the  $v_i$  are Nielsen reduced and that  $W^*(v_1, \dots, v_n) = x$ .

Let  $t_q$  denote one of the  $(v_i)^{\pm 1}$ . Two cancellation properties of a Nielsen reduced set of nontrivial elements are that if  $t_1 \dots t_p$  is a product with no  $t_q = t_{q+1}^{-1}$  then

- (i)  $|t_1 \dots t_p| \geq p$ , and
- (ii)  $|t_1 \dots t_p| \geq \max_{1 \leq q \leq p} |t_q|$ .

Since  $W^*(v_1, \dots, v_n) = x$  and we are assuming that no  $v_i$  which appears is 1, we conclude from (i) that  $|W^*| \leq |x|$ . By (ii), if  $\alpha_j$  occurs in  $W^*$ , then  $|v_j| \leq |x|$ .

Now by Whitehead's Theorem on the automorphisms of a free group [3], we can effectively list all words  $W' \in F$  such that  $|W'| \leq |x|$  and  $W'$  is the image of  $W$  under an automorphism of  $F$ . We can then enumerate all  $n$ -tuples  $(u'_1, \dots, u'_n)$  such that maximum  $|u'_i|$  does not exceed  $|x|$  and substitute all such  $n$ -tuples into the  $W'$  obtained above. Since  $W'$  is obtained from  $W$  by an automorphism,  $W'(u'_1, \dots, u'_n) = x$  if and only if there exist  $u_1, \dots, u_n$  such that  $W(u_1, \dots, u_n) = x$ .

So far, all that we have done is independent of the number of  $\alpha_i$ . The difficulty is that we have, in general, no algorithm to decide if a word on the  $\alpha$ 's can be obtained from  $W$  by an automorphism and then replacing some letters by 1. However, the situation is manageable if  $n = 2$ .

**THEOREM.** *Let  $F = \langle \alpha, \beta \rangle$ . Let  $W$  be a reduced word of  $F$ . Let  $G$  be a free group. Then there is an algorithm to decide, for  $x \in G$ , if  $x$  is of the form  $W$ .*

**PROOF.** Let  $\sigma_\alpha(Y)$  denote the exponent sum of  $\alpha$  in the word  $Y$ . Let  $a = \sigma_\alpha(W)$ , let  $b = \sigma_\beta(W)$ , and let  $d$  be the greatest common divisor of  $a$  and  $b$ . Since there exist integers  $s, t$ , such that  $d = as + bt$ , we have  $v^d = W(v^s, v^t)$ . Thus if  $x \in G$  is a  $d$ th power, then  $x$  is of the form  $W$ . Suppose  $x$  is not a  $d$ th power and  $W(u_1, u_2) = x$ . Then in the process of obtaining a Nielsen reduced pair  $(v_1, v_2)$ , neither  $v_1$  nor  $v_2$  is 1. For if we had, say  $v_1 = 1$ , then  $x = v_2^r$  where  $r$  is a multiple of  $d$ . Hence  $x$  is either a  $d$ th power or we can decide if  $x$  is of the form  $W$  by examining a finite number of substitutions.

## BIBLIOGRAPHY

1. G. Baumslag, W. W. Boone and B. H. Neumann, *Some unsolvable problems about elements and subgroups of groups*, Math. Scand. **7** (1959), 191–201.
2. W. Magnus, A. Karrass and D. Solitar, *Combinatorial group theory*, Interscience, New York, 1966.
3. J. H. C. Whitehead, *On equivalent sets of elements in a free group*, Proc. London Math Soc. **41** (1936), 48–56.
4. M. J. Wicks, *Commutators in free products*, London Math. Soc. **37** (1962), 433–444.

UNIVERSITY OF ILLINOIS, URBANA