

# AN EXTENSION OF HARTOGS' THEOREM FOR DOMAINS WHOSE BOUNDARY IS NOT SMOOTH

DAVID G. SCHAEFFER<sup>1</sup>

**ABSTRACT.** In this note we obtain an extension of Hartogs' theorem on analytic continuation inside a bounded domain in  $\mathbf{C}^n$  which requires no assumption on the smoothness of the boundary. The standard proof of Hartogs' theorem applies with the minor change of using a Whitney function as the boundary data.

Let  $\Omega$  be a bounded domain in  $\mathbf{C}^n$ ,  $n > 1$ , whose complement is connected. According to Hartogs' theorem, if  $\partial\Omega$  is a surface of class  $C^2$  and if  $u \in C^2(\partial\Omega)$  satisfies the tangential Cauchy-Riemann equations on  $\partial\Omega$ , then  $u$  is the restriction to  $\partial\Omega$  of a function  $U \in C^1(\bar{\Omega})$  analytic inside  $\Omega$ . In this note we observe that the smoothness condition  $\partial\Omega \in C^2$  may be discarded in favor of the very mild hypothesis that  $\Omega$  equals the interior of its closure, providing the boundary data are taken to be a Whitney function of class  $C^2$  which satisfies all the Cauchy-Riemann equations on  $\partial\Omega$ . We remark that an ordinary function  $u$  which satisfies the tangential Cauchy-Riemann equations on a smooth surface  $S$  may be assigned a normal derivative to form a Whitney function on  $S$  which satisfies all the Cauchy-Riemann equations.

We follow closely the proof of Hartogs' theorem in §2.3 of Hörmander [1]. If  $K$  is a compact set in  $\mathbf{C}^n$ , let  $\mathcal{E}^m(K)$  be the space of (complex-valued) Whitney functions on  $K$  of class  $C^m$ . (Chapter I of Malgrange [2] contains a concise introduction to Whitney functions.) We use the notation  $\nabla = (\partial/\partial z_1, \dots, \partial/\partial \bar{z}_1, \dots)$  for the gradient operator.

**THEOREM.** *Let  $\Omega$  be a bounded domain in  $\mathbf{C}^n$ ,  $n > 1$ , such that  $\Omega = \text{Int}(\bar{\Omega})$  and  $\Omega' = \mathbf{C}^n \sim \bar{\Omega}$  is connected. If  $u \in \mathcal{E}^2(\partial\Omega)$  and if  $\bar{\partial}u = \nabla(\partial u) = 0$ , then  $u$  may be extended to a function  $U \in \mathcal{E}^1(\bar{\Omega})$  analytic in  $\Omega$ .*

**PROOF.** By the Whitney extension theorem, there is a function  $v \in C^2(\mathbf{C}^n)$  which is an extension of  $u$ . Define a differential form of type  $(0, 1)$ ,

---

Received by the editors July 18, 1969.

*AMS Subject Classifications.* Primary 3222.

*Key Words and Phrases.* Several complex variables, pseudo-convexity, Hartogs' Theorem, Whitney functions.

<sup>1</sup> This research was partially supported by the National Science Foundation under NSF Grant GP-6761.

$$f = \bar{\partial}v \text{ in } \Omega, \quad f = 0 \text{ on } \mathbf{C}^n \sim \Omega.$$

Then  $f \in C^1(\mathbf{C}^n)$ ,  $\bar{\partial}f = 0$ , and  $f$  has compact support. By Theorem 2.3.1 of Hörmander, there is a function  $w \in C_c^1(\mathbf{C}^n)$  such that  $\bar{\partial}w = f$ . We note that  $\partial\Omega' = \partial\Omega$ , because  $\Omega = \text{Int}(\bar{\Omega})$ ; it follows by unique continuation that  $w|_{\partial\Omega} = 0$ . Of course the function  $U = v - w$  provides the desired analytic continuation of  $u$ .

It is a pleasure to acknowledge a stimulating conversation with Stephen Greenfield which led me to this problem, and I would like to thank Hugo Rossi for reading the paper before publication.

#### REFERENCES

1. L. Hörmander, *An introduction to complex analysis in several variables*, Van Nostrand, Princeton, N. J., 1966. MR 34 #2933.
2. B. Malgrange, *Ideals of differentiable functions*, Tata Institute of Fundamental Research, Bombay and Oxford Univ. Press, London, 1967. MR 35 #3446.

BRANDEIS UNIVERSITY