

SOME ASYMPTOTIC THEOREMS FOR ABSTRACT DIFFERENTIAL EQUATIONS

S. ZAIDMAN¹

ABSTRACT. We consider in this paper results on differential equations with time independent operators; uniqueness of solutions which are bounded in the Stepanoff norm as well as weak almost-periodic solutions are some of the topics here considered.

Introduction. In this paper, which is closely related with some of our previous publications, a number of results concerning differential equations in Hilbert and Banach spaces are derived. They concern asymptotic behaviour, boundedness and almost-periodicity.

1. Our first result, a very simple one, is “essentially” the Theorem 1 in [1]. Here it is given in its natural, operator case framework.

THEOREM 1. *Let H be a Hilbert space, A a closed linear operator in H with dense domain $D(A)$; A^* be its adjoint operator, and suppose that for a real β the relations*

$$(1.1) \quad \begin{aligned} \operatorname{Re}(Ax, x) &\leq \beta(x, x), & \forall x \in D(A), \\ \operatorname{Re}(A^*y, y) &\leq \beta(y, y), & \forall y \in D(A^*) \end{aligned}$$

are verified. Let $u(t)$, $t \geq 0 \rightarrow D(A)$, be a strong solution of equation

$$(1.2) \quad u'(t) = Au(t).$$

Then $\|u(t)\| \leq e^{\beta t} \|u(0)\|$ holds, $\forall t \geq 0$.

PROOF. Let us put $v(t) = e^{-\beta t} u(t)$. Then $v'(t) = (A - \beta I)v(t)$. It is easy to see that $A - \beta I$ is the infinitesimal generator of a strongly continuous one-parameter semigroup $T_\beta(t)$ such that $\|T_\beta(t)\| \leq 1$. We see also the representation $v(t) = T_\beta(t)v(0)$; consequently $\|v(t)\| \leq \|v(0)\| = \|u(0)\|$ and $\|u(t)\| \leq e^{\beta t} \|u(0)\|$.

Now, a very simple result about asymptotic behaviour (cf. Theorem 2 in [1]) is the

Received by the editors September 22, 1969.

AMS Subject Classifications. Primary 3495.

Key Words and Phrases. Hilbert space, linear closed operator, adjoint operator, strong solution, infinitesimal operator, one-parameter semigroup, Banach space, almost-periodic functions, compact operators, dual Banach space, space $L^p(\mathbb{R}^n)$, truncation operator, translation operator.

¹Supported by National Research Council of Canada.

THEOREM 2. *Let us have (1.1) with $\beta < 0$. Then if $f_0 \in H$ is given and $u(t), t \geq 0 \rightarrow D(A)$ is solution of*

$$(1.3) \quad u'(t) = Au(t) + f_0.$$

There exists $w_0 \in H$ such that $\lim_{t \rightarrow \infty} u(t) = w_0$.

PROOF. It follows easily that A^{-1} exists and belongs to $\mathcal{L}(H, H)$. We consider $w_0 = -A^{-1}f_0$, and put $v(t) = u(t) - w_0$. We have $v'(t) = Av(t)$; apply Theorem 1 and we get, as $\beta < 0$, $\lim_{t \rightarrow \infty} u(t) = w_0$.

Our next result is a simple generalization of Lemma 1 in our paper [2].

THEOREM 3. *Let \mathfrak{X} be a Banach space, and $T_t \in \mathcal{L}(\mathfrak{X}, \mathfrak{X}), t \geq 0$, be a one parameter strongly continuous semigroup, such that $\|T_t\| \leq Me^{\beta t}, \beta < 0, t \geq 0$. Let A be its infinitesimal generator and $u(t), -\infty < t < +\infty \rightarrow D(A)$, be a strong solution of $u'(t) = Au(t)$. Then if $\sup_{t \in \mathbb{R}} \int_t^{t+1} \|u(\sigma)\|^2 d\sigma < \infty$ it follows $u(t) = \theta$ for every real t .*

REMARK. Similar results are given in our paper [3].

PROOF. We see firstly the representation: $u(t) = T_{t-t_0}u(t_0), \forall t \geq t_0$.

Then we remark existence of a sequence $(t_n)_1^\infty$ such that $\lim_{n \rightarrow \infty} t_n = -\infty$ and such that $\sup_{n \in \mathbb{N}} \|u(t_n)\| = L < \infty$. Next for arbitrary real t we take n large enough in order to have $t_n < t$ and consequently $u(t) = T_{t-t_n}u(t_n)$. So we derive $\|u(t)\| \leq Me^{\beta(t-t_n)} \cdot L$; for $n \rightarrow \infty$ we get $u(t) = 0$.

2. In this section we give a complement to our result on almost-periodicity of certain relatively-compact valued vector functions (see [4]) by taking into account weakly almost-periodic solutions (of abstract differential equations). Remember that if \mathfrak{X} is a Banach space and \mathfrak{X}^* its strong dual, a continuous function $f(t), -\infty < t < +\infty \rightarrow \mathfrak{X}$ is weakly almost-periodic when $\langle x^*, f(t) \rangle$ is Bohr-almost-periodic for every $x^* \in \mathfrak{X}^*$. Our result is the following

THEOREM 4. *In the Banach space \mathfrak{X} , consider a strongly continuous one-parameter semigroup $T_t \in \mathcal{L}(\mathfrak{X}, \mathfrak{X})$ such that $\lim_{t \rightarrow \infty} T_t x = \theta, \forall x \in \mathfrak{X}$. Let also $Q \in \mathcal{L}(\mathfrak{X}, \mathfrak{X})$ be a compact operator commuting with $T_t, \forall t \geq 0$. Its inverse Q^{-1} exists on a dense set in \mathfrak{X} , and the adjoint $(Q^{-1})^*$ is defined on a dense set in \mathfrak{X}^* . Let A be the infinitesimal generator of T_t ; $f(t)$ a continuous weakly almost-periodic function $-\infty < t < +\infty \rightarrow \mathfrak{X}$; $u(t)$ a strong solution, on the whole real axis of equation $u'(t) = Au(t) + f(t)$, such that $\sup_{t \in \mathbb{R}} \|u(t)\| < \infty$. Then $u(t)$ is weakly almost-periodic.*

PROOF. We remark first, as a standard result, the representation formula

$$u(t) = T_{t-t_0}u(t_0) + \int_{t_0}^t T_{t-s}f(s)ds, \quad t \geq t_0.$$

Next we see that: if $g(t)$, $-\infty < t < +\infty \rightarrow \mathfrak{X}$, is a bounded function such that $\langle x^*, g(t) \rangle$ is almost-periodic for a dense set of elements in the dual space \mathfrak{X}^* , then $g(t)$ is weakly almost-periodic. The result is a corollary of the fact that uniform convergent on R^1 sequences of almost-periodic functions have almost periodic limit. A simple remark now is that $w(t) = Qu(t)$ has representation

$$w(t) = T_{t-t_0}w(t_0) + \int_{t_0}^t T_{t-s}(Qf)(s)ds, \quad t \geq t_0$$

and that range of $w(t)$ is relatively compact in \mathfrak{X} as $t \in R^1$. Then we have

LEMMA. *If $h(t)$ is continuous weakly almost-periodic, $t \in R^1 \rightarrow \mathfrak{X}$, and if Q is a compact operator $\in \mathcal{L}(\mathfrak{X}, \mathfrak{X})$ then Qh is strongly almost-periodic.*

In fact $h(t)$ is bounded, hence Qh has relatively compact range. Moreover $Qh(t)$ is weakly almost-periodic too; by well-known facts $Qh(t)$ is strongly almost-periodic.

By this Lemma, Qf is almost-periodic. We apply our Theorem 1 in [4] and obtain that $w(t) = Qu(t)$ is almost-periodic. Then $u(t) = Q^{-1}w(t) = Q^{-1}Qu(t)$. We take now $x^* \in D((Q^{-1})^*)$ (which is dense in \mathfrak{X}^*).

We have $\langle x^*, Q^{-1}Qu(t) \rangle = \langle (Q^{-1})^*x^*, w(t) \rangle$ which is almost-periodic Bohr. From the above made remarks, $u(t)$ is weakly almost-periodic.

3. Here we remember a certain natural generalization of almost-periodic functions.

DEFINITION 3.1. Let $h(t)$, $0 \leq t < \infty$, be a continuous function with values in the Banach space \mathfrak{X} . We say that $h(t)$ is in the class $\mathcal{B}_{\mathfrak{X}}^{\dagger}$ when the set of translates $(h(t+\eta))_{\eta \geq 0}$ is a relatively compact set in the space $C[0, \infty; \mathfrak{X}]$.

DEFINITION 3.2. Let $h(t)$ be a continuous function, $0 \leq t < \infty \rightarrow \mathfrak{X}$. We say that $h(t)$ is in the class $\mathcal{F}_{\mathfrak{X}}^{\dagger}$ when $\forall \epsilon > 0, \exists L_{\epsilon} > 0, N_{\epsilon} > 0$, such that in every interval $[a, a+L] \subset [0, \infty)$, $\exists \zeta_{\epsilon}$ with property

$$\sup_{t > N_{\epsilon}} \|h(t + \zeta_{\epsilon}) - h(t)\|_{\mathfrak{X}} < \epsilon.$$

In the Appendix of our paper [1] a proof of the inclusion $\mathcal{B}_{\mathfrak{X}}^{\dagger} \subset \mathcal{F}_{\mathfrak{X}}^{\dagger}$ is indicated.

DEFINITION 3.3. A continuous function, $0 \leq t < \infty \rightarrow \mathfrak{X}$, $h(t)$ is called weakly- $\mathcal{B}_{\mathfrak{X}}^{\dagger}$ (resp. weakly- $\mathcal{F}_{\mathfrak{X}}^{\dagger}$) if, for each $x^* \in \mathfrak{X}^*$, $\langle x^*, h(t) \rangle$ is in

class \mathfrak{B}^+ (resp \mathfrak{F}^+) corresponding to \mathfrak{X} = scalar field. It is easy to see that if $g(t) \in \mathfrak{F}_\mathfrak{X}^+$, then, $\forall x^* \in \mathfrak{X}^*$, $\langle x^*, g(t) \rangle \in \mathfrak{F}^+$. Also, we have the standard proof of the fact that uniform limits on $0 \leq t < \infty$ of sequences $(h_n(t))_1^\infty \subset \mathfrak{F}_\mathfrak{X}^+$ belong to the same class. We do now a simple observation, connected with Theorem 5 in [1]. We have precisely the

THEOREM 5. *Let \mathfrak{X} be a Banach space; $T_t, t \geq 0 \rightarrow \mathcal{L}(\mathfrak{X}, \mathfrak{X})$, be a strongly continuous one-parameter semigroup, such that $\|T_t\| \leq M, t \geq 0$. Let A be its infinitesimal generator and $u(t), t \geq 0 \rightarrow D(A)$, be a strong solution of the equation: $u'(t) = Au(t), t \geq 0$. Suppose that $u(t)$ has relatively compact trajectory; then $u(t) \in \mathfrak{F}_\mathfrak{X}^+$.*

PROOF. We have as usual, representation $u(t) = T_t u(0), t \geq 0$. We prove that $u(t) \in \mathfrak{B}_\mathfrak{X}^+$. Consider the set of vector-functions: $\{u(t+\eta)\}_{\eta \geq 0} = \{T_{t+\eta} u(0)\}_{\eta \geq 0}$. By relative compactness we may find a sequence $(\eta_n)_1^\infty \subset [0, \infty)$ such that $(T_{\eta_n} u(0))_{n=1}^\infty$ is a Cauchy sequence in \mathfrak{X} . Then $\{u(t+\eta_n)\}_{n=1}^\infty$ is a Cauchy sequence in $C[0, \infty; \mathfrak{X}]$. This follows from the obvious estimate:

$$\begin{aligned} \|T_{t+\eta_n} u(0) - T_{t+\eta_m} u(0)\| &\leq \|T_t\| \|T_{\eta_n} u(0) - T_{\eta_m} u(0)\| \\ &\leq M \|u(\eta_n) - u(\eta_m)\|. \end{aligned}$$

We complement this result by another one, on weak- $\mathfrak{F}_\mathfrak{X}^+$ solutions.

THEOREM 6. *Let \mathfrak{X} be a Banach space; $T_t, t \geq 0 \rightarrow \mathcal{L}(\mathfrak{X}, \mathfrak{X})$, be a strongly continuous one-parameter semigroup such that $\|T_t\| \leq M, t \geq 0$. Let A be its infinitesimal generator; suppose that for a complex λ_0 , operator $(\lambda_0 - A)^{-1}$ is a linear compact operator in \mathfrak{X} ; suppose also the adjoint operator A^* be densely defined in \mathfrak{X}^* . Consider then $u(t), t \geq 0 \rightarrow D(A)$ a strong solution of $u'(t) = Au(t), t \geq 0$, such that $\|u(t)\| \leq M, t \geq 0$. Then $u(t)$ is weakly- $\mathfrak{F}_\mathfrak{X}^+$.*

PROOF. We have again: $u(t) = T_t u(0), t \geq 0$. Denote by $v(t)$ the vector-function $(\lambda_0 - A)^{-1} u(t)$. Because T_t commutes with $(\lambda_0 - A)^{-1}$ we obtain $v(t) = T_t v(0)$, and moreover $v(t)$ has relatively compact trajectory. We apply the previous theorem and get $v(t) \in \mathfrak{F}_\mathfrak{X}^+$. Hence $u(t) = (\lambda_0 - A)v(t)$. Take then $x^* \in D(A^*)$. We have $\langle x^*, u(t) \rangle = \langle x^*, (\lambda_0 - A)v(t) \rangle = \langle (\lambda_0 - A)^* x^*, v(t) \rangle = \langle y^*, v(t) \rangle$. Applying the previous remarks our result follows.

We end this paper giving, in a concrete case an effective criterium in order that for a given semigroup T_t , the trajectory $\{T_t x\}_{t \geq 0}$ be relatively compact (see Theorem 5). Consider the space $L^p(R^p), 1 \leq p < \infty$. Remember a necessary and sufficient condition for a set $\mathfrak{A} \subset L^p(R^n)$ to be relatively compact:

$$(i) \int_{R^n} |u(x)|^p dx \leq M, \forall u \in \mathfrak{A},$$

(ii) $\lim_{\rho \rightarrow \infty} \int_{\mathbb{R}^n} |u_\rho(x)|^p dx = 0$ uniformly on $u \in \mathfrak{A}$; here $u_\rho(x) = u(x)$, $|x| > \rho$ and $u_\rho(x) = 0$ for $|x| \leq \rho$.

Call $t_\rho; \phi \rightarrow \phi_\rho, L^p \rightarrow L^p$, the truncation operator,

(iii) $\lim_{|h| \rightarrow 0} \int_{\mathbb{R}^n} |(\zeta_h u - u)(x)|^p dx = 0$, uniformly on $u \in \mathfrak{A}$. Here $(\zeta_h u)(x) = u(x+h)$ is the translation operator. Then we have

THEOREM 7. *Let $T_t, t \geq 0 \rightarrow \mathcal{L}(L^p(\mathbb{R}^n), L^p(\mathbb{R}^n)), 1 \leq p < \infty$, be a strongly continuous semigroup such that $\|T_t\| \leq M, t \geq 0$. Suppose that T_t commutes with the truncation operator t_ρ , for each $\rho > 0$ and with the translation operator ζ_h for each $h \in \mathbb{R}^n$. Then the set $\mathfrak{A} = \{T_t \phi_0\}_{t \geq 0}$ is, for fixed $\phi_0 \in L^p(\mathbb{R}^n)$, a relatively compact set in $L^p(\mathbb{R}^n)$.*

The proof is immediate if we apply the previous criterium.

REFERENCES

1. S. Zaidman, *Théorèmes qualitatifs pour des équations aux dérivées partielles*, Acad. R. P. Romine Stud. Cerc. Mat. 6 (1955), 645-666. (Roumanian) MR 17, 1211.
2. ———, *Uniqueness of bounded solutions for some abstract differential equations*, Ann. Univ. Ferrara Sez. VII (to appear).
3. ———, *Bounded solutions of some abstract differential equations*, Proc. Amer. Math. Soc. 23 (1969), 340-342.
4. ———, *On some almost-periodic functions*, Ann. Univ. Ferrara Sez. VII (to appear).

UNIVERSITY OF MONTREAL, MONTREAL, QUEBEC, CANADA