

TWO CELLS WITH N POINTS OF LOCAL NONCONVEXITY

NICK M. STAVRAKAS, W. R. HARE AND J. W. KENELLY

ABSTRACT. A subset S of the plane is a two cell provided S is homeomorphic to $\{x \mid \|x\| \leq 1\}$.

THEOREM. Let S be a two cell with exactly n points of local nonconvexity. Then S is expressible as a union of $n+1$ compact convex sets with mutually disjoint interiors.

I. **Introduction.** We will prove that a two cell S in R^2 which has exactly n points of local nonconvexity is expressible as a union of $n+1$ compact convex sets with mutually disjoint interiors. It follows immediately that S is $n+1$ polygonally connected, i.e., an L_{n+1} set and thus we have as a corollary a special case of a result of Valentine [1].

Throughout, the symbols \cup , \cap and \sim denote set union, set intersection and set difference respectively. The interior, closure, and boundary of a set $S \subset R^2$ are denoted by $\text{int } S$, $\text{cl } S$ and $\text{bd } S$, respectively. The convex hull of a set $S \subset R^2$ is denoted by $H(S)$. If $x, y \in S$ then $[xy]$ and (xy) denote the closed and open line segments joining x to y , respectively. The Euclidean norm is given by $\| \cdot \|$ and d denotes the Hausdorff metric on compact subsets of the plane as given in Valentine [2]. If S is a set, $|S|$ denotes its cardinality. We define the distance between a point x and a set S as $\inf \{ \|x - y\| \mid y \in S \}$; we denote this distance by $p(x, S)$. By the ϵ ball about a set S we mean $\{x \mid p(x, S) < \epsilon\}$.

DEFINITION 1. A point $x \in S$ is called a point of *local convexity* of S if there exists a neighborhood N of x such that $N \cap S$ is convex. If such a neighborhood does not exist, x is called a point of *local nonconvexity*.

DEFINITION 2. A set S is said to be *starshaped* relative to a point p if for each $x \in S$, $[xp] \subset S$.

DEFINITION 3. A segment $[xy]$ is said to be a *crosscut* of a set S provided $x, y \in \text{bd } S$ and $(xy) \subset \text{int } S$.

DEFINITION 4. A set $S \subset R^2$ is a *two cell* provided S is homeomorphic to $\{x \mid \|x\| \leq 1\}$.

Given a two cell S and a crosscut $[xy]$ of S , $[xy]$ induces a natural decomposition of S into two new two cells D^1xy and D^2xy such that

Received by the editors February 3, 1970.

AMS 1969 subject classifications. Primary 5225, 5014.

Key words and phrases. Convex set, local convexity, two cell.

Copyright © 1971, American Mathematical Society

$D^1xy \cap D^2xy = [xy]$. For a proof of this, see Newman [1]. Specifically, the points x and y divide $\text{bd } S$ into two disjoint relatively open connected subsets of $\text{bd } S$, say C^1xy and C^2xy , and D^1xy is the set whose boundary is given by $[xy] \cup C^1xy$ and D^2xy is the set whose boundary is given by $[xy] \cup C^2xy$. If S is a set, $C(S)$ and $L(S)$ will denote its points of local convexity and nonconvexity, respectively, and \emptyset will denote the empty set. The following two theorems constitute the main results of this paper.

THEOREM 1. *Let $S \subset R^2$ be a two cell such that $|L(S)| = n$, $n \geq 2$. Then there exists a crosscut $[xy]$ of S such that $x, y \in C(S)$ and $|L(D^i xy)| \leq n - 1$ for $i = 1, 2$.*

THEOREM 2. *Let $S \subset R^2$ be a two cell such that $|L(S)| = n$. Then $S = \bigcup_{i=1}^{n+1} C_i$, where C_i , $1 \leq i \leq n+1$, are compact, convex and have mutually disjoint interiors.*

Theorem 1 is of some independent interest, since it makes induction arguments readily accessible. Also, note that if $n = 0$ in Theorem 2, then the latter reduces to a special case of the important theorem of Tietze [1], which we state for later reference.

THEOREM 3. *Let S be a closed connected set in R^d , all of whose points are points of local convexity. Then S is convex.*

We also shall utilize the following result of Valentine [1].

THEOREM 4. *Let S be a closed connected set in R^d such that $L(S)$ is not empty. Then given $x \in S$, there exists $y \in L(S)$ such that $[xy] \subset S$.*

II. Preliminary results and proof of Theorems 1 and 2.

THEOREM 5. *Let $\{A_i\}_{i=1}^{\infty}$ be a sequence of compact sets in R^2 converging to a set A in the Hausdorff metric d such that for each i , $A_i = \bigcup_{j=1}^{n+1} C_j^i$ where each C_j^i is convex and compact for $1 \leq j \leq n+1$ and $\text{int } C_j^i \cap \text{int } C_k^i = \emptyset$, for $k \neq j$ and $1 \leq k, j \leq n+1$. Then $A = \bigcup_{j=1}^{n+1} C_j$ where C_j is compact and convex for $1 \leq j \leq n+1$ and $\text{int } C_j \cap \text{int } C_k = \emptyset$ for $j \neq k$, $1 \leq j, k \leq n+1$.*

PROOF. By the theorem of Blaschke we may assume without loss of generality that for each j the sequence $\{C_j^i\}_{i=1}^{\infty}$ converges to a set C_j . Now C_j is compact and convex, since C_j is the limit of compact convex sets. Since for each i and $j \neq k$ $\text{int } C_j^i \cap \text{int } C_k^i = \emptyset$, we have $\text{int } C_j \cap \text{int } C_k = \emptyset$, and this completes the proof.

THEOREM 6. *Let $S \subset R^2$ be a two cell such that $|L(S)| = 1$. Then $S = C_1 \cup C_2$ where C_1 and C_2 are compact, convex and $\text{int } C_1 \cap \text{int } C_2 = \emptyset$.*

PROOF. Throughout the proof let $L(S) = \{x\}$.

Case 1. Suppose there exist $z, q \in \text{bd } S$, such that $z \neq x, q \neq x, [zx] \cup [xq] \subset \text{bd } S$ and $[zx] \cap [xq] = x$. Then clearly we may choose a crosscut $[xy]$ where $y \in C(S)$ such that $x \in C(D^1xy) \cap C(D^2xy)$. Since $y \in C(S)$, we have $y \in C(D^1xy) \cap C(D^2xy)$. Then D^1xy and D^2xy are convex by Tietze's Theorem and $\text{int } D^1xy \cap \text{int } D^2xy = \emptyset$ by definition. Thus D^1xy and D^2xy are the required sets.

Case 2. Suppose for each $y \in \text{bd } S, y \neq x$, that $[xy] \not\subset \text{bd } S$. For this case, we need a lemma.

LEMMA 1. *Let S be a two cell satisfying Case 2. Then if $t \in \text{bd } S, t \neq x, (xt) \subset \text{int } S$.*

PROOF. Suppose the lemma is false. By Theorem 4, S is starshaped relative to x , so $[xt] \subset S$. By hypothesis, $(xt) \not\subset \text{bd } S$, so let $z \in (xt) \cap \text{int } S$. Since S is starshaped relative to x , for each $m \in (xt) \cap \text{int } S$ we have $(xm) \subset \text{int } S$. Then, since we are denying the lemma, we may not find a sequence $\{z_i\}_{i=1}^\infty$ such that $z_i \in (xt) \cap \text{int } S$ such that $\{z_i\}_{i=1}^\infty$ converges to t , for otherwise $(xt) = \bigcup_{i=1}^\infty (xz_i)$ and $\bigcup_{i=1}^\infty (xz_i) \subset \text{int } S$ by the last sentence, so $(xt) \subset \text{int } S$, a contradiction. Thus let $q \neq t$ be the closest point of $\text{bd } S \cap [zt]$ to x . Then $[tq] \subset \text{bd } S$ and $(qx) \subset \text{int } S$. Let $q_1 \in (tq)$. Since q_1 is a point of local convexity, and since a two cell is the closure of its interior, we may choose $\epsilon > 0$, so that $B(q_1, \epsilon) \cap S$ is convex and $D = \text{int}(B(q_1, \epsilon) \cap S) \neq \emptyset$, where $B(q_1, \epsilon) = \{p \mid \|p - q_1\| < \epsilon\}$. Let $\mathcal{C} = H(D \cup \{x\}) \sim \{x\}$. Then \mathcal{C} is an open set and $\mathcal{C} \subset \text{int } S$ since x is a seeing point. Now let $\{r_i\}_{i=1}^\infty$ be a sequence of boundary points converging to q , such that $r_i \neq q, r_i \notin (tq)$ for each i . Then $\{r_i\}_{i=1}^\infty$ must approach q from the open half space generated by the line containing $[xt]$, opposite from the open half space containing \mathcal{C} . Choose $\delta > 0$, so that $E = B(q, \delta) \cap S$ is convex. Then, clearly we have $\text{int}(E \cap \mathcal{C}) \neq \emptyset$. Since $\{r_i\}_{i=1}^\infty$ converges to q , there exists an integer k , such that if $i \geq k$, we have $r_i \in E$. Let $x_1 \in \text{int}(E \cap \mathcal{C})$ be such that $q \in (r_k x_1) \subset \text{int } S$. This contradicts that q is a boundary point.

Returning to Case 2, let f be the homeomorphism mapping the closed unit disc onto S . Then $\text{bd } S$ is homeomorphic to the unit circle. Let $f(x^*) = x$. Let $y^* \neq x^*$ be on the unit circle. Then x^* and y^* naturally divide the unit circle into two nonempty, relatively open, disjoint, connected subsets, say $B_{x^*y^*}^1$ and $B_{x^*y^*}^2$. Let $\{x_i^1\}_{i=1}^\infty$ and $\{x_i^2\}_{i=1}^\infty$ be sequences of points in $\text{bd } S$ such that the inverse images of x_i^1 and x_i^2 under f are in $B_{x^*y^*}^1$ and $B_{x^*y^*}^2$, respectively, for each i , and both sequences coverge to x . Now define the set S_i for each i to be that subset of S whose boundary is given by $[x_i^1x] \cup [xx_i^2] \cup f(B_i)$ where B_i is $B_{x_i^1x_i^2}^1$ if $x \notin B_{x_i^1x_i^2}^1$ and B_i is $B_{x_i^1x_i^2}^2$ if $x \notin B_{x_i^1x_i^2}^2$. Now since S is

simply connected, $S_i \subset S$, and S_i for each i is well defined by Lemma 1. Then $\{S_i\}_{i=1}^\infty$ converges to S in the Hausdorff metric. Since for each i , $x_i^1, x_i^2 \in C(S)$, we have $x_i^1, x_i^2 \in C(S_i)$. Then, the only possible point of local nonconvexity of S_i is x . For some integer k , we must have for each $i \geq k$, that $x \in L(S_i)$ for otherwise S would be convex. Now by Case 1 each S_i , $i \geq k$, is a union of two compact convex sets with mutually disjoint interiors. Thus, by Theorem 5, S is a union of two compact convex sets with mutually disjoint interiors.

Case 3. Suppose there exists $z \in \text{bd } S$ such that $z \neq x$, $[xz] \subset \text{bd } S$ and there does not exist $z_1 \in \text{bd } S$, $z_1 \neq z$, such that $[xz] \subset [xz_1] \subset \text{bd } S$. Further, suppose for each $y \in \text{bd } S \sim [xz]$ that $[xy] \not\subset \text{bd } S$. Then, following a similar proof as in Lemma 1 we show that if $y \in \text{bd } S \sim [xz]$, then $(xy) \subset \text{int } S$. Finally, as in Case 2, we construct a sequence of compact sets $\{S_i\}_{i=1}^\infty$ such that $|L(S_i)| = 1$ for each i beyond some k and $\{S_i\}_{i=1}^\infty$ converges to S in the Hausdorff metric, and then apply Theorem 5.

PROOF OF THEOREM 1. We begin with a lemma.

LEMMA 2. *Let S be a two cell such that $|L(S)|$ is finite. Let $[xy]$ be a crosscut of S such that $x, y \in C(S)$. Then there exists a convex subset Axy of S such that $(xy) \subset \text{int } Axy$.*

PROOF. Choose $\epsilon > 0$ so that $B(x, \epsilon) \cap S$ is convex, $B(y, \epsilon) \cap S$ is convex, $\text{cl } B(x, \epsilon) \cap L(S) = \emptyset$ and $\text{cl } B(y, \epsilon) \cap L(S) = \emptyset$. For each $z \in (xy)$, choose $\epsilon_z > 0$ so that $\text{cl } B(z, \epsilon_z) \subset \text{int } S$. Since $[xy]$ is compact select a finite subcover $\{B(x, \epsilon/2), B(y, \epsilon/2), B(z_1, \epsilon_{z_1}/2), \dots, B(z_n, \epsilon_{z_n}/2)\}$ of the cover $\{B(x, \epsilon/2), B(y, \epsilon/2), B(z, \epsilon_z/2) \mid z \in (xy)\}$. Choose $\delta > 0$ so that $\delta < \min\{\epsilon/2, \epsilon_{z_1}/2, \dots, \epsilon_{z_n}/2\}$. Let B be the closure of the $\delta/\sqrt{2}$ ball about the set $[xy]$. Then $B \cap S$ is clearly connected, closed and $B \cap S \cap L(S) = \emptyset$. Then $B \cap S$ is convex by Tietze's Theorem and this is the required set Axy .

Now suppose Theorem 1 is false. Let $r, s \in L(S)$ such that $C_r^1 \cap L(S) = \emptyset$ or $C_r^2 \cap L(S) = \emptyset$, which is possible since $|L(S)|$ is finite. Let us suppose that $C_r^1 \cap L(S) = \emptyset$. Let $\mathcal{C} = \{[xy] \mid [xy] \text{ is a crosscut and } x, y \in C(S)\}$. Since $|L(S)|$ is finite, every interior point of S is contained in an element of \mathcal{C} , so $\text{int } S \subset \cup \mathcal{C}$. Since we are assuming that Theorem 1 is false, if $[xy] \in \mathcal{C}$, then $x, y \in C_r^1$ or $x, y \in C_r^2$.

Consider any two crosscuts in \mathcal{C} , say $[x_1y_1]$ and $[x_2y_2]$ such that $x_1, y_1 \in C_r^1$ and $x_2, y_2 \in C_r^2$. These crosscuts can not intersect. To see this suppose $z \in (x_1y_1) \cap (x_2y_2)$. Then the set R whose boundary is given by $[x_1y_1]$ and the portion B of $\text{bd } S$ in C_{zy}^1 between x_1 and y_1 , is convex by Tietze's Theorem. Since $B \cap [x_2y_2] = \emptyset$, this forces x_2 or y_2

to be an interior point of R , and hence an interior point of S , a contradiction.

To continue, the interior of S is connected since it is the homeomorphic image of a connected set. Let $[xy] \in \mathcal{C}$. Let Axy be as in Lemma 2. Then letting $\mathcal{Q}_1 = \{\text{int } Axy \mid [xy] \in \mathcal{C}, x, y \in C_{rs}^1\}$ and $\mathcal{Q}_2 = \{\text{int } Axy \mid [xy] \in \mathcal{C}, x, y \in C_{rs}^2\}$, we have $\text{int } S \subset \cup \mathcal{Q}_1 \cup \cup \mathcal{Q}_2$ since $\cup \mathcal{C} = \cup \mathcal{Q}_1 \cup \cup \mathcal{Q}_2$. Since $\text{int } S$ is connected, we must have $\cup \mathcal{Q}_1 \cap \cup \mathcal{Q}_2 \neq \emptyset$, which says that there exists $[x_1y_1]$ and $[x_2y_2]$ in \mathcal{C} with $x_1, y_1 \in C_{rs}^1, x_2, y_2 \in C_{rs}^2$ and $\text{int } Ax_1y_1 \cap \text{int } Ax_2y_2 \neq \emptyset$. Let $z \in \text{int } Ax_1y_1 \cap \text{int } Ax_2y_2$. Then z lies in a crosscut having endpoints in C_{rs}^1 , and in a crosscut having endpoints in C_{rs}^2 , which says these crosscuts intersect, which is a contradiction by the last paragraph. Thus Theorem 1 holds.

PROOF OF THEOREM 2. We know Theorem 2 holds when $n = 0$, and $n = 1$ by Tietze's Theorem and Theorem 6, respectively. Thus, assume the theorem holds for $0 \leq k \leq n - 1$ and we will show the result holds for n . We begin by considering cases. Let $x \in L(S)$.

Case 1. Suppose there exist $z, q \in \text{bd } S$ such that $z \neq x, q \neq x, [zx] \cup [xq] \subset \text{bd } S$ and $[zx] \cap [xq] = \{x\}$. Then choose a crosscut $[xy]$ where $y \in C(S)$ such that $x \in C(D^1xy) \cap C(D^2xy)$. Since $y \in C(S), y \in C(D^1xy) \cap C(D^2xy)$. Now suppose $d = |L(D^1xy)|$. Then $d \leq n - 1$ and by hypothesis $D^1xy = \cup_{i=1}^{d+1} C_i$, where for each $i, 1 \leq i \leq d + 1, C_i$ is compact and convex and for $1 \leq i, j \leq d + 1, i \neq j, \text{int } C_i \cap \text{int } C_j = \emptyset$. Since $|L(D^1xy)| = d$ and since $x \in C(D^1xy) \cap C(D^2xy)$, we have $|L(D^2xy)| = n - d - 1$. Thus, by hypothesis $D^2xy = \cup_{i=1}^{n-d} B_i$, where for each $i, 0 \leq 1 \leq n - d, B_i$ is compact, convex and for $i \neq j, 1 \leq i, j \leq n - d, \text{int } B_i \cap \text{int } B_j = \emptyset$. Thus S is a union of $(d + 1) + (n - d) = n + 1$ compact convex sets with mutually disjoint interiors.

Case 2. Suppose for each $y \in \text{bd } S, y \neq x$, that $[xy] \not\subset \text{bd } S$. We shall need the following lemma.

LEMMA 3. Let S be a two cell such that $|L(S)| = n \geq 1$ satisfying Case 2. Let $\{x_i\}_{i=1}^\infty$ be a sequence in $\text{bd } S$ with $x_i \rightarrow x, x \in L(S)$ and $x_i \neq x$ for each i . Then there exists an integer k such that for each $i \geq k, (xx_i) \subset \text{int } S$.

PROOF. We proceed by induction. We know by Lemma 1 the lemma holds when $|L(S)| = 1$. So suppose the lemma holds for $1 \leq k \leq n - 1$. By Theorem 1, there exists a crosscut $[rs]$ of S such that $r, s \in C(S)$ and $|L(D_{rs}^i)| \leq n - 1$ for $i = 1, 2$. Suppose $x \in L(D_{rs}^1)$. Then since there exists an $\epsilon > 0$ such that $B(x, \epsilon) \cap S \subset D_{rs}^1$, the result holds. The same argument applies if $x \in D_{rs}^2$.

To complete Case 2, we argue the same way as in Case 2 of Theorem 6. We construct a sequence of sets $\{S_i\}_{i=1}^\infty$ such that $\{S_i\}_{i=1}^\infty$ converges to S in the Hausdorff metric and using Case 1 of this theorem

we show each S_i is representable as $n+1$ compact convex sets with mutually disjoint interiors and then apply Theorem 5. This completes Case 2.

Case 3. Suppose there exists $x \in L(S)$ and $z \in \text{bd } S$, $z \neq x$ such that $[zx] \subset \text{bd } S$ and there does not exist $z_1 \in \text{bd } S$, $z_1 \neq z$ such that $[xz] \subset [xz_1] \subset \text{bd } S$. Further suppose for each $y \in \text{bd } S \sim [xz]$, that $[x, y] \not\subset \text{bd } S$. Then using a similar proof as in Lemma 3, we prove that if $\{x_i\}_{i=1}^\infty$ is a sequence in $\text{bd } S \sim [xz]$ converging to x , then there exists an integer k , such that for each $i \geq k$, $(x_i x) \subset \text{int } S$ and use the same construction as in Case 2 to get the theorem. This completes the proof of Theorem 2.

III. Consequences of Theorem 2. As immediate corollaries of Theorem 2 we have

COROLLARY 1. *Let S be a two cell such that $|L(S)| = n$. Then S is an L_{n+1} set.*

COROLLARY 2. *Let S be a two cell such that $|L(S)| = n$. Then for every line L , $L \cap S$ is a union of at most $n+1$ closed line segments.*

Corollary 2 is a generalization of the familiar fact that the intersection of a line and a compact convex set is a point or a closed line segment.

Examples are easily constructable to show the number $n+1$ is best in Theorem 2.

REFERENCES

M. H. A. NEWMAN

1. *Elements of the topology of plane sets of points*, 2nd ed., Cambridge Univ. Press, Cambridge, 1951, pp. 144–145. MR 13, 483.

H. TIETZE

1. *Über Convexität im kleinen und im grossen und über gewisse den Punkten einer Menge zugeordnete Dimensionzahlen*, Math. Z. 28 (1928), 697–707.

F. A. VALENTINE

1. *Local convexity and L_n sets*, Proc. Amer. Math. Soc. 16 (1965), 1305–1310. MR 32 #2976.

2. *Convex sets*, McGraw-Hill Series in Higher Math., McGraw-Hill, New York, 1964. MR 30 #503.

CLEMSON UNIVERSITY, CLEMSON, SOUTH CAROLINA 29631