

SKELETA OF COMPLEXES WITH LOW MU_* PROJECTIVE DIMENSION¹

DAVID COPELAND JOHNSON

ABSTRACT. Let $MU_*(X)$ be the unitary bordism of a finite complex X . Let X^p be the p -skeleton of X . This note proves that certain properties of $MU_*(X)$ are shared by $MU_*(X^p)$ when the projective dimension of $MU_*(X)$ as a MU_* module is low (0, 1, or 2).

Introduction. Let X be a finite complex and let X^p be any skeleton of X . This note examines some algebraic properties which are inherited by the skeleton X^p from the complex X . Our work is motivated by the elementary example: if $H_*(X; Z)$ is free abelian, then $H_*(X^p; Z)$ is also free abelian.

$MU_*()$ is the complex bordism homology theory associated to the Thom spectrum MU . We denote the projective dimension of $MU_*(X)$ as a $MU_* \equiv MU_*(\text{Point})$ -module by $\text{hom dim}_{MU_*} MU_*(X)$. Connor and Smith prove: $\text{hom dim}_{MU_*} MU_*(X) = 0$ if and only if $H_*(X; Z)$ is free abelian [3, 3.10]. Our elementary example becomes the $n=0$ version of our first theorem.

THEOREM 1. *Let X be a finite complex. If $n=0, 1,$ or $2,$ then $\text{hom dim}_{MU_*} MU_*(X) \leq n$ if and only if $\text{hom dim}_{MU_*} MU_*(X^p) \leq n$ for every skeleton X^p of X .*

The "if" part of the theorem is trivial. The $n=1$ version is a folk theorem; we shall sketch a proof for completeness.

$k_*()$ is the connective k -theory; it is the homology theory derived from the connected unitary spectrum bu ([1], [8]). We prove:

THEOREM 2. *Let X be a finite complex. The following four conditions are equivalent.*

- (i) $k_*(X)$ is free abelian.
- (ii) $k_*(X^p)$ is free abelian for every skeleton X^p of X .
- (iii) $MU_*(X^p)$ is free abelian for every skeleton X^p of X .
- (iv) $MU_*(X)$ is free abelian and $\text{hom dim}_{MU_*} MU_*(X) \leq 1$.

Received by the editors June 17, 1971.

AMS 1970 subject classifications. Primary 55B20.

Key words and phrases. Unitary bordism, connective k -theory, homological dimension, finite complexes, Atiyah-Hirzebruch-Dold spectral sequence.

¹ During the preparation of this paper the author was supported by a University of Kentucky Faculty Summer Research Fellowship.

© American Mathematical Society 1972

As there exist finite complexes X with $MU_*(X)$ free abelian and with $\text{hom dim}_{MU_*} MU_*(X)$ arbitrarily high, we see that $MU_*(X)$ free abelian does not imply $MU_*(X^p)$ is also.

Postnikov fibres of homology theories. Let $M_*()$ be a homology theory and q an integer. We follow Dold [4] in defining the q th Postnikov fibre of $M_*()$ to be the homology theory $M(q)_*()$ with groups $M(q)_{p+q}(X)$ for a finite complex X given by

$$M(q)_{p+q}(X) = \text{Image}\{M_{p+q}(X^{p-1}) \rightarrow M_{p+q}(X^p)\}.$$

Recall the skeletal filtration exact couple for the Atiyah-Hirzebruch-Dold spectral sequence for $M_*(X)$ has $D_{p,q}^1 = M_{p+q}(X^p)$ and $E_{p,q}^1 = M_{p+q}(X^p, X^{p-1})$. So

$$M(q)_{p+q}(X) = \text{Image}\{D_{p-1,q+1}^1 \rightarrow D_{p,q}^1\}$$

which is $D_{p,q}^2$ in the derived exact couple. There is a natural homomorphism $M(q)_{p+q}(X) \rightarrow M(q-1)_{p+q}(X)$ which is $i_{p,q}^2: D_{p,q}^2 \rightarrow D_{p+1,q-1}^2$ in the derived exact couple [7, §XI-5]. The converse of the following folk lemma also holds, but we shall not need it.

LEMMA 3. $M(q)_{p+q}(X) \rightarrow M(q-1)_{p+q}(X)$ is a monomorphism modulo torsion. When it is monic, the Atiyah-Hirzebruch-Dold spectral sequence for $M_*(X)$ collapses.

PROOF. The first statement is seen by tensoring the derived exact couple with \mathcal{Q} , the rationals. For the second statement, note that if all the $i_{p,q}^2$'s are monic in the derived exact couple, then all the $k_{p,q}^2$'s are zero. Since each differential involves a $k_{p,q}^2$, the spectral sequence collapses.

Q.E.D.

We shall work with three homology theories: integral homology, connective k -theory, and complex bordism. All three satisfy the hypothesis of Lemma 4.

LEMMA 4. Let $M_*()$ be a homology theory such that $M_{2n+1} \equiv M_{2n+1}(\text{Point}) = 0$ for each integer n . Then

$$M(2r + 1)_*(X) \cong M(2r)_*(X)$$

for any finite complex X .

PROOF. This follows immediately from definitions and the fact that $M_{2r+1+s}(X^s, X^{s-1}) = 0$. Q.E.D.

Let $f: A_*() \rightarrow B_*()$ be a natural transformation of two homology theories. The inclusion of skeleta, $i: X^{p-1} \rightarrow X^p$, of a finite complex X

induces the commuting diagram:

$$\begin{array}{ccc}
 A_{p+q}(X^{p-1}) & \xrightarrow{A_*(i)} & A_{p+q}(X^p) \\
 \downarrow f_1 & & \downarrow f_2 \\
 B_{p+q}(X^{p-1}) & \xrightarrow{B_*(i)} & B_{p+q}(X^p)
 \end{array}$$

we may define

$$f(q): A(q)_{p+q}(X) = \text{Image } A_*(i) \rightarrow B(q)_{p+q}(X) = \text{Image } B_*(i)$$

by

$$f(q)(A_*(i)(y)) = f_2 A_*(i)(y) = B_*(i) f_1(y) \quad \text{for any } y \in A_{p+q}(X^{p-1}).$$

In particular: if spectra A and B represent the homology theories $A_*()$ and $B_*()$, a degree 0 morphism of spectra $f: A \rightarrow B$ induces natural transformations $f: A_*() \rightarrow B_*()$ and $f(q): A(q)_*() \rightarrow B(q)_*()$. Let

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A$$

be an exact triangle of spectra where f and g are of degree 0 and h is of degree -1 [10, §XI]. This induces a long exact sequence of homology theories:

$$\cdots \rightarrow A_n() \xrightarrow{f} B_n() \xrightarrow{g} C_n() \xrightarrow{h} A_{n-1}() \rightarrow \cdots$$

We start to ask whether $f(q), g(q), \dots$ are in a long exact sequence, but we recall that $h(q)$ has not been (and may not be) defined.

LEMMA 5. *If $A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} A$ is an exact triangle of spectra as above such that the homology theories $A_*()$, $B_*()$, and $C_*()$ satisfy the hypothesis of Lemma 4, then there is a long exact sequence of homology theories:*

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & A(2r)_n() & \xrightarrow{f(2r)} & B(2r)_n() & \xrightarrow{g(2r)} & C(2r)_n() \\
 & & & & & & \xrightarrow{h(2r)} \\
 & & & & & & A(2r)_{n-1}() \longrightarrow \cdots
 \end{array}$$

PROOF. For a finite complex X , let $i: X^{n-2r-1} \rightarrow X^{n-2r}$ and $j: X^{n-2r-2} \rightarrow X^{n-2r-1}$ be inclusions of skeleta. These induce the exact rows in the diagram below.

$$\begin{array}{ccccccc}
 C_{n+1}(X^{n-2r}, X^{n-2r-1}) & & & & & & \\
 = & & & & & & \\
 0 & \longrightarrow & C_n(X^{n-2r-1}) & \xrightarrow{C_n(i)} & C_n(X^{n-2r}) & & \\
 & & \downarrow 1 & & & & \\
 C_n(X^{n-2r-2}) & \xrightarrow{C_n(j)} & C_n(X^{n-2r-1}) & \longrightarrow & C_n(X^{n-2r-1}, X^{n-2r-2}) = 0 & & \\
 \downarrow h_1 & & \downarrow h_2 & & & & \\
 A_{n-1}(X^{n-2r-2}) & \xrightarrow{A_{n-1}(j)} & A_{n-1}(X^{n-2r-1}) & & & &
 \end{array}$$

Define

$$h(2r): C(2r)_n(X) = \text{Image } C_n(i) \rightarrow A(2r)_{n-1}(X) = \text{Image } A_{n-1}(j)$$

by

$$h(2r)(C_n(i)(y)) = h_2(y) = A_{n-1}(j) \circ h_1 \circ C_n(j)^{-1}(y) \quad \text{for any } y \in C_n(X^{n-2r-1}).$$

Verification of exactness is routine. Q.E.D.

LEMMA 6. *Let X be a finite complex. There is a finite complex Y and a map g: Y → Σ^sX (Σ^sX is the s-fold suspension of X) such that:*

- (i) *H_{*}(Y) is free abelian;*
 - (ii-1) *g_{*}: H(2r)_{*}(Y) → H(2r)_{*}(Σ^sX) is epic if hom dim_{MU_{*}} MU_{*}(X) ≤ 1;*
- or
- (ii-2) *g_{*}: k(2r)_{*}(Y) → k(2r)_{*}(Σ^sX) is epic if hom dim_{MU_{*}} MU_{*}(X) ≤ 2.*

PROOF. $H(2r)_*() = H_*(; Z)$ for r negative and $H(2r)_*()$ is the zero homology theory for r nonnegative. From Bott periodicity, there is the isomorphism $k(2r)_n() \cong k_{n-2r}()$ [1]. The lemma follows from the existence of MU-resolutions ([3, 2.4 and 3.11], [6]). Q.E.D.

PROOF OF THEOREM 1. *The n=2 case.* We assume X is a finite complex with $\text{hom dim}_{MU_*} MU_*(X) \leq 2$.

The Thom homomorphism $\zeta: MU_*() \rightarrow k_*()$ is induced by a degree 0 morphism of spectra $\zeta: MU \rightarrow bu$. By [10], we may construct an exact triangle of spectra

$$A \xrightarrow{f} MU \xrightarrow{\zeta} bu \xrightarrow{h} A$$

with $\text{degree}(f) = 0$ and $\text{degree}(h) = -1$. For any finite complex X, we have the long exact sequence

$$\begin{array}{ccccc} \longrightarrow & A_*(X) & \xrightarrow{f} & MU_*(X) & \xrightarrow{\zeta} & k_*(X) & \longrightarrow \\ & & & & & & \\ & & & & & h & \end{array}$$

($\text{degree}(h) = -1$). By [5], $\text{hom dim}_{MU_*} MU_*(X) \leq 2$ if and only if ζ is epic. We conclude: $\text{hom dim}_{MU_*} MU_*(X) \leq 2$ if and only if f is monic. If $H_*(Y; Z)$ is free abelian and if X is Y or Y^p , then f is monic. In particular if X is a one point space, this proves that $A_*()$ satisfies the hypothesis of Lemma 4.

We claim: $\text{hom dim}_{MU_*} MU_*(X^p) \leq 2$ if and only if $f(q): A(q)_{p+q}(X) \rightarrow MU(q)_{p+q}(X)$ is monic for all integers p and q. From the exact sequences for the pair (X^p, X^{p-1}) , we have exact rows in the diagram below.

$$\begin{array}{ccccccc} 0 & \longrightarrow & A(q)_{p+q}(X) & \longrightarrow & A_{p+q}(X^p) & \longrightarrow & A_{p+q}(X^p, X^{p-1}) \\ & & \downarrow f(q) & & \downarrow f_1 & & \downarrow f_2 \\ 0 & \longrightarrow & MU(q)_{p+q}(X) & \longrightarrow & MU_{p+q}(X^p) & \longrightarrow & MU_{p+q}(X^p, X^{p-1}). \end{array}$$

Since $H_*(X^p, X^{p-1}; Z)$ is free abelian, f_2 is monic. By the "5" Lemma, f_1 is monic if and only if $f(q)$ is monic. This proves the claim.

To prove the proposition, we pick $g: Y \rightarrow \Sigma^s X$ as in Lemma 6. Since $H^*(Y; Z)$ is free abelian, $f(2r)': A(2r)_* Y \rightarrow MU(2r)_*(Y)$ is monic and the top row in our diagram below is short exact. The bottom sequence is exact by Lemma 5.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & A(2r)_*(Y) & \xrightarrow{f(2r)'} & MU(2r)_*(Y) & \xrightarrow{\zeta(2r)'} & k(2r)_*(Y) \longrightarrow 0 \\
 & & \downarrow g_1 & & \downarrow g_2 & & \downarrow g_3 \\
 & & A(2r)_*(\Sigma^s X) & \xrightarrow{f(2r)} & MU(2r)_*(\Sigma^s X) & \xrightarrow{\zeta(2r)} & k(2r)_*(\Sigma^s X) \longrightarrow 0 \\
 & & \searrow & \xrightarrow{h(2r)} & \searrow & & \searrow
 \end{array}$$

g_3 is epic by Lemma 6. So $\zeta(2r)$ is epic and $f(2r)$ is monic. By desuspending (and by Lemma 4 for odd q), we have $f(q): A(q)_{p+q}(X) \rightarrow MU(q)_{p+q}(X)$ is monic for all p and q . By our claim this proves $\text{hom dim}_{MU_*} MU_*(X^p) \leq 2$.

The $n=1$ case. There is a degree 0 morphism of spectra $\mu: MU \rightarrow K(Z)$ ($K(Z)$ is the Eilenberg-Mac Lane spectrum) inducing the Thom homomorphism $\mu: MU_*() \rightarrow H_*(; Z)$. $\text{hom dim}_{MU_*} MU_*(X) \leq 1$ if and only if μ is epic for X a finite complex [3, 3.11]. Now one can mimic the proof of the $n=2$ case. Q.E.D.

PROOF OF THEOREM 2. (i) implies (ii). Since $k(2r+1)_n(X) \cong k(2r)_n(X) \cong k_{n-2r}(X)$ (Lemma 4 and the proof of Lemma 6), we have that $k(q)_{p+q}(X)$ is free abelian for all p and q . The exact sequence of the pair (X^p, X^{p-1}) induces the exact sequence:

$$0 \rightarrow k(q)_{p+q}(X) \rightarrow k_{p+q}(X^p) \rightarrow k_{p+q}(X^p, X^{p-1}).$$

$k_{p+q}(X^p, X^{p-1})$ and $k(q)_{p+q}(X)$ are free abelian; so $k_{p+q}(X^p)$ is also.

(ii) implies (iii). Since $k_*(X^p)$ is free abelian, $\text{hom dim}_{MU_*} MU_*(X^p) \leq 1$ [5]. So the Stong-Hattori homomorphism

$$sh: MU_*(X^p) \rightarrow k_*(MU \wedge X^p) \cong k_*(MU) \otimes_{Z[t]} k_*(X^p)$$

is monic [9]. $k_*(MU)$ is a free $Z[t]$ -module; so $k_*(MU \wedge X^p)$ is free abelian.

(iii) implies (iv). $MU(q)_{p+q}(X)$ is a subgroup of the free abelian $MU_{p+q}(X^p)$; so the monomorphism modulo torsion $MU(q)_{p+q}(X) \rightarrow MU(q-1)_{p+q}(X)$ is monic. By Lemma 3, the Atiyah-Hirzebruch-Dold spectral sequence for $MU_*(X)$ collapses. By [3, 3.11], this implies $\text{hom dim}_{MU_*} MU_*(X) \leq 1$.

(iv) implies (i). Since $\text{hom dim}_{MU_*} MU_*(X) \leq 1$, there is a monomorphism $m_i: k_{2n}(X) \rightarrow k_{2n+2}(X)$ [5]. We may consider $k_{2n}(X)$ as a subgroup of $K_0(X) = k_{2n+2r}(X)$ for r large. But $K_0(X)$ is a direct summand of $MU_0(X)$

[2, §9]; so $k_{2n}(X)$ is free abelian. Repeat the argument to get $k_{2n-1}(X) \cong \bar{k}_{2n}(\Sigma X^+)$ free abelian. Q.E.D.

REFERENCES

1. J. F. Adams, *On Chern characters and the structure of the unitary group*, Proc. Cambridge Philos. Soc. **57** (1961), 189–199. MR **22** #12525.
2. P. E. Conner and E. E. Floyd, *The relation of cobordism to K-theories*, Lecture Notes in Math., no. 28, Springer-Verlag, Berlin, 1966. MR **35** #7344.
3. P. E. Conner and L. Smith, *On the complex bordism of finite complexes*, Inst. Hautes Études Sci. Publ. Math. **37** (1970), 117–221.
4. A. Dold, *On general cohomology*, Lecture Notes, Nordic Summer School in Math., Matematisk Institut, Aarhus Universitet, Aarhus, 1968. MR **40** #8045.
5. D. C. Johnson and L. Smith, *On the relation of complex cobordism to connective K-theory*, Proc. Cambridge Philos. Soc. **70** (1971), 19–22.
6. P. S. Landweber, *On the complex bordism and cobordism of infinite complexes*, Bull. Amer. Math. Soc. **76** (1970), 650–654. MR **41** #2668.
7. S. Mac Lane, *Homology*, Die Grundlehren der math. Wissenschaften, Band 114, Academic Press, New York; Springer-Verlag, Berlin, 1963. MR **28** #122.
8. L. Smith, *On the relation of connective k-theory to homology*, Proc. Cambridge Philos. Soc. **68** (1970), 637–640.
9. ———, *A note on the Hattori-Stong theorem*, Illinois J. Math. (to appear).
10. R. Vogt, *Boardman's stable homotopy category*, Lecture Notes Series, vol. 21, Matematisk Institut, Aarhus Universitet, Aarhus, 1970.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506