

K_1 OF A COMMUTATIVE VON NEUMANN REGULAR RING

LESLIE G. ROBERTS

ABSTRACT. Let A be a commutative regular ring (in the sense of von Neumann), and let q be an ideal in A . Then $K_1(A, q) = U(A, q)$.

A ring A is called regular (in the sense of von Neumann) if for all $a \in A$, there exists $s \in A$ such that $a = asa$. Some properties of these rings are discussed in [2, Chapter I, §2, Exercises 16, 17, and Chapter II, §4, Exercise 16]. We will consider only A commutative, with unit. Let q be an ideal in A . Then we show that $K_1(A, q) = U(A, q)$, where $U(A, q)$ denotes the units of A which are congruent to 1 mod q . The groups $K_1(A, q)$ are thus as simple as possible, as might be expected since A is zero dimensional. If A has a finite number of ideals, then A is the direct product of a finite number of fields, and these results are well known, for example, by Theorem 9.1, p. 266, of [1]. However, if A has an infinite number of ideals, then our results seem to be new. By [1, Chapter 5, §2], the determination of $K_1(A, q)$ is equivalent to the determination of all normal subgroups of $GL(A)$. The notation is as in Chapter 5 of [1]. The result where $q \neq A$ was suggested by the referee.

We first consider the case $q = A$, where we write $K_1(A, A) = K_1(A)$. There is a surjective homomorphism $K_1(A) \rightarrow U(A)$ and this can be split by mapping $U(A) \rightarrow GL_1(A)$. Write $K_1(A) = U(A) \oplus SK_1(A)$. In order to prove that $SK_1(A) = 0$ it is sufficient to prove (in view of the Whitehead Lemma, p. 226 of [1]) that if $a \in GL_n(A)$, then there exist b and c in $E_n(A)$ such that bac is a diagonal matrix.

First I will make an observation that is valid for any ring A . Suppose $A = A_1 \times A_2 \times \cdots \times A_m$. Then $GL_n(A) = GL_n(A_1) \times GL_n(A_2) \times \cdots \times GL_n(A_m)$, and under this isomorphism $E_n(A) = E_n(A_1) \times \cdots \times E_n(A_m)$. If $a \in GL_n(A)$, then a corresponds to (a_1, a_2, \cdots, a_m) , where $a = a_1 + a_2 + \cdots + a_m$, and the matrix a_i has coefficients in A_i .

Assume again that A is a commutative regular ring (in the sense of von Neumann). I will prove by induction on n that if $a \in GL_n(A)$, then there exist $b, c \in E_n(A)$ such that bac is diagonal.

Received by the editors February 3, 1971.

AMS 1970 subject classifications. Primary 13D15, 18F30.

Key words and phrases. K_1 , von Neumann regular ring.

© American Mathematical Society 1972

Let a_{11} be the entry in the upper left hand corner of a . Since $a_{11}A$ is generated by an idempotent, we have a direct product decomposition $A = A_1 \times S_1$, such that a_{11} projects to a unit a_{11} in A_1 , and to zero in S_1 . Write $a = a_1 + b_1$, with $a_1 \in GL_n(A_1)$ and $b_1 \in GL_n(S_1)$. Let b_{21} be the entry in the second row, first column of b_1 . Write $S_1 = A_2 \times S_2$, where b_{21} projects to a unit in A_2 and to zero in S_2 . We then have $a = a_1 + a_2 + b_2$, with $a_i \in GL_n(A_i)$, $i = 1, 2$, and $b_2 \in GL_n(S_2)$, and b_2 has the first two entries in the first column zero. Continue down the first column in this manner. Eventually we get a decomposition $A = A_1 \times A_2 \times \cdots \times A_n$, and $a = a_1 + a_2 + \cdots + a_n$, where $a_i \in GL_n(A_i)$ and a_i has a unit (in A_i) as the i th entry in the first column. (Some of the rings A_i might be trivial.) By using elementary row and column transformations over each A_i , we reduce a to the form where it has a unit in the upper left hand corner, and all other entries in the first row and column are zero. The proof is now completed by induction on n .

Now we consider the case where $q \neq A$. By definition, $K_1(A, q) = GL(A, q)/E(A, q)$, where $GL(A, q) = \ker(GL(A) \rightarrow GL(A/q))$ and $E(A, q)$ is the normal subgroup of $E(A)$ generated by the q -elementary matrices. The sequence $0 \rightarrow E(A, q) \rightarrow E(A) \rightarrow E(A/q) \rightarrow 0$ is not exact for all A and q since $x \in \ker(E(A) \rightarrow E(A/q))$ need not lie in $E(A, q)$. It is, however, exact for the A considered here. First of all, it is exact if q is of finite type (hence idempotent generated) by Proposition 1.5, p. 451 of [1].

Suppose $x \in \ker(E(A) \rightarrow E(A/q))$. Then there is an ideal q' of finite type, with $q' \subset q$ and $x \in \ker(E(A) \rightarrow E(A/q'))$. Then $x \in E(A, q') \subset E(A, q)$. Hence we have an exact sequence $0 \rightarrow E(A, q) \rightarrow E(A) \rightarrow E(A/q) \rightarrow 0$. It now follows from the serpent diagram (p. 17 of [2]) that $0 \rightarrow K_1(A, q) \rightarrow K_1(A) \rightarrow K_1(A/q)$ is exact. Since A/q is also a von Neumann regular ring, we have that $K_1(A, q) = \ker(U(A) \rightarrow U(A/q))$ and this by definition equals $U(A, q)$.

BIBLIOGRAPHY

1. H. Bass, *Algebraic K-theory*, Benjamin, New York, 1968. MR 40 #2736.
2. N. Bourbaki, *Eléments de mathématique*. Fasc. XXVII. *Algèbre commutative*. Chap. 1: *Modules plats*. Chap. 2: *Localisation*, Actualités Sci. Indust., no. 1290, Hermann, Paris, 1961. MR 36 #146.

DEPARTMENT OF MATHEMATICS, QUEEN'S UNIVERSITY, KINGSTON, ONTARIO, CANADA