

## TWO PROPERTIES OF $R^N$ WITH A COMPACT GROUP TOPOLOGY

KEVIN J. SHARPE<sup>1</sup>

**ABSTRACT.** We let  $R_c^N$  be a compact additive group, and we prove that if  $A$  is an  $R_c^N$ -measurable set, then one of the interiors of  $A$  and  $A'$  in the usual topology for  $R^N$  (written  $R_u^N$ ) must be void. Also we show that the only functions from  $R^N$  to a Hausdorff space that are both  $R_u^N$ -continuous and  $R_c^N$ -measurable are the constant functions.

**1. Introduction.** If  $N$  is a positive integer and  $R_c^N$  a compact additive group (for the existence of which, see, for example, Halmos [1]), we can answer two questions (posed by Hawley [2] for  $R^1$ ):

(i) If  $A$  and  $A'$  have nonvoid interiors in the usual topology for  $R^N$  (written as  $R_u^N$ ), must  $A$  be  $R_c^N$ -nonmeasurable?

(ii) If  $f$  is  $R_u^N$ -continuous and  $R_c^N$ -continuous, must  $f$  be a constant function?

**2. Preliminaries.** If we let  $n$  be a positive integer, then the map sending  $x$  to  $nx$ , for each element  $x$  of  $R^N$ , is  $R^N$ -continuous as is also its inverse [5, p. 96, A] and [4, p. 141]. It now follows that  $n\mathcal{M} = \mathcal{M}$ , where  $\mathcal{M}$  is the set of all  $R_c^N$ - (Borel-) measurable sets.

We have then the following lemma, the first part of which is trivial:

**LEMMA 1.** *All translations and positive-integer multiples of  $R_c^N$ -measurable sets are again  $R_c^N$ -measurable.*

**3. Question (i).** As  $R_c^N$  is a compact group, we can construct on it a unique normalised Haar measure  $\mu$ . We define, for each positive integer  $n$ ,  $\mu^n(A)$  to be  $\mu(nA)$  for all  $R_c^N$ -measurable sets  $A$ .

**THEOREM 2.** *For each positive integer  $n$ ,  $\mu^n$  is identical to  $\mu$ .*

---

Received by the editors November 15, 1971.

AMS 1970 subject classifications. Primary 22C05; Secondary 28A05, 43A05, 54C05.

Key words and phrases. Compact group topologies for  $R^N$ , continuous functions on  $R^N$  with a compact group topology, Haar measure, measurable sets of a compact group topology for  $R^N$ .

<sup>1</sup> This work is part of a Ph.D. dissertation supervised by Dr. G. Elton of La Trobe University, Melbourne, Australia.

© American Mathematical Society 1972

PROOF. For  $n$  a positive integer,  $\mu^n$  is indeed a Haar measure, since, for instance,

$$\mu^n(A + x) = \mu(nA + nx) = \mu(nA) = \mu^n(A),$$

for any  $R_c^N$ -measurable set  $A$ , and element  $x$  of  $R^N$ . However, for each  $n$ ,

$$\mu^n(R^N) = \mu(nR^N) = \mu(R^N) = 1;$$

hence  $\mu^n = \mu$ .

The answer to question (i) is evident from the following theorem:

**THEOREM 3.** *An  $R_c^N$ -measurable set having nonvoid  $R_u^N$ -interior has Haar measure one for  $R_c^N$ .*

PROOF. Let  $A$  be an  $R_c^N$ -measurable set whose  $R_u^N$ -interior is nonvoid. Thus there is a nonvoid  $R_u^N$ -open ball in  $A$ , and we shift  $A$  (giving us  $B$ ) so that the ball,  $C$ , is centred on  $\mathbf{0}$ ; by Lemma 1,  $B$  is  $R_c^N$ -measurable and has the same measure as  $A$ . It is easy to show that, for positive integers  $n$ ,  $\bigcup_{n=1}^\infty nC = \lim_{n \rightarrow \infty} nC = R^N$ ; but  $nC \subseteq nB$  for each  $n$  means that  $\lim_{n \rightarrow \infty} nB = R^N$  ( $n \in Z^+$ ). Now by Lemma 1,  $nB$  is  $R_c^N$ -measurable for each positive integer  $n$ , and so the characteristic functions  $\chi_{nB}$  will be  $R_c^N$ -measurable. We have  $\lim_{n \rightarrow \infty} \chi_{nB} = \chi_{R^N}$  ( $n \in Z^+$ ), the integral  $\int_{R^N} \chi_{R^N} d\mu = 1$ , and  $|\chi_{nB}(x)| \leq \chi_{R^N}(x)$  for any  $x$  in  $R^N$  and positive integer  $n$ , allowing us to apply the Lebesgue Dominated Convergence Theorem to give

$$\lim_{n \rightarrow \infty} \int_{R^N} \chi_{nB} d\mu = \int_{R^N} \chi_{R^N} d\mu \quad (n \in Z^+).$$

Therefore, by Theorem 2,  $1 = \lim_{n \rightarrow \infty} \mu(nB) = \mu(B)$  ( $n \in Z^+$ ), and so  $\mu(A) = 1$ .

**4. Question (ii).** The answer to this question follows rather simply from that for its predecessor:

**THEOREM 4.** *The constant functions are the only maps from  $R^N$  to a Hausdorff space that are both  $R_u^N$ -continuous and  $R_c^N$ -measurable.*

PROOF. Let  $f$  be a  $R_u^N$ -continuous and  $R_c^N$ -measurable function to a Hausdorff space  $T$ , but which is not a constant function, and  $I_1$  and  $I_2$  be two nonvoid, disjoint subsets of  $T$  which are open relative to the image of  $R^N$  under  $f$ . The inverse maps of  $I_1$  and  $I_2$  under  $f$  are nonvoid, disjoint open subsets of  $R_u^N$  which are  $R_c^N$ -measurable, making the normalised Haar measure of  $R_c^N$  at least two.

## REFERENCES

1. P. R. Halmos, *Comment on the real line*, Bull. Amer. Math. Soc. **50** (1944), 877–878. MR **6**, 145.
2. D. N. Hawley, *Compact group topologies for  $R$* , Proc. Amer. Math. Soc. **30** (1971), 566–572.
3. E. Hewitt and K. A. Ross, *Abstract harmonic analysis*. Vol. I: *Structure of topological groups. Integration theory, group representations*, Die Grundlehren der math. Wissenschaften, Band 115, Academic Press, New York; Springer-Verlag, Berlin, 1963. MR **28** #158.
4. J. L. Kelly, *General topology*, Van Nostrand, Princeton, N.J., 1955. MR **16**, 1136.
5. L. S. Pontrjagin, *Topological groups*, 2nd ed., GITTL, Moscow, 1954; English transl., Gordon and Breach, New York, 1966. MR **17**, 171; MR **34** #1439.

DEPARTMENT OF MATHEMATICS, LA TROBE UNIVERSITY, MELBOURNE, AUSTRALIA