

EQUIVALENCE OF ALGEBRA NORMS ON $C(G)$

KJELD B. LAURSEN

In [1] Slobko shows that if G is a compact first countable group, if S_a denotes the translation operator on the algebra of continuous functions on G (with pointwise multiplication),

$$S_a f(x) = f(ax), \quad f \in C(G),$$

and if $|\cdot|$ is an algebra norm on $C(G)$ with respect to which the mapping $a \rightarrow S_a f$ is continuous for each f in $C(G)$ then $|\cdot|$ and the sup norm $\|\cdot\|$ are equivalent. The following simple argument shows that the countability assumption is redundant.

Define

$$\|f\| = \sup_{a \in G} |S_a f|, \quad f \in C(G).$$

Then since G is compact this defines an algebra norm on $C(G)$ and since

$$\begin{aligned} \|S_a\| &= \sup_{\|f\| \leq 1} \|S_a f\| = \sup_{\|f\| \leq 1} \sup_{b \in G} |S_b S_a f| \\ &= \sup_{\|f\| \leq 1} \sup_{a \in G} |S_a f| = 1, \end{aligned}$$

each mapping $S_a: C(G) \rightarrow C(G)$ is continuous with respect to $\|\cdot\|$. Hence by Theorem 1 of [1], $\|\cdot\|$ and $\|\cdot\|$ are equivalent. But clearly $\|f\| \geq |f|$, so $|\cdot|$ and $\|\cdot\|$ are equivalent.

REFERENCE

1. T. A. Slobko, *Algebra norms on $C(G)$* , Amer. J. Math. **92** (1970), 381–388.

UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024

Received by the editors October 18, 1971.

AMS 1970 subject classifications. Primary 46J10.