

COMPLEMENTING SETS OF n -TUPLES OF INTEGERS

MELVYN B. NATHANSON¹

ABSTRACT. Let S, A_1, A_2, \dots, A_p be finite nonempty sets of n -tuples of integers such that, if $a_i \in A_i$ for $i=1, 2, \dots, p$, then $a_1+a_2+\dots+a_p \in S$, and such that every $s \in S$ has a unique representation as a sum $s=a_1+a_2+\dots+a_p$ with $a_i \in A_i$. If S is the cartesian product of n sets of integers, then each A_i is also the cartesian product of n sets of integers, and conversely.

Let S, A_1, A_2, \dots, A_p be sets of n -tuples of integers. Define addition of n -tuples componentwise. Then S is the *sum* of A_1, A_2, \dots, A_p , denoted $S=A_1+A_2+\dots+A_p$, if $S=\{a_1+a_2+\dots+a_p \mid a_i \in A_i \text{ for } i=1, 2, \dots, p\}$. If S is the sum of A_1, A_2, \dots, A_p , and if for each $s \in S$ there exist unique n -tuples $a_i \in A_i$ such that $s=a_1+a_2+\dots+a_p$, then A_1, A_2, \dots, A_p are called *complementing sets* for S , denoted $S \cong A_1+A_2+\dots+A_p$. A set of n -tuples is *proper* if it is the cartesian product of n sets of integers. For positive integers u and v , let $S=\{0, 1, 2, \dots, u\} \times \{0, 1, 2, \dots, v\}$. If A_1 and A_2 are subsets of S such that $S \cong A_1+A_2$, then Hansen [1] proved that A_1 and A_2 are proper. This result extends to the general case of arbitrary n and p , and S the cartesian product of any n finite sets of integers.

THEOREM. Let S, A_1, A_2, \dots, A_p be finite nonempty sets of n -tuples of integers such that $S \cong A_1+A_2+\dots+A_p$. Then S is proper if and only if each A_i is proper.

PROOF. Suppose that S is proper. By translation, it is enough to consider only the case in which all coordinates of all n -tuples of S, A_1, \dots, A_p are nonnegative integers. Let $Z[X_1, \dots, X_n]$ be the ring of polynomials in n variables with integral coefficients. Define F, G_1, G_2, \dots, G_p in $Z[X_1, \dots, X_n]$ by

$$F = \sum_{(s_1, s_2, \dots, s_n) \in S} X_1^{s_1} X_2^{s_2} \dots X_n^{s_n},$$

$$G_i = \sum_{(a_1, a_2, \dots, a_n) \in A_i} X_1^{a_1} X_2^{a_2} \dots X_n^{a_n}.$$

Received by the editors July 30, 1971.

AMS 1970 subject classifications. Primary 10L05; Secondary 10A45, 10J99, 05A15.
Key words and phrases. Complementing sets, sumsets of integers, addition of n -tuples of integers.

¹Supported in part by an NSF Predoctoral Traineeship from the University of Rochester.

© American Mathematical Society 1972

Since S is proper, there exist finite sets of nonnegative integers S_1, S_2, \dots, S_n such that $S = S_1 \times S_2 \times \dots \times S_n$. Then

$$(1) \quad F = \prod_{j=1}^n \left(\sum_{s_j \in S_j} X_j^{s_j} \right).$$

The polynomial F is the product of irreducible polynomials, and by (1), each of these irreducibles is a polynomial in exactly one variable. Each G_i is also a product of irreducible polynomials. Since $A_1 + A_2 + \dots + A_p \cong S$, it follows that $F = G_1 G_2 \dots G_p$. Since $Z[X_1, \dots, X_n]$ is a unique factorization domain, each irreducible factor of each G_i is a factor of F , and so is a polynomial in exactly one variable. Thus there exist polynomials $g_{ij} \in Z[X_j]$ for $i=1, 2, \dots, p$ and $j=1, 2, \dots, n$ such that $G_i = g_{i1} g_{i2} \dots g_{in}$. Let A_{ij} be the finite set of nonnegative integers which are the powers of X_j occurring with nonzero coefficient in g_{ij} . Then $A_i = A_{i1} \times A_{i2} \times \dots \times A_{in}$ for $i=1, 2, \dots, p$, and A_i is proper. (Moreover, $A_{1j} + A_{2j} + \dots + A_{pj} \cong S_j$ for $j=1, 2, \dots, n$.)

Conversely, suppose that each A_i is proper. Then there exist finite sets of integers A_{ij} for $i=1, 2, \dots, p$ and $j=1, 2, \dots, n$ such that $A_i = A_{i1} \times A_{i2} \times \dots \times A_{in}$. Let $S_j = A_{1j} + A_{2j} + \dots + A_{pj}$ for $j=1, 2, \dots, n$. Then $S = S_1 \times S_2 \times \dots \times S_n$, and so S is proper.

REMARK. The theorem is false if S is the cartesian product of infinite sets. If N is the set of nonnegative integers, there exist sets A_1 and A_2 which are not proper but satisfy $A_1 + A_2 \cong N \times N$. Hansen [1] and Niven [2] have determined all sets A_1 and A_2 such that $A_1 + A_2 \cong N \times N$.

REFERENCES

1. R. T. Hansen, *Complementing pairs of subsets of the plane*, Duke Math. J. **36** (1969), 441–449. MR **39** #5719.
2. I. Niven, *A characterization of complementing sets of pairs of integers*, Duke Math. J. **38** (1971), 193–203. MR **43** #179.

UNIVERSITY OF ROCHESTER, ROCHESTER, NEW YORK 14627

Current address: Southern Illinois University, Carbondale, Illinois 62901