

EMBEDDING NUCLEAR SPACES IN PRODUCTS OF AN ARBITRARY BANACH SPACE

STEPHEN A. SAXON

ABSTRACT. It is proved that if E is an arbitrary nuclear space and F is an arbitrary infinite-dimensional Banach space, then there exists a fundamental (basic) system \mathcal{V} of balanced, convex neighborhoods of zero for E such that, for each V in \mathcal{V} , the normed space E_V is isomorphic to a subspace of F . The result for $F=l_p$ ($1 \leq p \leq \infty$) was proved by A. Grothendieck.

This paper is an outgrowth of an interest in varieties of topological vector spaces [2] stimulated by J. Diestel and S. Morris, and is in response to their most helpful discussions and questions. The main theorem, valid for arbitrary infinite-dimensional Banach spaces, was first proved by A. Grothendieck [3] (also, see [5, p. 101]) for the Banach spaces l_p ($1 \leq p \leq \infty$) and later by J. Diestel for the Banach space c_0 .

Our demonstration relies on two profound results of T. Kōmura and Y. Kōmura [4] and C. Bessaga and A. Pełczyński [1], respectively:

(i) A locally convex space is nuclear if and only if it is isomorphic to a subspace of a product space $(s)^I$, where I is an indexing set and (s) is the Fréchet space of all rapidly decreasing sequences.

(ii) Every infinite-dimensional Banach space contains a closed infinite-dimensional subspace which has a Schauder basis.¹

Recall that for a balanced, convex neighborhood V of zero in a locally convex space E , E_V is a normed space which is norm-isomorphic to $(M, p|_M)$, where p is the gauge of V and M is a maximal linear subspace of E on which p is a norm; \tilde{E}_V is the completion of E_V . Denote by (s) the nuclear Fréchet space of rapidly decreasing sequences, so that

$$(s) = \left\{ (\lambda_n) : \sup_n |n^k \lambda_n| < \infty, k = 1, 2, \dots \right\},$$

Received by the editors February 22, 1971 and, in revised form, October 25, 1971.
AMS 1970 subject classifications. Primary 46A05, 46A35, 46B15.

Key words and phrases. Nuclear spaces, (s) , \tilde{E}_V , product spaces, varieties, norm-isomorphic, Schauder basis.

¹ This result was stated without proof in Banach's book (1932).

© American Mathematical Society 1972

with the system of seminorms

$$\left\{ p_k((\lambda_n)) = \sup_n |n^k \lambda_n| : k = 1, 2, \dots \right\}.$$

LEMMA 1. *Let F be an arbitrary infinite-dimensional Banach space with a Schauder basis, and let U be a neighborhood of zero in (s) . There exists a balanced, convex neighborhood of zero $V \subset U$ such that $(s)_V$ is norm-isomorphic to F .*

PROOF. Let $\{x_n\}$ be a Schauder basis for F with coefficient functionals $\{f_n\} \subset F'$, and with $\|x_n\|=1$ ($n=1, 2, \dots$). By the uniform boundedness theorem, the (absolute) polar $\{f_n\}^0$ of $\{f_n\}$ is a neighborhood of zero, and its gauge p is thus a continuous seminorm. Therefore $K \|\sum_{n=1}^\infty a_n x_n\| \geq p(\sum_{n=1}^\infty a_n x_n) = \sup_n |a_n|$ for each $\sum_{n=1}^\infty a_n x_n \in F$, where K is some positive constant. For some $\varepsilon > 0$ and positive integer k ,

$$V' = \{x \in (s) : p_k(x) \leq \varepsilon\} \subset U.$$

Define a norm q on (s) so that, for each $(\lambda_n) \in (s)$,

$$\begin{aligned} K \left(\sum_{n=1}^\infty n^{-2} \right) p_{k+2}((\lambda_n)) &= K \left(\sum_{n=1}^\infty n^{-2} \right) \sup_n |n^{k+2} \lambda_n| \geq K \sum_{n=1}^\infty |n^k \lambda_n| \\ (*) \qquad \qquad \qquad &= K \sum_{n=1}^\infty \|n^k \lambda_n x_n\| \geq K \left\| \sum_{n=1}^\infty n^k \lambda_n x_n \right\| \\ &\equiv q((\lambda_n)) \geq p \left(\sum_{n=1}^\infty n^k \lambda_n x_n \right) = p_k((\lambda_n)). \end{aligned}$$

Therefore $V = \{x \in (s) : q(x) \leq \varepsilon\} \subset V' \subset U$ is a neighborhood of zero, and $(s)_V$ is norm-isomorphic to $((s), q)$, which is clearly norm-isomorphic to a dense subspace of F by (*).

REMARK. It is clear from the proof that the map

$$(\lambda_n) \rightarrow \left(\sum_{n=1}^\infty n^k \lambda_n x_n \right)_{k=1}^\infty,$$

embeds (s) in the product space $F \times F \times F \times \dots$, and thus by (i) and (ii), each nuclear space can be embedded in some product of any given infinite-dimensional Banach space.

LEMMA 2. *The space (s) is isomorphic to $(s) \times (s)$.*

PROOF. The map $(\lambda_n) \rightarrow ((\lambda_{2n-1}), (\lambda_{2n}))$ is a linear, bicontinuous bijection from (s) onto $(s) \times (s)$.

THEOREM. *Let E be an arbitrary nuclear space and F an arbitrary infinite-dimensional Banach space. Each neighborhood U of zero in E contains a balanced, convex neighborhood V of zero such that E_V is norm-isomorphic to a subspace of F .*

PROOF. Let I be any indexing set, and U a neighborhood of zero in the product space $E=(s)^I$. Thus there exists a finite subset A of I such that $U \supset W \times (s)^B$, where $B=I \setminus A$ and W is a neighborhood of zero in $(s)^A$. By (ii) there exists a closed infinite-dimensional subspace F_0 of F which has a Schauder basis, and by Lemmas 1 and 2, there is a balanced, convex neighborhood of zero $V \subset W$ such that the completion of $((s)^A)_V$ is norm-isomorphic to F_0 . Now $V' = V \times (s)^B$ is a neighborhood of zero contained in U such that $E_{V'}$ is norm-isomorphic to $((s)^A)_V$ and the conclusion of the theorem holds for $E=(s)^I$. But then the conclusion clearly holds for any subspace of $(s)^I$, and hence for any nuclear space E , by (i).

COROLLARY. *The variety ([2], [2a]) generated by any infinite-dimensional Banach space contains the variety of all nuclear spaces.*

The author thanks Professor Pełczyński for the privilege and benefit of discussing this material with him.

REFERENCES

1. C. Bessaga and A. Pełczyński, *On bases and unconditional convergence of series in Banach spaces*, *Studia Math.* **17** (1958), 151–164. MR **22** #5872.
2. J. Diestel, Sidney A. Morris and S. Saxon, *Varieties of locally convex topological vector spaces*, *Bull. Amer. Math. Soc.* **77** (1971), 799–803.
- 2a. ———, *Varieties of linear topological spaces*, *Trans. Amer. Math. Soc.* (to appear).
3. A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, *Mem. Amer. Math. Soc. No. 16* (1955). MR **17**, 763.
4. T. Kōmura and Y. Kōmura, *Über die Einbettung der nuklearen Räume in $(s)^A$* , *Math. Ann.* **162** (1965/66), 284–288. MR **32** #6190.
5. H. H. Schaefer, *Topological vector spaces*, Macmillan, New York, 1966. MR **33** #1738.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32603