

THE WHITNEY TECHNIQUE FOR POINCARÉ COMPLEX EMBEDDINGS

J. P. E. HODGSON

ABSTRACT. The following result is proved.

THEOREM. *Let $f: K^k \rightarrow X^n$ be a map of a finite connected k -dimensional Poincaré complex K to a finite simply-connected n -dimensional Poincaré complex X , then if $n \geq 2k + 1$, $k > 2$, f is homotopic to an embedding. The proof involves a technique akin to the Whitney procedure for eliminating double points.*

The object of this note is to prove the following result.

THEOREM 1. *Let $f: K^k \rightarrow X^n$ be a map between a finite connected k -dimensional Poincaré complex K and a simply-connected finite n -dimensional Poincaré complex X , then if $n \geq 2k + 1$ and $k > 2$, f is homotopic to an embedding.*

This improves the result of Levitt [3] by one dimension but at the cost of requiring X to be simply-connected. The method of proof is to follow Levitt's construction, introducing a 'singularity' in K to kill the obstruction to engulfing. Then by a Whitney process we can remove the singularity.

1. Construction of the stable model. Theorem 1 is a corollary of the following result.

LEMMA 2. *Let (X^{2n+1}, Y^{2n}) be a finite Poincaré pair of dimension $2n + 1$, with $\pi_1(Y) = 0$, then any map $f: (D^n, S^{n-1}) \rightarrow (X, Y)$ is homotopic to an embedding.*

This gives Theorem 1, by embedding the $(n-1)$ -skeleton of K in X using Levitt's theorem [3]; then by Wall [4] we have only the top cell left to embed. We note also that Levitt's theorem allows us to assume that $f|_{S^{n-1}}$ is an embedding.

The proof of Lemma 2 is modelled on that of Lemma 5.1 in [3], since modifications are required at several points we recapitulate in some detail.

Received by the editors March 12, 1971 and, in revised form, November 15, 1971.

AMS 1970 subject classifications. Primary 57B10.

Key words and phrases. Poincaré complex, splitting of Poincaré complexes, engulfing.

© American Mathematical Society 1972

First let $g: (X^{2n+1}, Y) \rightarrow (P^{2n+j+1}, Q^{2n+j})$ be a codimension j PL thickening, with $\partial P = Q \cup R$ and $\partial Q = \partial R$. Note that in homotopy terms we can think of (P, Q) as the mapping cone of a spherical fibration by [1, Chapter I, §4].

Now by general position we may homotope the composition

$$j \circ f: (D^n, S^{n-1}) \rightarrow (P, Q)$$

to a PL embedding: further by the remarks above we have a map

$$F_0: (D^n \times D^j, S^{n-1} \times D^j; D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (P, Q, R, \partial R)$$

obtained by considering the restriction of the spherical fibration to (D^n, S^{n-1}) .

Let (M, N) be a regular neighborhood of (D^n, S^{n-1}) in (P, Q) and $(\mathcal{M}, \mathcal{N})$ the mapping cylinder of the composition

$$\tilde{p}: (D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \xrightarrow{P} (D^n, S^{n-1}) \subset (M, N);$$

then F_0 can be extended to $F: (\mathcal{M}, \mathcal{N}) \rightarrow (P, Q)$ so that $F|(M, N)$ is an inclusion. Further if \tilde{p} is altered by a homotopy in (M, N) , then we can change F in such a way that $F|(D^n \times S^{j-1}, S^{n-1} \times S^{j-1})$ is unchanged and so that $F|(M, N)$ is still the inclusion. This follows from the fact that general position shows that a homotopy of $\text{Int}(D^n \times S^{j-1})$ (resp. $S^{n-1} \times S^{j-1}$) in M (resp. N) will miss $\text{Int } D^n$ (resp. S^{n-1}) so that we can achieve the following.

$$(GP) \quad F(\mathcal{M} - M) \subseteq P - M, \quad F(\mathcal{N} - N) \subset Q - N.$$

From (GP) we obtain a map of pairs

$$\tilde{p}: (D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (\partial M - \text{Int } N, \partial N)$$

and by the remarks following Lemma 2, we may assume $\tilde{p}|S^{n-1} \times S^{j-1}$ is an embedding. Let B be a regular neighborhood of its image in ∂N , then

$$\tilde{p}: (D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (\partial M - \text{Int } N, B)$$

is a $(j-1)$ -connected map. Thus by Theorem 2.3 of [2] we can find $(A^{2n+j}, B^{2n+j-1}) \subset (\partial M - \text{Int } N, \partial N)$ and a simple homotopy equivalence

$$q: (D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (A, B)$$

such that q followed by the inclusion $(A, B) \subset (\partial M - \text{Int } N, \partial N)$ is homotopic $\text{rel}(S^{n-1} \times S^{j-1})$ to \tilde{p} . Now we can apply Theorem 5.2 of [3] to the map q to change \tilde{p} by a homotopy and get $\tilde{p}(D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \subseteq (W, V)$ where $(W, V) \subset (\partial M - \text{Int } N, \partial N)$ is a pair of dimension $(n+j-1, n+j-2)$ and the map

$$(D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (W, V)$$

is a homotopy equivalence. Now let (W_0, V_0) be a disjoint copy of (W, V) and let $i: (W_0, V_0) \rightarrow (M, N)$ be the map corresponding to the inclusion $(W, V) \subset (M, N)$. Then the homotopy equivalence

$$(D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (W, V)$$

induces a homotopy equivalence

$$\alpha: (\mathcal{M}, \mathcal{N}, D^n \times S^{j-1}, S^{n-1} \times S^{j-1}) \rightarrow (\mathcal{M}_i, \mathcal{N}_i, W_0, V_0)$$

where $(\mathcal{M}_i, \mathcal{N}_i)$ is the mapping cylinder of $i: (W_0, V_0) \rightarrow (M, N)$. Thus we can get a map

$$F: (\mathcal{M}_i, \mathcal{N}_i, W_0, V_0) \rightarrow (P, Q, R, \partial R)$$

such that $F \circ \alpha$ is homotopic to F as a map of the 4-tuple $(\mathcal{M}, \mathcal{N}, D^n \times S^{j-1}, S^{n-1} \times S^{j-1})$ into $(P, Q, R, \partial R)$ and with $F|(M, N)$ still the inclusion. Let (T, U) be the part of $(\mathcal{M}_i, \mathcal{N}_i)$ corresponding to $(W_0, V_0) \times I$; that is $T = \text{cl}(\mathcal{M}_i - M)$, $U = \text{cl}(\mathcal{N}_i - N)$.

Now by (GP) we can suppose

$$F(T) \subseteq P - (\text{Int } M \cup \text{Int } N) \quad \text{and} \quad F(U) \subseteq Q - \text{Int } N.$$

Set $P_0 = P - (\text{Int } M \cup \text{Int } N)$, $Q_0 = Q - \text{Int } N$ so $F(T, U) \subset (P_0, Q_0)$, and this implies that the pair $(W, V) \subset (\partial M - \text{Int } N, \partial N)$ can be deformed into $(R, \partial R)$ keeping V in Q_0 and W in P_0 .

This completes the construction of the stable model.

2. Engulfing. In Levitt's original proof of the embedding theorem [3], the transition from the stable model to the required embedding was effected by engulfing (W, V) in a collar on $(R, \partial R)$, a procedure which corresponds to constructing a compressible embedding of $(D^n \times D^j, S^{n-1} \times D^j)$ in (P, Q) . Unfortunately, there are obstructions to doing the engulfing in our case (in fact the counterexample of Irwin shows that these will in general not vanish), so that we require a slightly different approach. We shall show how one can embed a complex (K^n, S^{n-1}) in (P, Q) where K is of the homotopy type of a wedge of S^1 's, and then we will perform surgery to eliminate the S^1 's. The reader may perhaps be helped by thinking of the S^1 's as the union of the two paths joining the double points that one has in the Whitney procedure for elimination of double points; this is in fact the spirit of the construction of K , which involves attempting to engulf W in a collar on R .

It follows from Lemma 5.3 of [3], that in Q_0 we can engulf V in a collar C , on ∂R in Q , whose inner boundary meets ∂N in a regular neighbourhood \bar{V} of V with $C_1 \cap \text{Int } N = \emptyset$. Further W is $R \cup C_1$ -inessential, but as we remarked above the counterexample of Irwin shows that we cannot in


general engulf W in a collar C on $R \cup C_1$; let us however recapitulate the proof of the engulfing theorem so as to see what modifications are possible.

We are given a homotopy of W to $R \cup C_1$ in P_0 ; choose a simplicial approximation and put this homotopy in general position; then its singular set is of dimension

$$\begin{aligned} &2(\dim \text{homotopy}) - \dim(\text{ambient space}) \\ &= 2(n + j) - (2n + j - 1) = j - 1. \end{aligned}$$

Thus the homotopy collapses onto a subcomplex of dimension j , but as (P_0, R) is only $(j-1)$ -connected we cannot deform this subcomplex into R . The idea for continuing is to ‘punch holes’ in the $(j-1)$ -simplexes of the singular set by adding 1-cells to D^k ; (geometrically this one cell represents a double point) so that the singular set collapses to dimension $(j-2)$.

For each $(j-1)$ -cell σ_i ($i=1, \dots, l$) of the singular set, let b_i be its barycentre; then there exist two disjoint paths ω_i, ω'_i from b_i to W (and thus to D^n) such that $\omega_i \cap W$ (resp. $\omega'_i \cap W$) is the barycentre of a simplex τ_i (resp. τ'_i) of maximal dimension in W , and such that the track of τ_i (resp. τ'_i) in the homotopy gives a neighbourhood of ω_i (resp. ω'_i), in the image of the homotopy (see diagram).



Now suppose we have a simplicial decomposition of P_0 so that ω_i and ω'_i are subcomplexes, and let (Z_i, Z'_i) be disjoint relative regular-neighbourhoods of $(\omega_i \cup \omega'_i)$.

Now $M' = M \cup \bigcup_{i=1}^l Z_i$ is homotopy equivalent to a complex $K = D^k \vee \bigvee_{i=1}^l S^1$ and we shall embed the pair (K, S^{k-1}) in (X, Y) . To do this using Levitt's program requires us to embed a complex $W' \subset \partial M'$, with W' of the homotopy type of the normal Spivak fibration of X in P restricted to M' , and then we must engulf W' in a collar on R . It is clear that W' is homotopy equivalent to $W \cup_{i=1}^l I \times S^{j-1}$ where the pairs $(I \times S^{j-1}, S^0 \times S^{j-1})$ are to be embedded in $(\partial Z_i - Z'_i, \partial Z_i)$. These embeddings exist by the

‘Stalling’s’ theorem used in Levitt’s paper [3, Theorem 5.2]. W' is obtained from this union by collapsing the I factors of the $S^{j-1} \times I$'s to $S^{j-1} \times 0$ in the Z_i 's.

We claim that W' can be engulfed in a collar C on $R \cup C'$, so that the inner boundary of C meets M' in a regular neighbourhood of W' . This will follow if we can show that W' can be deformed into $R \cup C'$ in $P - \text{Int } M'$ by a homotopy whose singular set is of dimension $\leq j-2$ (since once this is achieved the proof in [5] gives the required engulfing). To do this we note that we already have such a homotopy for $W' \cap W (= W - \bigcup_{i=1}^l \tau_i \cup \tau_i)$ so that it suffices to provide an extension over the $S^{j-1} \times I$. Now there is a homotopy of the $S^{j-1} \times I$ to R since the $S^{j-1} \times I$ represent part of the Spivak normal fibration of X in P , but now general position gives us the required condition on the dimension of the singular set. So the existence of C follows from engulfing this homotopy.

We thus obtain a splitting of (X, Y) mod boundary given by $(X, Y) \simeq (X_0, Y_0) \cup_{F^2} (X_1, Y_1)$ where

$$Y_2 = \text{Closure}(\overline{P - C - M'}) \cap M',$$

$$X_1 = M', \quad Y_1 = Y_2 \cup N,$$

$$X_0 = \text{Closure}(\overline{P - C - M'}), \quad Y_0 = Y_2 \cup \text{Closure}(\overline{Q - C - N}),$$

The verification that this is indeed a splitting is straightforward.

Surgery. Now choose S^1, \dots, S^l embedded in Y_2 , generating $\pi_1(X_1)$ and corresponding to the S^1, \dots, S^l in $D^n \vee \bigvee_{i=1}^l S^i$. This is possible since Levitt’s theorem tells us that a map $f: S^1 \rightarrow Y_2$ is homotopic to an embedding since $\dim Y_2 \geq 4$. But now S^1 is null homotopic in X_0 since X_0 is 1-connected by Van Kampen’s theorem (this requires $k > 2$). Hence, again by Levitt’s theorem we can get disjoint embeddings $(D_i^2, S_i^1) \rightarrow (X_0, Y_2)$ representing these null homotopies. We show how to use each of these embeddings to do surgery on X_1 and X_0 . Suppose $l=1$, and Y_2 is split as $Y_2 \simeq (A, B) \cup_B (C, B)$. Where $C \sim S^1$, then (X_0, Y_0) is split as

$$(X_0, Y_0) = (D, E) \cup_{F'} (G, H)$$

where $D \sim D^2$ and $\text{cl}(E-F) \sim C$ so we replace X_1 by $X'_1 = X_1 \cup_C D$. (X_0, Y_0) is replaced by (G, H) and Y_2 is replaced by $Y'_2 = A \cup_B F$. $Y'_1 = Y'_2 \cup \text{cl}(Y_1 - Y_2)$.

This surgery has the effect of constructing $X'_1 \sim D^n$, and thus embedding (D^n, S^{n-1}) in (X, Y) in the required homotopy class, (X, Y) being split as $(X'_1, Y'_1) \cup_{F^2} (G, H)$, thus proving the lemma.

REFERENCES

1. W. F. Browder, *Surgery of simply connected manifolds*, Notes, Princeton University, Princeton, N.J., 1969.
2. J. P. E. Hodgson, *Obstructions to concordance for thickenings*, *Invent. Math.* **5** (1968), 292–316. MR **37** #5879.
3. N. Levitt, *On the structure of Poincaré duality spaces*, *Topology* **7** (1968), 369–388. MR **40** #2089.
4. C. T. C. Wall, *Poincaré complexes. I*, *Ann. of Math. (2)* **86** (1967), 213–245. MR **36** #880.
5. E. C. Zeeman, *Seminar on combinatorial topology*. Chap. 7, *Engulfing*, Inst. Haute Études Sci., Paris, 1963; revised 1965 (mimeographed).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA,
PENNSYLVANIA 19104