

ON POLYNOMIAL APPROXIMATION IN $A_q(D)$

THOMAS A. METZGER

ABSTRACT. Let D be a bounded Jordan domain with rectifiable boundary and define $A_q(D)$, the Bers space, as the space of holomorphic functions f , such that

$$\iint_D |f| \lambda_D^{2-q} dx dy$$

is finite, where λ_D is the Poincaré metric for D . It is shown that the polynomials are dense in $A_q(D)$ for $q > 3/2$.

1. Introduction. Let D be a bounded Jordan domain. Define $A_q(D)$ ($q > 1$), the Bers space, as the Banach space of holomorphic functions $f(z)$, such that

$$(1.1) \quad \|f\|_q = \iint_D |f(z)| \lambda_D^{2-q}(z) dx dy < \infty,$$

when $\lambda_D(z)$ is the Poincaré metric for D . If ϕ is a Riemann mapping function from D onto U , the unit disk, and $\psi = \phi^{-1}$, then

$$(1.2) \quad \iint_D \lambda_D^{2-q}(z) dx dy = \iint_U |\psi'(\zeta)|^q (1 - |\zeta|^2)^{q-2} d\zeta d\bar{\zeta}.$$

Since for Jordan domains the rectifiability of the boundary is equivalent to $\psi' \in H^1(U)$, the Hardy class (cf. [3, p. 44]), it follows immediately by a theorem of Carleson [3, p. 157] that if the boundary of D is rectifiable then (1.2) is finite for all $q > 1$. Hence D bounded implies that the polynomials belong to $A_q(D)$, for all $q > 1$.

The question of polynomial density in $A_q(D)$ has been considered by various authors. In the case $q \geq 2$, Bers [2] and Knopp [4] proved that the polynomials were dense in $A_q(D)$ without any assumptions on the mapping functions ψ or ϕ . Later Sheingorn [6] proved that the polynomials are dense in $A_q(U^*)$, for all $q > 1$, where U^* is a particular Jordan region which appears in the proof of the "Main Lemma" of Knopp [4]. Finally Metzger

Received by the editors May 17, 1972 and, in revised form, June 15, 1972.

AMS (MOS) subject classifications (1970). Primary 30A82; Secondary 30A58, 41A10.

Key words and phrases. Polynomial density, Bers spaces.

© American Mathematical Society 1973

and Sheingorn [5] proved that if either $\psi' \in H^p$, for some $p > 1$, or D is a Smirnov domain, then the polynomials are dense in $A_q(D)$, for all $q > 1$.

In this paper we show

THEOREM 1. *If D is a bounded Jordan domain with a rectifiable boundary, then the polynomials are dense in $A_q(D)$, for $q > 3/2$.*

2. Proof of Theorem 1. Clearly we can assume that the origin lies in D and we choose $\psi(z) = 0$, $\psi'(0) > 0$. We now note

LEMMA 1. *Suppose D is a bounded Jordan domain with rectifiable boundary. The polynomials are dense in $A_q(D)$ if and only if ϕ'^q can be approximated by polynomials in $A_q(D)$.*

The necessity is clear since $\phi'^q \in A_q(D)$ for all $q > 1$ and the sufficiency follows easily from Lemma 1 of [5].

We define $\mathcal{H}_p(D) = \{f: f \text{ is holomorphic on } D \text{ and } \|f\|_p^* < \infty\}$, where

$$(2.1) \quad \|f\|_p^* = \left(\iint_D |f|^p dx dy \right)^{1/p}.$$

An application of Hölder's inequality yields

$$\begin{aligned} \|F\|_q &= \iint_U |F \circ \psi| |\psi'^q| (1 - |z|^2)^{q-2} dx dy \\ &\leq \left(\iint_U |F \circ \psi|^p |\psi'|^2 dx dy \right)^{1/p} \\ &\quad \cdot \left(\iint_U |\psi'|^{(qp-2)/(p-1)} (1 - |z|^2)^{(qp-2p)/(p-1)} dx dy \right)^{(p-1)/p} \\ &= \|F\|_p^* \left(\iint_D \lambda_D^{(qp-2)/(p-1)} dx dy \right)^{(p-1)/p} < \infty, \end{aligned}$$

if $(qp-2)/(p-1) > 1$, i.e. if $p > 1/(q-1)$, by (1.2) and the fact that $(qp-2)/(p-1) - 2 = (qp-2p)/(p-1)$. Thus polynomial approximation in $\mathcal{H}_p(D)$ implies polynomial approximation in $A_q(D)$ and, by Lemma 1, it follows that we need only show that $\phi' \in \mathcal{H}_p(D)$ for $p > q/(q-1)$.

To see that this holds, we first note that $q < 2$ implies $p > 2$ and by changing variables we get

$$\iint_D |\phi'|^p dx dy = \iint_U |\psi'|^{2-p} dx dy = I.$$

Since ψ is a bounded schlicht function with $\psi(0)=0$, it follows that there exists an $M (>0)$ such that $|\psi'(z)| \geq M(1-|z|^2)$ for all z in U . Hence I is finite if $p < 3$, i.e., if $q > 3/2$ and this completes the proof.

REMARK. If one lets G be a discrete group of conformal transformations on D and defines $A_q(D, G)$ as in Bers [1] then Theorem 2 of that paper yields the density of the Poincaré theta series of the polynomials in $A_q(D, G)$. (Cf. [1] and [5] for a precise formulation of the result plus all definitions and notations.)

BIBLIOGRAPHY

1. L. Bers, *Automorphic forms and Poincaré series for infinitely generated Fuchsian groups*, Amer. J. Math. **87** (1965), 196–214. MR **30** #4837.
2. ———, *A non-standard integral equation with applications to quasiconformal mappings*, Acta Math. **116** (1966), 113–134. MR **33** #273.
3. P. L. Duren, *Theory of H^p spaces*, Pure and Appl. Math., vol. 38, Academic Press, New York, 1970. MR **42** #3552.
4. M. I. Knopp, *A corona theorem for automorphic forms and related results*, Amer. J. Math. **91** (1969), 599–618. MR **40** #4450.
5. T. A. Metzger and M. Sheingorn, *Polynomial approximations in the Bers' spaces* (to appear).
6. M. Sheingorn, *Poincaré series of polynomials bounded away from zero on a fundamental region*, Amer. J. Math. (to appear).

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION,
TEXAS 77843