

## MULTIPLIERS FOR THE SPACE OF ALMOST-CONVERGENT FUNCTIONS ON A SEMIGROUP

CHING CHOU AND J. PETER DURAN

**ABSTRACT.** Let  $S$  be a countably infinite left amenable cancellative semigroup,  $FL(S)$  the space of left almost-convergent functions on  $S$ . The purpose of this paper is to show that the following two statements concerning a bounded real function  $f$  on  $S$  are equivalent: (i)  $f \cdot FL(S) \subset FL(S)$ ; (ii) there is a constant  $\alpha$  such that for each  $\varepsilon > 0$  there exists a set  $A \subset S$  satisfying (a)  $\varphi(\chi_A) = 0$  for each left invariant mean  $\varphi$  on  $S$  and (b)  $|f(x) - \alpha| < \varepsilon$  if  $x \in S \setminus A$ .

1. Let  $S$  be a semigroup,  $m(S)$  the space of bounded real functions on  $S$  with the sup norm.  $\varphi \in m(S)^*$  is called a left invariant mean on  $S$  if  $\|\varphi\| = 1$ ,  $\varphi \geq 0$  and  $\varphi(l_s f) = \varphi(f)$  for  $s \in S$  and  $f \in m(S)$ , where  $l_s f \in m(S)$  is defined by  $(l_s f)(t) = f(st)$ ,  $t \in S$ . The set of left invariant means on  $S$  is denoted by  $ML(S)$ . If  $ML(S)$  is nonempty, then  $S$  is said to be left amenable [2]. A bounded real function  $f$  on a left amenable semigroup is called left almost-convergent if  $\varphi(f)$  equals a fixed constant  $d(f)$  as  $\varphi$  runs through  $ML(S)$  [2]. The set of all left almost-convergent functions, denoted by  $FL(S)$ , is a vector subspace of  $m(S)$  and it contains constant functions. But, in general, it is not closed under multiplication. The purpose of this paper is to study this aspect of  $FL(S)$  and our main result is the following.

**THEOREM.** *Let  $S$  be a countable left-cancellative left amenable semigroup without finite left ideals. Then the following two statements concerning a function  $f \in m(S)$  are equivalent:*

- (i)  *$f$  is a multiplier of  $FL(S)$ , i.e.,  $f \cdot FL(S) \subset FL(S)$ ;*
- (ii)  *$f$  is  $S$ -convergent to a constant  $\alpha$ , i.e., for a given  $\varepsilon > 0$  there exists a set  $A \subset S$  such that*
  - (a)  *$\varphi(\chi_A) = 0$  for each  $\varphi \in ML(S)$ , and*
  - (b)  *$|f(x) - \alpha| < \varepsilon$  if  $x \in S \setminus A$ .*

2. From now on  $S$  will always denote a left-cancellative left amenable semigroup without finite left ideals.

---

Presented to the Society, January 17, 1972; received by the editors May 31, 1972.  
AMS (MOS) subject classifications (1970). Primary 43A07; Secondary 46N05.

*Key words and phrases.* Amenable semigroups, almost-convergence, multipliers, invariant means, weak Cauchy sequences.

© American Mathematical Society 1973

REMARKS. (1) A set  $A \subset S$  is said to be left almost-convergent if its characteristic function  $\chi_A$  is left almost-convergent. In this case we denote  $d(\chi_A)$  by  $d(A)$ . Roughly speaking, a set  $A \subset S$  is left almost-convergent if it is evenly distributed in  $S$  with respect to the semigroup structure and  $d(A)$  indicates the density of  $A \subset S$ . In particular,  $d(A)=0$  means that  $A$  is sparsely distributed in  $S$ . (The set  $A$  in the statement (ii) of the above Theorem is such a set.) For example, when  $S=N$ , the additive semigroup of positive integers, a set  $A \subset N$  is left almost-convergent if and only if

$$\lim_n (1/n) \text{Card}(\{k, k+1, \dots, k+n-1\} \cap A) = d(A)$$

exists uniformly in  $k$  [7].

(2) Since  $S$  contains no finite left ideals,  $d(B)=0$  for each finite subset  $B \subset S$  (cf. [5]). Therefore if  $f$  converges to  $\alpha$  at infinity, i.e., if given  $\varepsilon > 0$  there exists a finite set  $B \subset S$  such that  $|f(x) - \alpha| < \varepsilon$  whenever  $x \notin B$ , then  $f$   $S$ -converges to  $\alpha$ . On the other hand, the space of  $S$ -convergent functions is much smaller than  $FL(S)$ . Indeed,  $FL(S)$  separates points of  $\beta S$  [6], the Stone-Čech compactification of the discrete set  $S$ , while it is easy to see that  $f \in m(S)$  is  $S$ -convergent if and only if  $f$  is a constant on the set

$$K(S) = \text{cl}_{\beta S} \cup \{\text{supp } \varphi : \varphi \in ML(S)\},$$

(cf. [1]). Here we consider a bounded real function on  $S$  as a continuous function on  $\beta S$  and a mean on  $m(S)$  as a probability measure on  $\beta S$ . In particular, if  $\varphi \in ML(S)$ ,  $\text{supp } \varphi$  denotes the support of the measure  $\varphi$ .

(3) As in [2], let  $EG$  denote the smallest class of groups which contains all finite groups, all abelian groups and is closed under the following four ways of constructing new groups from given ones: (a) subgroup; (b) factor group; (c) group extension; and (d) direct limits.

Each group in  $EG$  is amenable and they constitute all the known amenable groups [2]. If we assume that  $S$  is an infinite group in  $EG$ , then a stronger result is known [1]:  $f \in m(S)$  is  $S$ -convergent if and only if

(a)  $f \cdot \chi_A \in FL(S)$  for each left almost-convergent set  $A$ ,

(b)  $f^n \in FL(S)$ ,  $n=1, 2, \dots$ ;

in particular, if  $A \subset S$  and  $A \cap B$  is left almost-convergent for each  $\chi_B \in FL(S)$ , then  $d(A)=1$  or  $0$ . It is not clear whether our Theorem yields the same conclusion. The proof of our Theorem is completely different from the proof in [1].

**3. Proof of the Theorem.** (ii) $\Rightarrow$ (i) is easy, cf. [1].

(i) $\Rightarrow$ (ii). Let  $f$  be a multiplier of  $FL(S)$ . To show that  $f$  is  $S$ -convergent, it suffices to show that  $f \equiv d(f)$  on  $\text{supp } \varphi$ . We claim that this follows from the following assertion:

( $\alpha$ ) If  $g \in FL(S)$  and  $\varphi \in ML(S)$ , then  $\varphi(fg) = \varphi(f)\varphi(g)$ .

Indeed, if  $(\alpha)$  holds and if  $\varphi \in ML(S)$  then

$$\varphi((f - d(f))^2) = (\varphi(f) - d(f))^2 = 0.$$

Therefore  $f \equiv d(f)$  on  $\text{supp } \varphi$  as we wanted.

PROOF OF  $(\alpha)$ . We shall consider  $l^1(S)^* = m(S)$  and  $l^1(S)^{**} = m(S)^*$ . If  $\psi \in l^1(S)$  and  $h \in m(S)$ , then  $h(\psi) = \psi(h) = \sum_{t \in S} h(t)\psi(t)$ . Since  $S$  is left amenable and countable there exists a sequence  $\varphi_n$  in  $l^1(S)$  such that  $\|\varphi_n\|_1 = 1$ ,  $\varphi_n \geq 0$ , and  $\lim_n(\varphi_n(h) - \varphi_n(l_x h)) = 0$  for each  $h \in m(S)$  and each  $x \in S$  [5, Lemma 5.1]. We shall need the following two well-known facts (cf. [3, §9]):

( $\beta$ ) If  $g \in FL(S)$ , then  $\lim_n \varphi_n(g) = d(g)$ .

( $\gamma$ )  $FL(S) =$  the closed linear span of  $\{l_x h - h : x \in S, h \in m(S)\} \cup \{\chi_S\}$ .

Let  $x \in S$  be fixed. Set  $\psi_n = \varphi_n \cdot f - l_x(\varphi_n \cdot f)$ , i.e.,  $\psi_n(t) = \varphi_n(t)f(t) - \varphi_n(xt)f(xt)$ ,  $t \in S$ . Then  $\psi_n \in l^1(S)$ . We claim that  $\psi_n$  is a weak Cauchy sequence in  $l^1(S)$ . Indeed, if  $h \in m(S)$ ,

$$\begin{aligned} \psi_n(l_x h) &= \sum_{t \in S} (\varphi_n(t)f(t)h(xt) - \varphi_n(xt)f(xt)h(xt)) \\ &= \sum_{t \in S} \varphi_n(t)f(t)(h(xt) - h(t)) + \sum_{t \in S \setminus xS} \varphi_n(t)f(t)h(t) \\ &= \varphi_n(f \cdot (l_x h - h)) + \varphi_n(fh \cdot \chi_{S \setminus xS}) \equiv a_n + b_n. \end{aligned}$$

Note that  $f \cdot (l_x h - h) \in FL(S)$ , since  $f$  is a multiplier of  $FL(S)$  and  $l_x h - h \in FL(S)$ . Therefore by ( $\beta$ )  $\lim_n a_n = d(f \cdot (l_x h - h))$ . Note also that  $1 \geq \varphi(\chi_{tS}) = \varphi(l_t \chi_{tS}) \geq \varphi(\chi_S) = 1$ , i.e.,  $\varphi(\chi_{S \setminus tS}) = 0$  for each  $t \in S$ . Therefore  $\chi_{S \setminus tS}$  is left almost-convergent to zero. By ( $\beta$ ) again, we get

$$|b_n| \leq \|f\|_\infty \|h\|_\infty \varphi_n(\chi_{S \setminus tS}) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore we have obtained:

( $\delta$ )  $\lim_n \psi_n(l_x h) = d(f \cdot (l_x h - h))$ ,  $h \in m(S)$ .

Since  $S$  is left-cancellative, each  $k \in m(S)$  is of the form  $l_x h$  for some  $h \in m(S)$ . Therefore  $\lim_n \psi_n(h)$  exists for each  $h \in m(S)$ , i.e.,  $\psi_n$  is a weak Cauchy sequence as we claimed. Since  $l^1(S)$  is weakly sequentially complete [4, p. 374], there exists  $\psi \in l^1(S)$  such that  $\psi = \lim_n \psi_n$  in the weak topology. Certainly,  $\psi(t) = \lim_n \psi_n(t)$  for  $t \in S$ . On the other hand,

$$\lim_n \varphi_n(t) = \lim_n \varphi_n(\chi_{tS}) = 0,$$

since  $\chi_{tS} \in FL(S)$  and  $d(\chi_{tS}) = 0$  (cf. Remark (1)). Hence

$$\psi(t) = \lim_n (\varphi_n(t)f(t) - \varphi_n(xt)f(xt)) = 0.$$

So,  $\psi \equiv 0$ . By ( $\delta$ ),  $d(f \cdot (l_x h - h)) = 0 = d(f) \cdot d(l_x h - h)$ . It is of course true

that  $d(f \cdot c\chi_S) = d(f) \cdot d(c\chi_S)$ . Hence  $(\alpha)$  follows from  $(\gamma)$  and the above observation.

## REFERENCES

1. C. Chou, *The multipliers for the space of almost convergent sequences*, Illinois J. Math. **16** (1972), 687–694.
2. M. M. Day, *Amenable semigroups*, Illinois J. Math. **1** (1957), 509–544. MR **19**, 1067.
3. ———, *Semigroups and amenability*, Proc. Sympos. Semigroups (Wayne State Univ., Detroit, Mich., 1968), Academic Press, New York, 1969, pp. 5–53. MR **42** #411.
3. N. Dunford and J. T. Schwartz, *Linear operators. I: General theory*, Pure and Appl. Math., vol. 7, Interscience, New York, 1958. MR **22** #8302.
5. E. Granirer, *On amenable semigroups with a finite-dimensional set of invariant means. I*, Illinois J. Math. **7** (1963), 32–48. MR **26** #1744.
6. S. P. Lloyd, *Subalgebras in a subspace of  $C(X)$* , Illinois J. Math. **14** (1970), 259–267. MR **41** #8995.
7. G. G. Lorentz, *A contribution to the theory of divergent sequences*, Acta Math. **80** (1948), 167–190. MR **10**, 367.

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK AT BUFFALO,  
AMHERST, NEW YORK 14226

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PUERTO RICO AT MAYAGUEZ,  
MAYAGUEZ, PUERTO RICO 00708