

PRIME GENERALIZED ALTERNATIVE RINGS I WITH NONTRIVIAL IDEMPOTENT

HARRY F. SMITH

ABSTRACT. A generalized alternative ring I is a nonassociative ring R in which the identities $(wx, y, z) + (w, x, [y, z]) - w(x, y, z) - (w, y, z)x$; $([w, x], y, z) + (w, x, yz) - y(w, x, z) - (w, x, y)z$; and (x, x, x) are identically zero. It is demonstrated here that if R is a ring of this type with characteristic different from two and three, then R semiprime with idempotent e implies that R has a Peirce decomposition relative to e . Furthermore, if R is prime and $e \neq 0, 1$; then R must be alternative.

1. Introduction. Let R be a nonassociative ring. As is customary, for $x, y, z \in R$ we denote by (x, y, z) the associator $(x, y, z) = (xy)z - x(yz)$ and by $[x, y]$ the commutator $[x, y] = xy - yx$. R is called power-associative if for every $x \in R$ the subring generated by x is associative.

In [2] Kleinfeld defines a generalized alternative ring I to be a nonassociative ring R such that for all $w, x, y, z \in R$ the following identities are satisfied:

- (1) $(wx, y, z) + (w, x, [y, z]) = w(x, y, z) + (w, y, z)x$,
- (2) $([w, x], y, z) + (w, x, yz) = y(w, x, z) + (w, x, y)z$,
- (3) $(x, x, x) = 0$.

It is easily verified that a ring of this type is power-associative.

Consider R to be a power-associative ring of characteristic different from two and define $x \circ y = \frac{1}{2}(xy + yx)$ for $x, y \in R$. If R contains an idempotent e , then Albert has shown in [1] that $R = R_1(e) + R_{1/2}(e) + R_0(e)$ where $R_i(e) = \{x \in R : x \circ e = ix\}$. In fact, $ex = x = xe$ for $x \in R_1(e)$ and $ex = 0 = xe$ for $x \in R_0(e)$. This decomposition of R is known as the Albert decomposition. Throughout this work we will denote $R_i(e)$ by just R_i .

Suppose now one also has $(R, e, e) = (e, R, e) = (e, e, R) = 0$. If, as in the associative case, one takes $x = exe + (ex - exe) + (xe - exe) + (x - ex - xe + exe)$, one sees that $R = R_{11} + R_{10} + R_{01} + R_{00}$ where $R_{ij} = \{x \in R : ex = ix, xe = jx\}$. This further decomposition of R is referred to as the Peirce decomposition.

Presented to the Society, September 1, 1971 under the title *On generalized alternative rings I and II*; received by the editors October 6, 1972.

AMS (MOS) subject classifications (1970). Primary 17A30.

Key words and phrases. Generalized alternative ring I , semiprime ring, Peirce decomposition, prime ring, alternative ring.

© American Mathematical Society 1973

A nonassociative ring R is said to be semiprime if R contains no nonzero ideal I such that $I^2=0$. If, in addition, R contains no nonzero ideals I and H such that $IH=0$, then R is said to be prime.

2. Preliminaries. Henceforth we assume R to be a generalized alternative ring I with characteristic different from two and three. Under the additional assumption that R contains an idempotent e , we will make use of the following results established by Kleinfeld in [2]:

(i) $(e, R, e)=0$.

(ii) In the Albert decomposition, $R_{1/2}R_i \subseteq R_{1/2}$ and $R_iR_{1/2} \subseteq R_{1/2}$ for $i=0, 1$. Furthermore, $B=\{x \in R_{1/2}: xR \subseteq R_{1/2}, Rx \subseteq R_{1/2}\}$ is an ideal of R such that $(e, e, R) \subseteq B$ and every element of B squares to zero.

(iii) If R permits a Peirce decomposition, then for $i, j, k, l=0$ or 1 , $R_{ij}R_{kl}=0$ when $j \neq k$ and $R_{ij}R_{jl} \subseteq R_{il}$, with two exceptions, namely $R_{01}R_{01} \subseteq R_{10}$ and $R_{10}R_{10} \subseteq R_{01}$. Furthermore, if $(R_{01}, R_{11}, R_{11})=(R_{11}, R_{11}, R_{10})=(R_{10}, R_{00}, R_{00})=(R_{00}, R_{00}, R_{01})=0$, then $I_i = \sum (R_{ii}, R_{ii}, R_{ii}) + \sum (R_{ii}, R_{ii}, R_{ii})R_{ii}$ for $i=0, 1$ are ideals of R .

The following identities will also be necessary for our calculations:

$$(4) \quad (wx, y, z) - (w, xy, z) + (w, x, yz) = w(x, y, z) + (w, x, y)z,$$

$$(5) \quad (w, xy, z) - (w, x, zy) + (w, x, y)z - (w, y, z)x = 0,$$

$$(6) \quad (w, xy, z) - (xw, y, z) + w(x, y, z) - y(w, x, z) = 0,$$

$$(7) \quad (x, x, yz) = y(x, x, z) + (x, x, y)z.$$

A straightforward verification shows that (4), known as the Teichmüller identity, holds for all w, x, y, z in any nonassociative ring. Identity (5) is obtained by subtracting (4) from (1), while (6) is obtained by subtracting (4) from (2). Identity (7) follows from letting $w=x$ in identity (2).

3. Semiprime and prime rings.

THEOREM 1. *Let R be a generalized alternative ring I with characteristic different from two and three. If R contains an idempotent e , then $I=(e, e, R)$ is an ideal of R such that $I^2=0$. Furthermore, $[I, R]=0$ and $R_{1/2}I=(R_{1/2})^2I=0$.*

PROOF. By (ii), $B=\{x \in R_{1/2}: xR \subseteq R_{1/2}, Rx \subseteq R_{1/2}\}$ is an ideal of R such that $(e, e, R) \subseteq B$ and every element of B squares to zero. Let $K=(e, e, B) \subseteq B$. We first show that K is an ideal of R such that $0=K^2=[K, R]=R_{1/2}K=(R_{1/2})^2K$.

We begin by making the following observations. Let $(e, y, z)=a_1+a_{1/2}+a_0$ and $(e, e, [y, z])=b_{1/2}$ where $a_i \in R_i$ for $i=0, \frac{1}{2}, 1$ and $b_{1/2} \in R_{1/2}$. Then (1) yields $(e, y, z) + (e, e, [y, z]) = e(e, y, z) + (e, y, z)e$ or $a_1 + a_{1/2} + a_0 + b_{1/2} = a_1 + ea_{1/2} + a_1 + a_{1/2}e$, whence $a_1 = a_0 = b_{1/2} = 0$. Since

from [1] we know $y, z \in R_{1/2}$ implies $y \circ z \in R_1 + R_0$, we also have $0 = (e, e, yz + zy) + (e, e, yz - zy) = 2(e, e, yz)$ or $(e, e, yz) = 0$ if $y, z \in R_{1/2}$.

Now if we let $w = x = z$ in (2) and then apply (3), we have $(z, z, yz) = (z, z, y)z$. If we then let $w = x = y$ in (5) and apply (3), we have $(y, y^2, z) = 2(y, y, z)y$. Taking $y = e$ and $z = x$, this gives $(e, e, x) = 2(e, e, x)e$. Let $b = (e, e, x)$. Then $b = 2be$ and, since $b = eb + be$, this implies $eb = \frac{1}{2}b = be$. In particular, we have $ek = \frac{1}{2}k = ke$ for every $k \in K$. Now taking $k, k' \in K \subseteq R_{1/2}$ and using our last observation above together with (7), we have $0 = (e, e, kk') = k(e, e, k') + (e, e, k)k' = \frac{1}{4}kk' + \frac{1}{4}kk' = \frac{1}{2}kk'$ or $K^2 = 0$.

We next show K to be an ideal of R . That K is additive is clear. Let $b \in B, k \in K$. Since $B \subseteq R_{1/2}$ implies $(e, e, B^2) = 0$, using (7) and the fact that $K^2 = 0$ we have $0 = (e, e, bk) = b(e, e, k) + (e, e, b)k = b(e, e, k) = \frac{1}{4}bk$ or $BK = 0$. Since $b, b' \in B$ implies $(b + b')^2 = 0$ or $bb' = -b'b$, it follows too that $KB = 0$. Let $x_i \in R_i$ for $i = 0, \frac{1}{2}, 1$. Noting that previous calculations show if $b' = (e, e, x_{1/2})$, then $(e, e, 4b') = b' = (e, e, x_{1/2})$, we now use (7), the fact that $BK = 0 = KB$, and the fact that B is an ideal of R to compute as follows:

$$\begin{aligned} (e, e, b)x_i &= (e, e, bx_i) - b(e, e, x_i) = (e, e, bx_i) \in K \quad \text{for } i = 0, 1; \\ (e, e, b)x_{1/2} &= (e, e, bx_{1/2}) - b(e, e, x_{1/2}) = (e, e, bx_{1/2}) - b(e, e, 4b') \\ &= (e, e, bx_{1/2}) \in K; \\ x_i(e, e, b) &= (e, e, x_i b) - (e, e, x_i)b = (e, e, x_i b) \in K \quad \text{for } i = 1, 1; \\ x_{1/2}(e, e, b) &= (e, e, x_{1/2} b) - (e, e, x_{1/2})b = (e, e, x_{1/2} b) - (e, e, 4b')b \\ &= (e, e, x_{1/2} b) \in K. \end{aligned}$$

Thus it follows K is an ideal of R .

Suppose now we are given $x \in R$. Let $x = x_1 + x_{1/2} + x_0$ where $x_i \in R_i$ for $i = 0, \frac{1}{2}, 1$. Then, using (7) while keeping in mind that $k = (e, e, 4k)$ for $k \in K$ and $0 = (e, e, [y, z])$ for $y, z \in R$, one has for $i = 0, 1$ that $x_i k = x_i(e, e, 4k) + (e, e, x_i)(4k) = 4(e, e, x_i k) = (4k)(e, e, x_i) + (e, e, 4k)x_i = kx_i$. Also keeping in mind that $0 = (e, e, yz)$ for $y, z \in R_{1/2}$ and $BK = 0 = KB$, one has $0 = (e, e, x_{1/2} k) = x_{1/2}(e, e, k) + (e, e, x_{1/2})k = x_{1/2}(e, e, k) = \frac{1}{4}x_{1/2} k$ as well as $0 = (e, e, kx_{1/2}) = k(e, e, x_{1/2}) + (e, e, k)x_{1/2} = \frac{1}{4}kx_{1/2}$. Thus $[K, R] = 0$ and, in particular, $R_{1/2}K = 0$.

Next let $x, y \in R_{1/2}$ and $k \in K$. Then (1) gives $(xy, e, k) + (x, y, [e, k]) = x(y, e, k) + (x, e, k)y$. But $[K, R] = 0$ implies $(x, y, [e, k]) = 0$, while K an ideal of R with $R_{1/2}K = 0$ implies $x(y, e, k) = 0 = (x, e, k)y$. Hence $(xy, e, k) = 0$. Let $xy = a_1 + a_{1/2} + a_0$ where $a_i \in R_i$ for $i = 0, \frac{1}{2}, 1$. Then $0 = (xy, e, k) = [(xy)e]k - \frac{1}{2}(xy)k = (a_1 + a_{1/2}e)k - \frac{1}{2}a_1 k - \frac{1}{2}a_0 k = a_1 k - \frac{1}{2}a_1 k - \frac{1}{2}a_0 k$, using the fact from (ii) that $R_{1/2}R_1 \subseteq R_{1/2}$. Thus $a_1 k = a_0 k$.

Now from our initial observations we have $(e, x, y) \in R_{1/2}$. Hence

$(e, x, y) = (ex)y - e(xy) = (ex)y - a_1 - ea_{1/2}$ gives $[(ex)y]_1 = a_1$ and $[(ex)y]_0 = 0$. Then $a_1 + a_{1/2} + a_0 = xy = (ex)y + (xe)y$ implies $[(xe)y]_1 = 0$. Thus $(ex)y \in R_1 + R_{1/2}$ while $(xe)y \in R_{1/2} + R_0$. Now since from (ii) we know $xe \in R_{1/2}$, our previous argument shows $[(xe)y]_1 k = [(xe)y]_0 k$. But $[(xe)y]_1 = 0$, so $[(xe)y]_1 k = 0 = [(xe)y]_0 k$. Hence $[(xe)y]k = [(xe)y]_1 k + [(xe)y]_{1/2} k + [(xe)y]_0 k = 0$, since $R_{1/2} K = 0$. In similar fashion we have $[(ex)y]k = 0$. But then $x \in R_{1/2}$ gives $(xy)k = [(xe + ex)y]k = [(xe)y]k + [(ex)y]k = 0$ or $(R_{1/2})^2 K = 0$.

Finally, it is clear that $K = (e, e, B) \subseteq (e, e, R)$. But we have also shown above that if $b = (e, e, x_{1/2})$, then $(e, e, 4b) = b = (e, e, x_{1/2})$. Thus $(e, e, R) \subseteq (e, e, 4B) = K$, that is $K = (e, e, R)$.

COROLLARY. *Let R be a generalized alternative ring I with characteristic different from two and three. If R contains an idempotent e , then R semiprime implies that R has a Peirce decomposition relative to e .*

PROOF. Linearization of (3) gives $(y, x, x) + (x, y, x) + (x, x, y) = 0$. By (i), one always has $(e, R, e) = 0$. Since, by Theorem 1, $I = (e, e, R)$ is an ideal of R such that $I^2 = 0$, R semiprime now implies in addition that $(e, e, R) = 0 = (R, e, e)$.

THEOREM 2. *Let R be a prime generalized alternative ring I with characteristic different from two and three. If R contains an idempotent $e \neq 0, 1$; then R is alternative.*

PROOF. Since R a prime ring implies that R is semiprime, by the above Corollary R has a Peirce decomposition relative to e . Throughout this proof we use the convention that $w_{ij}, x_{ij}, y_{ij}, z_{ij} \in R_{ij}$ for $i, j = 0$ or 1 . Then using the multiplication table described by (iii), (5) gives $(x_{11}, ey_{11}, z_{10}) - (x_{11}, e, z_{10}y_{11}) + (x_{11}, e, y_{11})z_{10} - (x_{11}, y_{11}, z_{10})e = 0$ and $(x_{00}, y_{00}e, z_{01}) - (x_{00}, y_{00}, z_{01}e) + (x_{00}, y_{00}, e)z_{01} - (x_{00}, e, z_{01})y_{00} = 0$, that is $(R_{11}, R_{11}, R_{10}) = 0 = (R_{00}, R_{00}, R_{01})$. Similarly (6) and (iii) yield $(x_{01}, y_{11}e, z_{11}) - (y_{11}x_{01}, e, z_{11}) + x_{01}(y_{11}, e, z_{11}) - e(x_{01}, y_{11}, z_{11}) = 0$ and $(x_{10}, ey_{00}, z_{00}) - (ex_{10}, y_{00}, z_{00}) + x_{10}(e, y_{00}, z_{00}) - y_{00}(x_{10}, e, z_{00}) = 0$, that is $(R_{01}, R_{11}, R_{11}) = 0 = (R_{10}, R_{00}, R_{00})$. Thus by (iii), $I_i = \sum (R_{ii}, R_{ii}, R_{ii}) + \sum (R_{ii}, R_{ii}, R_{ii})R_{ii}$ are ideals for $i = 0, 1$.

We now use a technique which was first used by Kleinfeld for prime associator-dependent rings. Consider $H = R_{10}R_{01} + R_{10} + R_{01} + R_{01}R_{10}$. Using the multiplication table described by (iii), it follows that to show H is an ideal of R it suffices to verify that $R_{10}R_{01}$ is an ideal of R_{11} and $R_{01}R_{10}$ is an ideal of R_{00} . But from (5) and (iii) we have $(x_{11}, y_{10}e, z_{01}) - (x_{11}, y_{10}, z_{01}e) + (x_{11}, y_{10}, e)z_{01} - (x_{11}, e, z_{01})y_{10} = 0$ or $(R_{11}, R_{10}, R_{01}) = 0$, while (6) and (iii) yield $(x_{10}, ey_{01}, z_{11}) - (ex_{10}, y_{01}, z_{11}) + x_{10}(e, y_{01}, z_{11}) - y_{01}(x_{10}, e, z_{11}) = 0$ or $(R_{10}, R_{01}, R_{11}) = 0$. Thus $R_{10}R_{01}$ is an ideal of R_{11} .

Analogously one has $(R_{00}, R_{01}, R_{10})=0=(R_{01}, R_{10}, R_{00})$ or $R_{01}R_{10}$ is an ideal of R_{00} .

Now (5) and (iii) give $(w_{11}, x_{10}y_{11}, z_{11})-(w_{11}, x_{10}, z_{11}y_{11})+(w_{11}, x_{10}, y_{11})z_{11}-(w_{11}, y_{11}, z_{11})x_{10}=0$ or $(R_{11}, R_{11}, R_{11})R_{10}=0$. Also, $[(R_{11}, R_{11}, R_{11})R_{11}]R_{10}\subseteq((R_{11}, R_{11}, R_{11}), R_{11}, R_{10})+(R_{11}, R_{11}, R_{11})(R_{11}R_{10})=0$, using (iii) and earlier calculation that $(R_{11}, R_{11}, R_{10})=0$. Hence $I_1R_{10}=0$. Then $I_1(R_{10}R_{01})\subseteq(I_1, R_{10}, R_{01})+(I_1R_{10})R_{01}=0$, since we have shown previously $(R_{11}, R_{10}, R_{01})=0$. From these last calculations and (iii), it now follows that $I_1H=0$. Similarly one may show $I_0H=0$.

Suppose $H=0$. Then $R_{10}=0=R_{01}$ implies R_{11} and R_{00} are ideals of R such that $R_{11}R_{00}=0$. Since R is a prime ring and $e \in R_{11}$, we must have $R_{00}=0$. But then $R=R_{11}$ implies $e=1$, contrary to assumption. Thus it must be the case $H \neq 0$. But $H \neq 0$ and R a prime ring imply $I_1=0=I_0$, or that R_{11} and R_{00} are associative subrings of R .

We may now join the calculations on p. 316 of [2] to conclude that for any $x_1, x_2, x_3 \in R$ one has $(x_1, x_2, x_3)=\text{sgn } \sigma(x_{\sigma 1}, x_{\sigma 2}, x_{\sigma 3})$ where σ is any permutation of three elements. Since this implies R is alternative, the proof is complete.

In conclusion, the author would like to express his appreciation to Professor Erwin Kleinfeld under whose direction this research was accomplished.

BIBLIOGRAPHY

1. A. A. Albert, *Power-associative rings*, Trans. Amer. Math. Soc. **64** (1948), 552–593. MR **10**, 349.
2. E. Kleinfeld, *Generalization of alternative rings*. I, J. Algebra **18** (1971), 304–325. MR **43** #308.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242