

AUTOMORPHISMS OF VON NEUMANN ALGEBRAS AS POINT TRANSFORMATIONS¹

CHARLES RADIN

ABSTRACT. Given a concrete separable C^* -algebra \mathfrak{A} with unit and a faithful normal finite trace τ on \mathfrak{A}'' , we introduce a notion of τ -almost every state on \mathfrak{A} which has the proper relationship with Segal's noncommutative integration theory. We then prove that any $*$ -automorphism of \mathfrak{A} is implemented by some point transformation in the state space of \mathfrak{A} , defined τ -almost everywhere. This generalizes the classical result of von Neumann-Maharam.

0. Notation and introduction. Throughout this paper, \mathfrak{A} will denote a norm separable C^* -algebra² with unit I of operators on a Hilbert space \mathcal{H} . \mathfrak{S} will denote the space of states on \mathfrak{A} , and τ a faithful normal finite trace on \mathfrak{A}'' .

In §1 we define the notion of τ -almost every state in \mathfrak{S} or sets of full τ -measure in \mathfrak{S} . In §2 it is shown that for each fixed element A in \mathfrak{A}'' we can define a set of full τ -measure, $S(A)$, in \mathfrak{S} and a natural limiting procedure to define $\rho(A)$ for all ρ in $S(A)$. The assigned values $\{\rho(A)|\rho \text{ in } S(A)\}$ are unique to the extent that any two such procedures must agree on a common subset of full τ -measure. When used in the obvious way to define τ -almost uniform convergence for sequences in \mathfrak{A}'' , it follows from Theorem B in the appendix³ that our definition coincides with that of Segal [3, Definition 2.3].

In §3 we generalize to noncommutative⁴ \mathfrak{A} a result of Maharam [1, Lemma 4]; namely we show that any $*$ -automorphism α of \mathfrak{A}'' (not necessarily leaving τ invariant) is induced by a transformation $\tilde{\alpha}$ of \mathfrak{S}

Received by the editors September 18, 1972.

AMS (MOS) subject classifications (1970). Primary 46L05, 46L10; Secondary 28A65.

Key words and phrases. Automorphisms of measure algebras, noncommutative integration.

¹ Research supported by AFOSR under Contract F44620-71-C-0108.

² For definitions and basic results in operator theory see, for example, [2].

³ Theorems A and B in the Appendix are slight modifications of proofs presented by E. Nelson in a course on functional analysis at Princeton University in the fall of 1971.

⁴ We are motivated by the abelian case with: $\mathcal{A} = L_2[0, 1]$; $\mathfrak{A} = C(X)$, the multiplication algebra of continuous functions on $[0, 1]$; $\mathfrak{A}'' = L_\infty[0, 1]$, the multiplication algebra of (equivalence classes of) essentially bounded measurable functions, and $\tau(\cdot) = \int \cdot dx$.

© American Mathematical Society 1973

defined τ -almost everywhere. $\tilde{\alpha}$ is unique to the extent that any two such transformations must agree on a common subset of full τ -measure.

1. **Definition of τ -almost every state.** A subset S of \mathfrak{S} will be said to consist of τ -almost every state in \mathfrak{S} , or to be a set of full τ -measure, if there exists a sequence $\{P_n\}$ of projections P_n in \mathfrak{A}'' such that:

- (a) $\{P_n\}$ is nondecreasing,
- (b) $\tau(I - P_n) \rightarrow_n 0$,
- (c) $S \supseteq S(\{P_n\}) \equiv \bigcup_n \overline{\text{convex hull } \{\omega_\psi \mid \psi \in \mathcal{H}, \|\psi\| = 1, P_n\psi = \psi\}^{w^*}}$,

where ω_ψ is the vector state on \mathfrak{A} generated by ψ . A sequence $\{P_n\}$ of projections P_n from \mathfrak{A}'' , satisfying (a) and (b), will be called an *exhaustion*.⁵ The fundamental property of the class of sets of full τ -measure, which follows immediately from [3, Theorem 6], is that it is closed under countable intersection.

2. **Isolation of discontinuities.** At this point we introduce the main technical tool. It is Tomita's [4] generalization to the noncommutative case of the following fact: Given f in $\mathcal{L}_\infty[0, 1]$, we can isolate its discontinuities, in that given $\varepsilon > 0$ there exists a set $E(\varepsilon)$ with Borel measure less than ε off which f coincides with some g from $C[0, 1]$. Therefore for any pure state ρ on $C[0, 1]$, corresponding to a point mass δ_x with x in $[0, 1] \setminus E(\varepsilon)$, or in fact for any state ρ "concentrated" on $[0, 1] \setminus E(\varepsilon)$, we can unambiguously define $\rho(f)$ as $\rho(g)$. And if we are dealing with f and $\chi_{E(\varepsilon)}$ (the characteristic function for $E(\varepsilon)$) only as elements of $L_\infty[0, 1]$, the assignment of $\{\rho(f) \mid \rho \text{ "concentrated" on } [0, 1] \setminus E(\varepsilon)\}$ would be ambiguous only up to measure zero.

In the general (not necessarily commutative) case, let A in \mathfrak{A}'' be given. From [4, Theorem 6], for each $n=1, 2, \dots$, there exists A_n in \mathfrak{A} and a projection P_n in \mathfrak{A}'' such that $\tau(I - P_n) < 1/2^n$ and $AP_n = A_nP_n$. Defining $R_n = \bigwedge_{i \geq n} P_i$, it follows from [2, 2.1.5 and 2.5.2] that $\tau(I - R_n) \rightarrow_n 0$ so that $\{R_n\}$ is an exhaustion, and $AR_n = A_nR_n$. Therefore, for $\rho \in S(\{R_n\})$ we have $\rho = w^* \lim_\beta \sum_{\psi \in \beta} \omega_\psi$ with $R_{n_\rho} \psi = \psi$ for some n_ρ (where the coefficients for a convex sum of ω_ψ 's are absorbed into the ψ 's, and " β " labels the convex sums), and we define $\rho(A) = \rho(A_{n_\rho})$. If through different choices in the procedure we had arrived at a different exhaustion $\{Q_n\}$, and thus an assignment of values $\{\rho(A) \mid \rho \in S(\{Q_n\})\}$, since $\{R_n \wedge Q_n\}$ is an exhaustion the two assignments would agree on the common subset $S(\{R_n \wedge Q_n\})$ of full τ -measure.

⁵ There is a more general definition for semifinite τ in the Appendix. The terminology is that of Nelson; see footnote 3.

In applications there may be a distinguished sequence $\{B_n\}$, with B_n in \mathfrak{A} , such that $B_n \rightarrow_n A$ in the strong operator topology. Then it is not difficult to prove using [3, Corollary 13.1] and Theorem B in the appendix that there exists a subset S of $S(\{R_n\})$, of full τ -measure, and a subsequence $\{B_{n_i}\}$ of $\{B_n\}$ such that for all ρ in S , $\lim_i \rho(B_{n_i})$ exists and equals $\rho(A)$ as we have already defined it.

3. Application to *-automorphisms of \mathfrak{A}'' . We now use the procedure of §2 for our stated objective. Let α be a *-automorphism of \mathfrak{A}'' and $\{A_i\}$ a countable dense subset of \mathfrak{A} . Let $\{P_n(i)\}$ be an exhaustion appropriate for $\alpha^{-1}(A_i)$, as discussed in §2. From [3, Theorem 6] there exists a subset S of $\bigcap_i S(\{P_n(i)\})$ of full τ -measure, and definitions of $\rho[\alpha^{-1}(A_i)]$ for all ρ in S and all i . For ρ in S we define $[\tilde{\alpha}\rho](A_i)$ as $\rho[\alpha^{-1}(A_i)]$. It is easy to see that $\tilde{\alpha}(\rho)$ can be uniquely extended by continuity to a state $\tilde{\alpha}(\rho)$ in \mathfrak{S} . The map $\tilde{\alpha}: S \rightarrow \mathfrak{S}$ is the one claimed in §0. It is clear that any two maps constructed in the above manner must agree on a common subset of full τ -measure. We summarize this result in the following

THEOREM. *Let \mathfrak{A} be a concrete, separable C^* -algebra with unit, and τ a faithful, normal, finite trace on \mathfrak{A}'' . Then any *-automorphism of \mathfrak{A}'' is implemented by some point transformation of the state space of \mathfrak{A} , defined τ -almost everywhere. The point transformation is unique to the extent that any two such transformations must coincide τ -almost everywhere.*

Appendix. Throughout this appendix M will denote a countably decomposable semifinite von Neumann algebra on a Hilbert space \mathcal{H} . We define an *exhaustion* as a sequence $\{P_n\}$ of projections P_n from M such that

- (a') $(I - P_n)$ is eventually finite with respect to M , and nonincreasing.
- (b') $(I - P_n) \rightarrow_n 0$ in the strong operator topology.

Note that if M possesses a faithful normal finite trace then this definition coincides with that of §1.

We say that a sequence $\{A_n\}$, with A_n in M , “converges nearly everywhere to zero in the sense of Segal” if given $\epsilon > 0$ there exists an exhaustion $\{P_n(\epsilon)\}$ such that $\|A_n P_n(\epsilon)\| < \epsilon$ for all n . (This is a trivial modification of Segal’s definition [2, Definition 2.3] which eliminates the significance of any finite number of terms of the sequence.)

THEOREM A (NELSON).³ *A sequence $\{A_n\}$, with A_n in M , converges nearly everywhere to zero in the sense of Segal if and only if there exists an exhaustion $\{P_n\}$ such that $\|A_n P_j\| \rightarrow_n 0$ for all j .*

PROOF. "If" direction. Let $\varepsilon > 0$ be given. For each n there exists a subsequence $\{P_{n_i}\}$ such that $\|A_n P_{n_i}\| < \varepsilon$ for all i . The result then follows from [3, Theorem 6].

"Only if" direction. Let $\varepsilon = 1/n$, and let $\{P_j\}$ be an exhaustion for the strongly dense domain⁶ given by the intersection over n of those defined by $\{P_j(1/n)\}$. Note from the proof of [3, Theorem 6] that for each j and k there exists $n(j; k)$ such that $P_j \leq P_{n(j; k)}(1/k)$. Therefore, if $n \geq n(j; k)$ we have

$$\|A_n P_j\| \leq \|A_n P_{n(j; k)}(1/k)\| \leq \|A_n P_n(1/k)\| < 1/k. \quad \text{Q.E.D.}$$

THEOREM B (NELSON).³ A sequence $\{A_n\}$, with A_n in M , converges nearly everywhere to zero in the sense of Segal if and only if there exists an exhaustion $\{P_j\}$ such that $\|P_j A_n P_j\| \rightarrow_n 0$ for all j .

PROOF. "Only if" direction. Obvious from Theorem A.

"If" direction. Let $\varepsilon > 0$ be given. Choose a faithful normal semifinite trace ϕ on M^+ such that $\phi(I - P_1) < \infty$. (This is easily seen to be possible, for example by using countable decomposability in the proof of [2, 2.5.3].) Then $\phi(I - P_j) < \infty$ for all j . By taking a subsequence if necessary we may assume that $\phi(I - P_j) < 1/2^j$. Define $\mathcal{H}_j = P_j \mathcal{H}$ and $L_j = \mathcal{H}_j \cap \bigcap_{k \geq j} A_k^{-1}(\mathcal{H}_k)$. It is not difficult to see that $\{L_j\}$ is a nondecreasing sequence of closed linear subspaces of \mathcal{H} , and that if we define Q_j as the projection onto L_j then Q_j is in M . Using [3, Lemma 3.1] and [2, 2.1.5] we see that

$$\phi(I - Q_j) \leq 1/2^j + \sum_{k \geq j} 2/2^k \rightarrow_j 0.$$

Therefore $\{Q_j\}$ is an exhaustion. Since $A_n Q_n = P_n A_n Q_n$ and $Q_n \leq P_n$ for all n , we have $\|A_n Q_n\| = \|P_n A_n Q_n\| \leq \|P_n A_n P_n\| < \varepsilon$ for all n . Q.E.D.

REFERENCES

1. D. Maharam, *Automorphisms of products of measure spaces*, Proc. Amer. Math. Soc. **9** (1958), 702-707. MR **20** #3963.
2. S. Sakai, *C*-algebras and W*-algebras*, Springer-Verlag, New York, 1971.
3. I. E. Segal, *A noncommutative extension of abstract integration*, Ann. of Math. (2) **57** (1953), 401-457; *ibid.* (2) **58** (1953), 595-596. MR **14**, 991; MR **15**, 204.
4. M. Tomita, *Spectral theory of operator algebras*. I, Math. J. Okayama Univ. **9** (1959), 63-98. MR **22** #8358.

JOSEPH HENRY LABORATORIES OF PHYSICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540

⁶ The notation is that of [3, Definition 2.1].