

ERRATA TO VOLUME 33

Franklin D. Tall, *A set-theoretic proposition implying the metrizable-ness of normal Moore spaces*, Proc. Amer. Math. Soc. **33** (1972), 195–198.

On page 196, line 21, insert \cup before $\{$. N. Smith has pointed out that Theorem 6 is only true for regular μ .

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO
M5S 1A1, CANADA

R. S. Gupta, *An extremal problem for functions with positive real part*, Proc. Amer. Math. Soc. **33** (1972), 455–462.

Page 461, lines 9 to 11, should read:

$$p(z) = \lambda \frac{1 + ze^{i\theta_1}}{1 - ze^{i\theta_1}} + (1 - \lambda) \frac{1 + ze^{i\theta_2}}{1 - ze^{i\theta_2}}$$

where $\lambda \in [0, 1]$ and $\theta_1, \theta_2 \in [0, 2\pi]$ are obtained from the equations

$$\lambda \frac{1 + re^{i\theta_1}}{1 - re^{i\theta_1}} + (1 - \lambda) \frac{1 + re^{i\theta_2}}{1 - re^{i\theta_2}} = Re^{i\phi}$$

and

$$\frac{1 + re^{i\theta_2}}{1 - re^{i\theta_2}} = a + e^{i(\phi - \alpha)} \left(\frac{1 + re^{-i\theta_1}}{1 - re^{-i\theta_1}} - a \right).$$

REFERENCE

1. R. S. Gupta, *An extremal problem for functions with positive real part*, Proc. Amer. Math. Soc. **33** (1972), 455–462.

74/114, DHANKUTTI, KANPUR 208001, INDIA

© American Mathematical Society 1974