

THE DISCRETENESS OF THE SPECTRUM OF SELF-ADJOINT, EVEN ORDER, ONE-TERM, DIFFERENTIAL OPERATORS

ROGER T. LEWIS

ABSTRACT. An open question which was asked by I. M. Glazman is answered. It is well known that the condition

$$\lim_{x \rightarrow \infty} x^{2n-1} \int_x^\infty r^{-1} = 0$$

is sufficient for the discreteness and boundedness from below of the spectrum of selfadjoint extensions of $(-1)^n (ry^{(n)})^{(n)}$. This paper shows that the condition is also necessary.

Let \tilde{L} denote any selfadjoint extension of

$$(-1)^n (ry^{(n)})^{(n)} \quad (x \geq 0, r(x) > 0).$$

THEOREM 1. *A necessary and sufficient condition that the spectrum of \tilde{L} be discrete and bounded below is that*

$$(1) \quad \lim_{x \rightarrow \infty} x^{2n-1} \int_x^\infty r^{-1} = 0.$$

The proof of the sufficiency part of the above theorem is due to V. A. Tkachenko and is exhibited in a book by I. M. Glazman [2, pp. 120, 121]. Glazman states that if $r(x)$ is monotonic, then relation (1) is necessary for the discreteness of the spectrum, but in the general case the problem of the necessity of condition (1) for $n > 1$ remains open. The development below completes the proof of Theorem 1.

Define

$$L_{2n}y = \sum_{k=0}^n (-1)^{n-k} (p_k y^{(n-k)})^{(n-k)}$$

where we assume that p_k is $n-k$ times continuously differentiable. The domain of L_{2n} is the set of all y such that y and $L_{2n}y$ are elements of $L^2[a, b]$ for a, b satisfying $0 < a < b < \infty$ and $y^{(k)}$ is absolutely continuous on compact subintervals of $(0, \infty)$ for $k=0, 1, \dots, 2n-1$. Let \mathcal{D} denote

Presented to the Society, November 16, 1973; received by the editors April 9, 1973 and, in revised form, June 12, 1973.

AMS (MOS) subject classifications (1970). Primary 34B25.

the set of all y in the domain of L_{2n} such that y has compact support interior to $[a, \infty)$ for some $a > 0$. Let T denote the closure of the restriction of L_{2n} to \mathcal{D} .

The differentiable operator L_{2n} is said to be *oscillatory on $[a, \infty)$* if for any given $b > a$ there are numbers c and d and a function $y \neq 0$ such that $L_{2n}(y) = 0$, $b \leq c < d$, and

$$y^{(k)}(c) = 0 = y^{(k)}(d) \quad \text{for } k = 0, 1, \dots, n - 1.$$

Otherwise, L_{2n} is said to be *nonoscillatory on $[a, \infty)$* .

THEOREM 2 (GLAZMAN [2]). *The following statements are equivalent.*

(i) *The spectrum of every selfadjoint extension of T is bounded below and discrete.*

(ii) *For every real number λ , $L_{2n} - \lambda$ is nonoscillatory on $[0, \infty)$.*

THEOREM 3 (AHLBRANDT [1]). *Suppose that r and p are positive, real-valued functions which are Lebesgue integrable on arbitrary compact subintervals of $[0, \infty)$. Then $(-1)^n(r^{-1}y^{(n)})^{(n)} - py$ is nonoscillatory on $[a, \infty)$ for some $a > 0$ if and only if $(-1)^n(p^{-1}y^{(n)})^{(n)} - ry$ is nonoscillatory on $[a, \infty)$.*

The following theorem is an extension by the author [3] of a theorem due to Glazman [2].

THEOREM 4. *If $p(x) \leq 0$, $0 < r(x) \leq Mx^\alpha$ for some $\alpha < 2n - 1$, and*

$$\limsup_{x \rightarrow \infty} x^{2n-1-\alpha} \int_x^\infty |p| > M \cdot A_n^2$$

where

$$A_n^{-1} = \frac{(2n - 1)^{1/2}}{(n - 1)!} \sum_{k=1}^n (-1)^{k-1} \binom{n - 1}{k - 1} / (2n - k)$$

then $(-1)^n(ry^{(n)})^{(n)} + py$ is oscillatory on $[a, \infty)$ for $a > 0$.

PROOF OF THEOREM 1. (Necessity) Suppose that

$$\lim_{x \rightarrow \infty} x^{2n-1} \int_x^\infty r^{-1} \neq 0;$$

then

$$\limsup_{x \rightarrow \infty} x^{2n-1} \int_x^\infty r^{-1} = \beta > 0.$$

There is a constant $c > 0$ such that $\lambda \geq c$ implies that $\beta > A_n^2/\lambda$. Hence, by Theorem 4,¹ $(-1)^n(\lambda^{-1}y^{(n)})^{(n)} - r^{-1}y$ is oscillatory on $[a, \infty)$ for $a > 0$ and

¹ Theorem 11/31 (9)/ , p. 100 of Glazman [2] could also be used if one lets $q = -\lambda r^{-1}$.

$\lambda \geq c$ which implies that $(-1)^n (ry^{(n)})^{(n)} - \lambda y$ is oscillatory on $[a, \infty)$ by Theorem 3. Consequently, by Theorem 2, the spectrum of \tilde{L} is not necessarily discrete and bounded below.

REFERENCES

1. Calvin D. Ahlbrandt, *Equivalent boundary value problems for self-adjoint differential systems*, J. Differential Equations **9** (1971), 420–435. MR **44** #1860.
2. I. M. Glazman, *Direct methods of qualitative spectral analysis of singular differential operators*, Fizmatgiz, Moscow, 1963; English transl., Israel Program for Scientific Translations, Jerusalem, 1965 and Davey, New York, 1966. MR **32** #2938; #8210.
3. Roger T. Lewis, *Oscillation and nonoscillation criteria for some self-adjoint, even order, linear differential operators*, Pacific J. Math. (to appear).

DEPARTMENT OF MATHEMATICS, SLIPPERY ROCK STATE COLLEGE, SLIPPERY ROCK,
PENNSYLVANIA 16057