

MULTIPLIERS FOR l_1 -ALGEBRAS WITH APPROXIMATE IDENTITIES

CHARLES D. LAHR

ABSTRACT. Let S be a commutative semigroup with multiplier semigroup $\Omega(S)$. Assume that $l_1(S)$ is semisimple and possesses a bounded approximate identity. If $l_1(S)^0$ denotes the annihilator of $l_1(S)$ in $l_1(\Omega(S))$, then the multiplier algebra of $l_1(S)$ is topologically isomorphic to $l_1(\Omega(S))/l_1(S)^0$, and this quotient algebra of $l_1(\Omega(S))$ is itself an l_1 -algebra.

1. Introduction. Let S be a commutative semigroup with multiplier semigroup $\Omega(S)$. If $l_1(S)$ is semisimple and possesses an approximate identity of norm one, then it is proved in [4] that the multiplier algebra $\mathcal{M}(l_1(S))$ is isometrically isomorphic to $l_1(\Omega(S))$. The purpose of this paper is to describe $\mathcal{M}(l_1(S))$ when $l_1(S)$ possesses an approximate identity bounded by some number $R > 1$. The main result is contained in Theorem 4.2 where it is proved that in general $\mathcal{M}(l_1(S))$ is topologically a quotient algebra of $l_1(\Omega(S))$. Interestingly enough, it turns out that this quotient algebra is also an l_1 -algebra.

The paper is organized as follows: §2 is devoted to notation and background material; in §3 it is shown that it is always possible to imbed a certain subsemigroup of $\Omega(S)$ in the structure semigroup of $l_1(S)$; and finally, §4 contains the main result.

2. Preliminaries. Throughout this paper S is a commutative semigroup, and $(l_1(S), *)$ is semisimple. Unless otherwise stated, $l_1(S)$ possesses a bounded approximate identity of norm R . That is, there exists a net $\{E_d\} \subset l_1(S)$ such that $\|\alpha * E_d - \alpha\| \rightarrow 0$ for all $\alpha \in l_1(S)$ and $\|E_d\| \leq R$ for all d and some positive number R . If Γ denotes the structure semigroup associated with $(l_1(S), *)$ [6], then there is an isomorphism i_s of S onto a dense subsemigroup of Γ [3]. Moreover, the fact that $l_1(S)$ has a bounded approximate identity implies that S contains a set of relative units and Γ contains a finite set of relative units [4]. Let $U = \{\gamma_1, \gamma_2, \dots, \gamma_n\}$ be a minimal set of relative units for Γ , with $\gamma_i^2 = \gamma_i$ for all i [2]; thus, for each

Presented to the Society, January 25, 1973; received by the editors November 1, 1972.

AMS (MOS) subject classifications (1970). Primary 43A10, 43A20.

Key words and phrases. Convolution measure algebra, l_1 -algebra, multiplier.

© American Mathematical Society 1974

$\gamma \in \Gamma$ there exists j such that $\gamma\gamma_j = \gamma$. Further define $\hat{\Gamma}$ to be the set of all nonzero continuous semicharacters on Γ .

A bounded linear operator T from $l_1(S)$ into $l_1(S)$ is called a multiplier of $l_1(S)$ if $T(\alpha * \beta) = \alpha * T(\beta)$ for all $\alpha, \beta \in l_1(S)$. The set of multipliers of $l_1(S)$ is a commutative Banach algebra of operators under operator norm $\|\cdot\|$ and is denoted $\mathcal{M}(l_1(S))$ [7]. Taylor shows [6] that $l_1(S)$ is isometrically isomorphic to a subalgebra of $M(\Gamma)$ (norm denoted by $\|\cdot\|$), and it is proved in [4] that $\mathcal{M}(l_1(S))$ is also isomorphic to a subalgebra of $M(\Gamma)$ with $\|\cdot\|$ equivalent to $\|\cdot\|$ in this case.

3. Multipliers of S . A function $\sigma: S \rightarrow S$ having the property that $\sigma(xy) = x\sigma(y)$ for all $x, y \in S$ is called a multiplier of S . Let $\Omega(S)$ denote the set of all multipliers of S and consider S to be a subsemigroup of $\Omega(S)$. The natural isomorphism i_x of S into Γ can be extended to an isomorphism of $\Omega(S)$ into Γ if and only if Γ has an identity, that is, if and only if $l_1(S)$ has a weak bounded approximate identity of norm one [4]. In this section we show that, by utilizing the relative units which belong to S because $l_1(S)$ has an approximate identity, then it is possible to imbed in Γ a subsemigroup of $\Omega(S)$ containing S . The importance of this subsemigroup will be seen in the next section. We begin with a technical lemma.

LEMMA 3.1. *Assume that S is imbedded in Γ . For each $i, 1 \leq i \leq n$, there exist nets $\{x_{\rho(i)}\} \subset S$ and $\{u_{\rho(i)}\} \subset S$ such that*

- (a) $\lim x_{\rho(i)} = \gamma_i$, with $\gamma_i x_{\rho(i)} = x_{\rho(i)}$ for all $\rho(i)$,
- (b) $u_{\rho(i)} x_{\rho(i)} = x_{\rho(i)}$, $\gamma_i u_{\rho(i)} = u_{\rho(i)}$ for all $\rho(i)$.

PROOF. Let i be arbitrary, $1 \leq i \leq n$. Since S is dense in Γ , there exists a net $\{x_{\rho'(i)}\} \subset S$ such that $\lim x_{\rho'(i)} \rightarrow \gamma_i$. Then there exists a subnet $\{x_{\rho'(i)}\}$ such that $\gamma_i x_{\rho'(i)} = x_{\rho'(i)}$ for all $\rho'(i)$. To see that such a subnet exists observe that there exist j and a subnet $\{x_{d(i)}\}$ of $\{x_{\rho'(i)}\}$ such that $\gamma_j x_{d(i)} = x_{d(i)}$ for all $d(i)$. But $\gamma_i \gamma_j = \lim \gamma_j x_{d(i)} = \lim x_{d(i)} = \gamma_i$ implies that U is not a minimal set of relative units for Γ unless $j=i$.

Now, since $l_1(S)$ has an approximate identity, S has a set of relative units [4], and so there exists a net $\{u_{\rho'(i)}\} \subset S$ such that $u_{\rho'(i)} x_{\rho'(i)} = x_{\rho'(i)}$ for all $\rho'(i)$. Further, let $\{u_{\rho(i)}\}$ be a subnet of $\{u_{\rho'(i)}\}$ such that $\gamma_i u_{\rho(i)} = u_{\rho(i)}$ for all $\rho(i)$. To substantiate the existence of this subnet, once again observe that there exist j and a subnet $\{u_{d(i)}\}$ of $\{u_{\rho'(i)}\}$ such that $\gamma_j u_{d(i)} = u_{d(i)}$ for all $d(i)$. But

$$\begin{aligned} \gamma_j \gamma_i &= \lim \gamma_j x_{d(i)} = \lim \gamma_j u_{d(i)} x_{d(i)} \\ &= \lim u_{d(i)} x_{d(i)} = \lim x_{d(i)} = \gamma_i \end{aligned}$$

shows that U is not a minimal set of relative units for Γ unless $j=i$. Thus, the nets $\{x_{\rho(i)}\}$ and $\{u_{\rho(i)}\}$ have the properties stated in the lemma. This completes the proof.

It is possible to use the semicharacters of Γ to decompose Γ into the following subsemigroups: $H_\gamma = \{\nu \in \Gamma : \chi(\nu) = 0 \text{ if and only if } \chi(\gamma) = 0 \text{ for all } \chi \in \hat{\Gamma}\}$; $A_\gamma = \{\nu \in \Gamma : \text{if } \chi \in \hat{\Gamma} \text{ and } \chi(\gamma) \neq 0 \text{ then } \chi(\nu) \neq 0\}$. Also note that the characteristic function of A_γ is always a semicharacter of Γ [1]. In the next theorem we show that U determines a subset of $\Omega(S)$.

THEOREM 3.2. *Assume that S is imbedded in Γ and that nets $\{x_{\rho(i)}\} \subset S$, $\{u_{\rho(i)}\} \subset S$ are as described in Lemma 3.1, $i=1, 2, \dots, n$. Then for each $x \in S$ there exists $\rho(i)_x$ such that $\rho(i) \geq \rho(i)_x$ implies $xu_{\rho(i)} = x\gamma_i$, $i=1, 2, \dots, n$. Thus, for each i , $1 \leq i \leq n$, $S\gamma_i \subset S$ and hence multiplication by γ_i determines a multiplier of S .*

PROOF. Let $x \in S$ and i be arbitrary, $1 \leq i \leq n$. Since the characteristic function ϕ of $A_{x\gamma_i}$ belongs to $\hat{\Gamma}$ and $\lim xx_{\rho(i)} = x\gamma_i$, then $\phi(xx_{\rho(i)}) \rightarrow \phi(x\gamma_i) = 1$ implies that there exists $\rho(i)_x$ such that $\phi(x_{\rho(i)}) = \phi(x_{\rho(i)})\phi(x) = \phi(x_{\rho(i)}x) = 1$ for all $\rho(i) \geq \rho(i)_x$. Hence, by the nature of $\{u_{\rho(i)}\}$, $\phi(u_{\rho(i)}) = 1$ for all $\rho(i) \geq \rho(i)_x$. We thus have that $\rho(i) \geq \rho(i)_x$ implies that $x_{\rho(i)} \in A_{x\gamma_i}$, $u_{\rho(i)} \in A_{x\gamma_i}$, $xx_{\rho(i)} \in A_{x\gamma_i}$. Also note that because $\gamma_i u_{\rho(i)} = u_{\rho(i)}$ and $u_{\rho(i)} \in A_{x\gamma_i}$, it is easy to verify that $xu_{\rho(i)} \in H_{x\gamma_i}$ for all $\rho(i) \geq \rho(i)_x$. We assert that $xu_{\rho(i)} = x\gamma_i$ for all $\rho(i) \geq \rho(i)_x$. If this statement is false, then there exist $\rho(i)$ and $\chi \in \hat{\Gamma}$ such that $\chi(xu_{\rho(i)}) \neq \chi(x\gamma_i)$. Since $xu_{\rho(i)} \in H_{x\gamma_i}$, it follows that $\chi(xu_{\rho(i)}) \neq 0$ and $\chi(x\gamma_i) \neq 0$. Also, since $x_{\rho(i)} \in A_{x\gamma_i}$, then $\chi(x_{\rho(i)}) \neq 0$; and since $x_{\rho(i)}u_{\rho(i)} = x_{\rho(i)}$, then $\chi(u_{\rho(i)}) = 1$. Thus, $\gamma_i u_{\rho(i)} = u_{\rho(i)}$ implies that $\chi(\gamma_i) = 1$; but this leads to the contradiction that $\chi(x) = \chi(xu_{\rho(i)}) \neq \chi(x\gamma_i) = \chi(x)$. Therefore, the theorem is proved.

For each i , $1 \leq i \leq n$, define a function $\sigma_{\gamma_i}: S \rightarrow S$ by $\sigma_{\gamma_i}(x) = x\gamma_i$ for all $x \in S$, where the product is formed in Γ . Then $\sigma_{\gamma_i} \in \Omega(S)$ for all i . Further, define $D = \{\sigma_{\gamma_i} : \sigma \in \Omega(S), i=1, 2, \dots, n\}$; then D is a subsemigroup of $\Omega(S)$.

THEOREM 3.3. *There is an algebra isomorphism i_D of D into Γ which extends the natural isomorphism i_s of S into Γ .*

PROOF. Let $i_s(x) = \tilde{x}$ for all $x \in S$, and let $\sigma_{\gamma_i} \sigma \in D$, $\sigma \in \Omega(S)$. By Lemma 3.1 there exists a net $\{x_{\rho(i)}\} \subset S$ such that $x_{\rho(i)} \rightarrow \gamma_i$. Consider $\{\sigma(x_{\rho(i)})\} \subset \Gamma$ and in accordance with the compactness of Γ let $\{\sigma(x_{d(i)})\}$ be a convergent subnet with $\lim \sigma(x_{d(i)}) = \gamma_i^\sigma$, $\gamma_i^\sigma \in \Gamma$. Now, if $x \in S$, then $\tilde{x}\gamma_i^\sigma = \lim \tilde{x}\sigma(x_{d(i)}) = \lim \sigma(x) \tilde{x}_{d(i)} = \sigma(x) \tilde{x}\gamma_i$. Suppose that $\{x_\rho\} \subset i_s(S)$ is any net such that $\lim x_\rho = \gamma_i$, and further suppose that $\{\sigma(x_\rho)\}$ is a convergent subnet of $\{\sigma(x_\rho)\}$ with $\lim \sigma(x_\rho) = \gamma \in \Gamma$; then $\tilde{x}\gamma_i^\sigma = \tilde{x}\gamma$ for all $x \in S$ implies that $\chi(\gamma_i^\sigma) = \chi(\gamma)$ for all $\chi \in \hat{\Gamma}$ and hence $\gamma = \gamma_i^\sigma$. We may now define $i_D: D \rightarrow \Gamma$ by $i_D(\sigma_{\gamma_i} \sigma) = \gamma_i^\sigma$; i_D is one-to-one since $\gamma_i^\sigma = \gamma_i^{\sigma'}$ implies $\gamma_i \sigma(x) = \gamma_i \sigma'(x)$ for all $x \in S$ and hence $\sigma_{\gamma_i} \sigma = \sigma_{\gamma_i} \sigma'$. Thus, i_D is an isomorphism of D into Γ that extends i_s .

When Γ has an identity, $D = \Omega(S)$ and the isomorphism i_D is simply an imbedding of $\Omega(S)$ into Γ ; i_D then induces an isomorphism of $l_1(\Omega(S))$ into $M(\Gamma)$.

4. Multipliers of $l_1(S)$ induced by $l_1(\Omega(S))$. In this paragraph and in Proposition 4.1 we drop the restriction that $l_1(S)$ has an approximate identity. Since $S \subset \Omega(S)$, $l_1(S)$ can be viewed as a subalgebra of $l_1(\Omega(S))$. Let $l_1(S)^0$ denote the annihilator of $l_1(S)$ in $l_1(\Omega(S))$, that is $l_1(S)^0 = \{\tau \in l_1(\Omega(S)) : \tau * \alpha = 0 \text{ for all } \alpha \in l_1(S)\}$. It follows that $l_1(S)^0$ is a closed ideal in $l_1(\Omega(S))$, in which case $l_1(\Omega(S))/l_1(S)^0$ is a Banach algebra under quotient norm $\| \cdot \|_Q$. Now, there is a natural homomorphism $\tau \mapsto T_\tau$ of $l_1(\Omega(S))$ into $\mathcal{M}(l_1(S))$ given by $T_\tau(\alpha) = \tau * \alpha$ for all $\alpha \in l_1(S)$. However, in general this homomorphism cannot be expected to be one-to-one. But the following is true.

PROPOSITION 4.1. (1) *The natural map of $l_1(S)$ into $l_1(\Omega(S))/l_1(S)^0$ is one-to-one and $l_1(\Omega(S))/l_1(S)^0$ is semisimple.*

(2) *The induced homomorphism $\tau + l_1(S)^0 \mapsto T_\tau$ of $l_1(\Omega(S))/l_1(S)^0$ into $\mathcal{M}(l_1(S))$ is one-to-one and $\| \| T_\tau \| \| \leq \| \tau + l_1(S)^0 \|_Q$.*

PROOF. (1) If $\alpha + l_1(S)^0 = l_1(S)^0$ for some $\alpha \in l_1(S)$, then $\alpha * \delta_x = 0$ for all $x \in S$, in contradiction to the semisimplicity of $l_1(S)$ unless $\alpha = 0$.

Also if $\tau + l_1(S)^0 \neq l_1(S)^0$, then choose $\alpha \in l_1(S)$ such that $0 \neq \tau * \alpha \in l_1(S)$. Semisimplicity of $l_1(S)$ implies the existence of $\chi \in \hat{\Gamma}$ such that $(\tau * \alpha)^\wedge(\chi) \neq 0$, which in turn implies $\hat{\tau}(\chi) \neq 0$.

(2) To see that $\tau + l_1(S)^0 \mapsto T_\tau$ is a one-to-one map, observe that $T_\tau = 0$ implies that $\tau * \alpha = T_\tau(\alpha) = 0$ for all $\alpha \in l_1(S)$, in which case $\tau \in l_1(S)^0$.

If $\tau \in l_1(\Omega(S))$, $\tau' \in l_1(S)^0$, then for all $\alpha \in l_1(S)$ we have $\| \tau * \alpha \| = \| \tau * \alpha + \tau' * \alpha \| \leq \| \tau + \tau' \| \| \alpha \|$; thus, $\| \| T_\tau \| \| \leq \| \tau + l_1(S)^0 \|_Q$. This completes the proof.

If $l_1(S)$ has a bounded approximate identity that is not of norm one, nonzero annihilators of $l_1(S)$ are quite readily available. If \bar{e} denotes the identity of $\Omega(S)$ and

$$E = \sum_{1 \leq i \leq n} \delta_{\sigma_{\gamma_i}} - \sum_{1 \leq i < j \leq n} \delta_{\sigma_{\gamma_i \gamma_j}} + \cdots + (-1)^{n+1} \delta_{\sigma_{\gamma_1 \gamma_2 \cdots \gamma_n}},$$

then $(\delta_{\bar{e}} - E) \in l_1(S)^0$ and so, for every $\tau \in l_1(\Omega(S))$, $(\tau - \tau * E) \in l_1(S)^0$. On the other hand, if $l_1(S)$ has a bounded approximate identity of norm one, then because $\mathcal{M}(l_1(S))$ is isometrically isomorphic to $l_1(\Omega(S))$ [4], $l_1(S)^0 = 0$ according to Proposition 4.1.

Assume for the remainder of this paper that $l_1(S)$ has a bounded approximate identity and that E is defined as above. Consider

$$E * l_1(\Omega(S)) = \{E * \tau : \tau \in l_1(\Omega(S))\};$$

it is straightforward to show that $E * l_1(\Omega(S))$ is a closed subalgebra of $l_1(\Omega(S))$ and hence complete. Moreover, we are now in a position to prove the main result, namely that $\mathcal{M}(l_1(S))$ is topologically isomorphic to $l_1(\Omega(S))/l_1(S)^0$.

THEOREM 4.2. (1) $E * l_1(\Omega(S))$ is isometrically isomorphic to $l_1(D) \subset l_1(\Gamma)$.

(2) $l_1(\Omega(S))/l_1(S)^0$ is isomorphic to $E * l_1(\Omega(S))$, and hence $\|\cdot\|_Q$ is equivalent to $\|\cdot\|$. Moreover, $l_1(\Omega(S))/l_1(S)^0$ is an l_1 -algebra.

(3) $\mathcal{M}(l_1(S))$ is isomorphic to $l_1(\Omega(S))/l_1(S)^0$ and $\|\!\|\cdot\!\|$ is equivalent to $\|\cdot\|_Q$.

PROOF. Statements (1) and (2) are fairly obvious.

(1) Using the notation of Theorem 3.3, $i_D: D \rightarrow \Gamma$ is defined by $i_D(\sigma_i \sigma) = \gamma_i^\sigma$ for all $\sigma \in \Omega(S)$, $i=1, \dots, n$. The isomorphism i_D induces a mapping from $E * l_1(\Omega(S))$ into $l_1(\Gamma)$ by defining, for all $\sigma \in \Omega(S)$,

$$E * \delta_\sigma \mapsto \sum_{1 \leq i \leq n} \delta_{\gamma_i \sigma} - \sum_{1 \leq i < j \leq n} \delta_{\gamma_i \sigma \gamma_j \sigma} + \dots + (-1)^{n+1} \delta_{\gamma_1 \sigma \gamma_2 \sigma \dots \gamma_n \sigma}.$$

Identifying D with $i_D(D)$ this induced homomorphism is clearly an isometry of $E * l_1(\Omega(S))$ onto $l_1(D)$.

(2) Define a homomorphism from $l_1(\Omega(S))/l_1(S)^0$ to $E * l_1(\Omega(S))$ by $\tau + l_1(S)^0 \mapsto E * \tau$. This mapping is clearly one-to-one and onto. Thus, because $l_1(\Omega(S))/l_1(S)^0$ is a semisimple Banach algebra complete under $\|\cdot\|$ and $\|\cdot\|_Q$, then the two norms must be equivalent [5]. The fact that $l_1(\Omega(S))/l_1(S)^0$ is an l_1 -algebra follows from (1).

(3) By Theorem 3.1 of [4], $\mathcal{M}(l_1(S))$ is isomorphic to a subalgebra of $M(\Gamma)$; we now have that $l_1(\Omega(S))/l_1(S)^0$ is topologically isomorphic to $l_1(D)$, a closed subalgebra of $l_1(\Gamma)$, where D is a subsemigroup of Γ . If Λ denotes the uniformly closed subspace of $l_\infty(S)$ generated by \hat{S} , then the measure $\mu \in M(\Gamma)$ is a multiplier of $l_1(S)$ if and only if m_μ is continuous on Λ in the weak topology generated by $l_1(S)$, where, $m_\mu: \Lambda \rightarrow \Lambda$ is such that $m_\mu(f)$ is the Arens product of $\mu \in M(\Gamma)$ and $f \in \Lambda$ [4, Definition 3.3 and Corollary 3.8]. In order to prove that $\mathcal{M}(l_1(S))$ is isomorphic to $l_1(D)$, we need only show that if $\mu \in M(\Gamma)$ is such that m_μ is continuous on Λ , then μ determines a multiplier of $l_1(D)$ and hence belongs to $l_1(D)$ since $l_1(D)$ has an identity. Now, because D is a subset of Γ , Λ can be identified with a subset of $l_1(D)^*$ as in [4, Proposition 4.8: replace $\Omega(S)$ by D]. Likewise, Theorem 4.9 of [4] establishes the desired isomorphism between $\mathcal{M}(l_1(S))$ and $l_1(D)$ by replacing $\Omega(S)$ by D throughout that part of the proof. Theorem 3.1 of [4] and (2) then prove that $\|\!\|\cdot\!\|$ is equivalent to $\|\cdot\|_Q$.

EXAMPLE 4.3. Let S be the set of integers under maximum multiplication and consider the subgroup S_0 of $S \times S$ consisting of the negative axes $\{(x, 0), (0, y): x, y \in S, x \leq 0, y \leq 0\}$. Γ contains relative units γ_1 and γ_2 with products: $\gamma_1\gamma_2 = (0, 0)$; $\gamma_1(n, 0) = (n, 0)$ for all $n \leq 0$; $\gamma_1(0, m) = (0, 0)$ for all $m \leq 0$; $\gamma_2(0, m) = (0, m)$ for all $m \leq 0$; and $\gamma_2(n, 0) = (0, 0)$ for all $n \leq 0$. If $\sigma \in \Omega(S_0)$, then σ restricted to $X_1 = \{(x, 0): x \in S, x \leq 0\}$ is a multiplier of X_1 : suppose $x \in X_1$ is such that $\sigma(x) \notin X_1$; then $\sigma\sigma_{\gamma_1} = \sigma_{\gamma_1}\sigma$ implies that $\sigma(x) = (0, 0) \in X_1$ and hence a contradiction. Similarly, σ restricted to $X_2 = \{(0, x): x \in S, x \leq 0\}$ is a multiplier of X_2 . Thus, each $\sigma \in \Omega(S_0)$ is of the form (σ', σ'') where $\sigma' \in \Omega(X_1)$ and $\sigma'' \in \Omega(X_2)$ and

$$\begin{aligned}(\sigma', \sigma'')(x, 0) &= (\sigma'(x), 0), (\sigma', \sigma'')(0, x) \\ &= (0, \sigma''(x)) \quad \text{for all } x \in S, x \leq 0.\end{aligned}$$

Also, $(\sigma', \sigma'')(v', v'') = (\sigma'v', \sigma''v'')$ for all $\sigma', v' \in \Omega(X_1)$, $\sigma'', v'' \in \Omega(X_2)$. Now because $\Omega(X_i)$ is just X_i with an identity adjoined for $i=1, 2$, then

$$\begin{aligned}\Omega(S_0) &= \{(\sigma_{\gamma_1}, \sigma_{\gamma_2})\} \cup \{(\sigma_{\gamma_1}, \sigma_x): x \in S, x \leq 0\} \\ &\quad \cup \{(\sigma_x, \sigma_{\gamma_2}): x \in S, x \leq 0\} \\ &\quad \cup \{(\sigma_x, \sigma_y): x, y \in S, x \leq 0, y \leq 0\}.\end{aligned}$$

However, only the subsemigroup $\{(\sigma_{\gamma_1}, \sigma_0)\} \cup \{(\sigma_0, \sigma_{\gamma_2})\} \cup S_0$ is embedded in Γ . For a fuller discussion of S_0 and Γ see [3, Example 4.14].

REFERENCES

1. W. W. Comfort, *The Šilov boundary induced by a certain Banach algebra*, Trans. Amer. Math. Soc. **98** (1961), 501–517. MR **22** #11282.
2. E. Hewitt and H. S. Zuckerman, *The l_1 -algebra of a commutative semigroup*, Trans. Amer. Math. Soc. **83** (1956), 70–97. MR **18**, 465.
3. C. D. Lahr, *Approximate identities for convolution measure algebras*, Pacific J. Math. **47** (1973), 147–159.
4. ———, *Multipliers for certain convolution measure algebras*, Trans. Amer. Math. Soc. (to appear).
5. C. E. Rickart, *General theory of Banach algebras*, University Series in Higher Math., Van Nostrand, Princeton, N.J., 1960. MR **22** #5903.
6. J. L. Taylor, *The structure of convolution measure algebras*, Trans. Amer. Math. Soc. **119** (1965), 150–166. MR **32** #2932.
7. J. K. Wang, *Multipliers of commutative Banach algebras*, Pacific J. Math. **11** (1961), 1131–1149. MR **25** #1462.

TRAFFIC RESEARCH DEPARTMENT, BELL TELEPHONE LABORATORIES, INC., HOLMDEL, NEW JERSEY 07733

Current address: Department of Mathematics and Physics, Savannah State College, Savannah, Georgia 31404