

GENERALIZED CONTRACTIONS AND FIXED POINT THEOREMS

CHI SONG WONG¹

ABSTRACT. Let T be a self-mapping on a complete metric space (X, d) . Then T has a fixed point if there exist self-mappings $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5$ on $[0, \infty)$ such that (a) $\alpha_1(t) + \alpha_2(t) + \alpha_3(t) + \alpha_4(t) + \alpha_5(t) < t$ for $t > 0$, (b) each α_i is upper semicontinuous from the right, (c)

$$d(T(x), T(y)) \leq a_1d(x, T(x)) + a_2d(y, T(y)) + a_3d(x, T(y)) \\ + a_4d(y, T(x)) + a_5d(x, y)$$

for all pairs of distinct x, y in X , where $a_i = \alpha_i(d(x, y))/d(x, y)$. Related results are obtained for two mappings and mappings on a bounded convex subset of a uniformly convex Banach space.

0. Introduction. Let (X, d) be a (nonempty) complete metric space. Let T be a self-mapping on X . It is the purpose of this paper to obtain some fixed point theorems for certain T which are controlled by five functions α_i 's from $(0, \infty)$ to $[0, \infty)$: for any distinct x, y in X ,

$$d(T(x), T(y)) \leq a_1d(x, T(x)) + a_2d(y, T(y)) + a_3d(x, T(y)) \\ + a_4d(y, T(x)) + a_5d(x, y),$$

where $a_i = \alpha_i(d(x, y))/d(x, y)$. These control functions will satisfy certain regular and contractive conditions such as the upper semicontinuity from the right for each α_i and

$$\sum_{i=1}^5 \alpha_i(t) < t \quad \text{for } t > 0.$$

Most of the results can be modified for two self-mappings S, T on X which are controlled by five functions α_i 's in the above fashion (with

Received by the editors June 19, 1972 and, in revised form, June 30, 1972.

AMS (MOS) subject classifications (1970). Primary 47H10; Secondary 54H25.

Key words and phrases. Asymptotic center, commuting family, contraction, error control, uniform convex Banach space, upper semicontinuity, weakly compact convex set.

¹ This research was partially supported by the National Research Council of Canada Grant A8518 and a grant from the Canadian Mathematical Congress. The author is on leave of absence from Southern Illinois University, Carbondale, Illinois 62901 to the University of Windsor, Windsor, Ontario, Canada.

© American Mathematical Society 1974

$T(x)$ replaced by $S(x)$). Among others, our results generalize certain results of D. W. Boyd and J. S. W. Wong [1], Felix E. Browder ([2], [3]), G. Göhde [5]. When T is continuous and each α_i is increasing, we can characterize T topologically.

1. Self-mappings on a complete metric space.

THEOREM 1. *Let T be a self-mapping on a complete metric space (X, d) . Suppose that there exist functions α_i , $i=1, 2, 3, 4, 5$, of $(0, \infty)$ into $[0, \infty)$ such that*

- (a) *each α_i is upper semicontinuous from the right;*
- (b) *$\alpha_1(t) + \alpha_2(t) + \alpha_3(t) + \alpha_4(t) + \alpha_5(t) < t$, $t > 0$;*
- (c) *for any distinct x, y in X ,*

$$d(T(x), T(y)) \leq a_1 d(x, T(x)) + a_2 d(y, T(y)) + a_3 d(x, T(y)) \\ + a_4 d(y, T(x)) + a_5 d(x, y),$$

where $a_i = \alpha_i(d(x, y))/d(x, y)$.

Then T has a unique fixed point.

PROOF. Let x_0 be a point in X . Define

$$x_{n+1} = T(x_n), \quad b_n = d(x_n, x_{n+1}), \quad n = 0, 1, 2, \dots$$

We shall first prove that T has a fixed point. So we may assume that $b_n > 0$ for each n . By (c),

$$(1) \quad b_0 b_1 = b_0 d(T(x_1), T(x_0)) \\ \leq \alpha_1(b_0) b_1 + \alpha_2(b_0) b_0 + \alpha_4(b_0) d(x_0, x_2) + \alpha_5(b_0) b_0.$$

Since $d(x_0, x_2) \leq b_0 + b_1$, we have from (1)

$$(2) \quad b_1 \leq \frac{\alpha_2(b_0) + \alpha_4(b_0) + \alpha_5(b_0)}{b_0 - \alpha_1(b_0) - \alpha_4(b_0)} b_0.$$

Similarly,

$$(3) \quad b_2 \leq \frac{\alpha_1(b_1) + \alpha_3(b_1) + \alpha_5(b_1)}{b_1 - \alpha_2(b_1) - \alpha_3(b_1)} b_1.$$

By symmetry of x, y in (c), we may assume that $\alpha_1 = \alpha_2$ and $\alpha_3 = \alpha_4$. So from (2), (3) and induction,

$$(4) \quad b_{n+1} \leq \alpha(b_n), \quad n = 0, 1, 2, \dots,$$

where

$$\alpha(t) = \frac{\alpha_1(t) + \alpha_3(t) + \alpha_5(t)}{t - \alpha_2(t) - \alpha_4(t)} t, \quad t > 0.$$

From (b), $\alpha(t) < t$ for $t > 0$. So $\{b_n\}$ is decreasing and therefore converges to some point b in $[0, \infty)$. If $b > 0$, then

$$(5) \quad b = \lim_{n \rightarrow \infty} b_{n+1} \leq \limsup_{n \rightarrow \infty} \alpha(b_n).$$

From (a), α is upper semicontinuous from the right. So from (5), $b \leq \alpha(b)$, a contradiction. So $b = 0$. We shall prove that $\{x_n\}$ is Cauchy. Suppose not. Then there exist $r > 0$ and sequences $\{p(n)\}, \{q(n)\}$ such that for each $n = 0, 1, 2, \dots$,

$$(6) \quad p(n) > q(n) > n, \quad d(p(n), q(n)) \geq r,$$

and (by the well-ordering principle)

$$(7) \quad d(x_{p(n)-1}, x_{q(n)}) < r.$$

Let $n \geq 0, c_n = d(x_{p(n)}, x_{q(n)})$. Then

$$r \leq c_n \leq d(x_{p(n)-1}, x_{q(n)}) + d(x_{p(n)-1}, x_{p(n)}) \leq r + b_{p(n)-1}.$$

Since $\{b_n\}$ converges to 0, $\{c_n\}$ converges to r from the right. By (c),

$$\begin{aligned} c_n d(T(x_{p(n)}), T(x_{q(n)})) &\leq \alpha_1(c_n) b_{p(n)} + \alpha_2(c_n) b_{q(n)} \\ &\quad + \alpha_3(c_n) d(x_{p(n)}, x_{q(n)+1}) \\ &\quad + \alpha_4(c_n) d(x_{q(n)}, x_{p(n)+1}) + \alpha_5(c_n) c_n. \end{aligned}$$

So by letting $n \rightarrow \infty$, we obtain

$$r^2 \leq (\alpha_3(r) + \alpha_4(r) + \alpha_5(r))r,$$

a contradiction to (b). Hence $\{x_n\}$ is Cauchy. By completeness of (X, d) , $\{x_n\}$ converges to some point x in X . We shall prove that x is a fixed point of T . Since each $b_n > 0$, there is a subsequence $\{x_{k(n)}\}$ of $\{x_n\}$ such that $x_{k(n)} \neq x$ for each n . Let $n \geq 0, d_n = d(x, x_{k(n)})$. Then from (c),

$$\begin{aligned} d(x_{k(n)+1}, T(x)) &= d(T(x_{k(n)}), T(x)) \\ &\leq [\alpha_1(d_n) b_{k(n)} + \alpha_2(d_n) d(x, T(x)) + \alpha_3(d_n) d(x_{k(n)}, T(x)) \\ &\quad + \alpha_4(d_n) d(x, x_{k(n)+1}) + \alpha_5(d_n) d_n] / d_n. \end{aligned}$$

So

$$d(x, T(x)) \leq \frac{\alpha_2(d_n) + \alpha_3(d_n)}{d_n} d(x, T(x)) + o(n),$$

where $\{o(n)\}$ converges to 0. Since $\alpha_2(t) + \alpha_3(t) < t/2$ for $t > 0$,

$$d(x, T(x)) \leq d(x, T(x))/2.$$

So $T(x)=x$. If T has two distinct fixed points x, y in X , then

$$d(x, y) = d(T(x), T(y)) \leq (\alpha_3(d(x, y)) + \alpha_4(d(x, y)) + \alpha_5(d(x, y))) < d(x, y),$$

a contradiction. Hence T has a unique fixed point in X .

When $\alpha_1=\alpha_2=\alpha_3=\alpha_4=0$, Theorem 1 is reduced to a result of D. W. Boyd and J. S. W. Wong [1, Theorem 1]. We expand their proof by considering the functions $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha$ instead of a single mapping α .

THEOREM 2. *Let S, T be self-mappings on a complete metric space (X, d) . Suppose that there exist upper semicontinuous functions $\alpha_1=\alpha_2, \alpha_3=\alpha_4, \alpha_5$ of $(0, \infty)$ into $[0, \infty)$ such that*

- (a) $\alpha_1(t)+\alpha_2(t)+\alpha_3(t)+\alpha_4(t)+\alpha_5(t) < t, t > 0$;
- (b) for any distinct x, y in X ,

$$d(S(x), T(y)) \leq a_1d(x, S(x)) + a_2d(y, T(y)) + a_3d(x, T(y)) + a_4d(y, S(x)) + a_5d(x, y),$$

where $a_i = \alpha_i(d(x, y))/d(x, y)$.

Then either S or T has a fixed point.

OUTLINE OF A PROOF. Let x_0 be a point in X . Define

$$x_{2n+1} = T(x_{2n}), x_{2n+2} = S(x_{2n+1}), \quad n = 0, 1, 2, \dots$$

We may assume that $d(x_n, x_{n+1}) > 0$ for each n , otherwise some x_n is a fixed point of S or T . With the notation of Theorem 1, (1)–(7) still hold. However we cannot write $d(x_{p(n)+1}, x_{q(n)+1})$ as $d(S(x_{p(n)}), T(x_{q(n)}))$ or $d(S(x_{q(n)}), T(x_{p(n)}))$ unless $p(n)+q(n)$ is odd. When $p(n)+q(n)$ is even, we may estimate $d(x_{p(n)+1}, x_{q(n)+1})$ by

$$d(S(x_{q(n)}), T(x_{p(n)-1})) + d(x_{p(n)}, x_{p(n)+1})$$

or

$$d(S(x_{p(n)-1}), T(x_{q(n)})) + d(x_{p(n)}, x_{p(n)+1}).$$

But then $\{d(x_{p(n)-1}, x_{q(n)})\}$ converges to r from the left. So we assume that each α_i is upper semicontinuous.

There are simple examples of S, T which satisfy the conditions of Theorem 2, but S has two fixed points and T has none.

2. Approximations and errors.

THEOREM 3. *In Theorem 1, suppose further that each α_i is increasing. Then*

- (i) $d(T^n(x), x_0) \leq \alpha^n(d(x, x_0)), x \in X, n=0, 1, 2, \dots$, where x_0 is the

fixed point of T , $\alpha(0)=0$ and for $t>0$,

$$\alpha(t) = \frac{\alpha_1(t) + \alpha_2(t) + \alpha_3(t) + \alpha_4(t) + 2\alpha_5(t)}{2t - \alpha_1(t) - \alpha_2(t) - \alpha_3(t) - \alpha_4(t)} t;$$

(ii) α is increasing, continuous from the right and for any t in $[0, \infty)$, $\{\alpha^n(t)\}$ converges to 0. Hence $\{T^n(x)\}$ converges uniformly to the fixed point of T on any bounded subset of X .

PROOF. (i) Let x be a point in $X \setminus \{x_0\}$, $n \geq 0$, $b_n = d(T^n(x), x_0)$. By (c) in Theorem 1,

$$\begin{aligned} b_0 b_1 &\leq b_0 d(T(x), T(x_0)) \\ &\leq \alpha_1(b_0) d(x, T(x)) + \alpha_3(b_0) b_0 + \alpha_4(b_0) b_1 + \alpha_5(b_0) b_0 \\ &\leq \alpha_1(b_0)(b_0 + b_1) + \alpha_3(b_0) b_0 + \alpha_4(b_0) b_1 + \alpha_5(b_0) b_0. \end{aligned}$$

So

$$(8) \quad b_1 \leq \frac{\alpha_1(b_0) + \alpha_3(b_0) + \alpha_5(b_0)}{b_0 - \alpha_1(b_0) - \alpha_4(b_0)} b_0.$$

Similarly (interchange the roles of x, x_0),

$$(9) \quad b_2 \leq \frac{\alpha_2(b_1) + \alpha_4(b_1) + \alpha_5(b_1)}{b_1 - \alpha_2(b_1) - \alpha_3(b_1)} b_1.$$

Because of the symmetry of x, y , (c) in Theorem 1 still holds if we replace $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5$ respectively by

$$\frac{\alpha_1 + \alpha_2}{2}, \frac{\alpha_1 + \alpha_2}{2}, \frac{\alpha_3 + \alpha_4}{2}, \frac{\alpha_3 + \alpha_4}{2}, \alpha_5.$$

Thus from (8), (9) and induction, $b_{n+1} = \alpha(b_n)$, $n=0, 1, 2, \dots$. So by induction

$$d(T^n(x), x_0) = b_n \leq \alpha^n(b_0) = \alpha^n(d(x, x_0)), \quad n = 0, 1, 2, \dots$$

(ii) Each α_i is increasing and continuous from the right, so is α . Let $t > 0$. By (b) in Theorem 1, $\alpha(t) < t$. So $\{\alpha^n(t)\}$ is decreasing and therefore converges to some t_0 in $[0, \infty)$. $t_0 = 0$, otherwise, by the right continuity of α ,

$$t_0 = \lim_{n \rightarrow \infty} \alpha^{n+1}(t) \leq \alpha \left(\lim_{n \rightarrow \infty} \alpha^n(t) \right) = \alpha(t_0),$$

a contradiction to $\alpha(t) < t$ for $t > 0$.

When $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 0$ and X is bounded, Theorem 3 is reduced to a result of Felix E. Browder [3, Theorem 1] with $M = X$. As Browder

pointed out, it is the explicit control over the error term in the Picard theorem which contributes so much to the widespread usefulness. Theorem 3 gives a uniform error control sequence $\{\alpha^n(d_0)\}$ over the error sequence $\{d(T^n(x), x_0)\}$ of functions $d(T^n(x), x_0)$ on any bounded subset of X with a diameter d_0 .

3. Topological contractions. Let (X, \mathcal{T}) be a metrizable topological space. Suppose that \mathcal{T} is metrically topologically complete, i.e. \mathcal{T} admits a complete metric for X . A self-mapping T on X is a topological contraction if there is an admissible complete metric d for X such that T is a contraction with respect to d , i.e. there exists a number k in $[0, 1)$ such that $d(T(x), T(y)) \leq kd(x, y)$ for all x, y in X . Very often, a complete metric space (X, d) is given and then a topology \mathcal{T} for X is induced; is there a practical way of recognizing topological contractions on (X, \mathcal{T}) in the language of d ? The following result gives a partial answer to this question.

THEOREM 4. *In Theorem 3, suppose further that T is continuous. Then T is a topological contraction on X .*

PROOF. From Theorems 1 and 3, T has a unique fixed point x_0 and for any x in X , $\{T^n(x)\}$ converges to x_0 . By a result of P. R. Meyer [6, Theorem 1], we need only to find a neighborhood V of x_0 such that $\{T^n(V)\}$ converges to x_0 . Let V be the open ball in X with center at x_0 and radius 1. Then for any x in V , $d(T^n(x), x_0) \leq \alpha^n(1)$ for each x in V . So for any neighborhood U of x_0 , $T^n(V)$ is included in U for large n 's.

4. Self-mappings on a bounded closed convex subset of a uniformly convex Banach space. In Theorem 1, it can be proved that if X is compact and if " $<$ " in (b) is interchanged with " \leq " in (c), then T has a unique fixed point. However, compactness is too strong for many purposes. The following result is perhaps more interesting.

THEOREM 5. *Let K be a bounded closed convex subset of a uniformly convex Banach space B . Let S, T be continuous self-mappings on K such that either*

$$\inf\{d(S(x), x) : x \in K\} = 0 \quad \text{or} \quad \inf\{d(T(x), x) : x \in K\} = 0.$$

Suppose further that there exist functions α_i , $i=1, 2, \dots, 5$, of $K \times K$ into $[0, \infty)$ such that $\alpha_1 = \alpha_2$, $\alpha_3 = \alpha_4$, $\sum_{i=1}^5 \alpha_i \leq 1$ and for any distinct x, y in K ,

$$d(S(x), T(y)) \leq a_1 d(x, S(x)) + a_2 d(y, T(y)) + a_3 d(x, T(y)) \\ + a_4 d(y, T(x)) + a_5 d(x, y),$$

where $a_i = \alpha_i(x, y)$.

Then S or T has a fixed point.

PROOF. By symmetry, we may assume that

$$\inf\{d(S(x), x) : x \in K\} = 0.$$

So there is a sequence $\{x_n\}$ in K for which $\{d(x_n, S(x_n))\}$ converges to 0, where d is the metric for B induced by the norm $\| \cdot \|$ for B . Let

$$(10) \quad r_m(y) = \sup\{d(y, x_n) : n \geq m\}, \quad m = 0, 1, 2, \dots$$

Since K is uniformly convex, there exists a unique c_m in K such that

$$(11) \quad r_m(c_m) = \inf\{r_m(y) : y \in K\}.$$

By a recent result of M. Edelstein [4, Theorem 1], $\{c_m\}$ converges to some point c in K . c is called the asymptotic center of $\{x_n\}$. If $\{x_n\}$ converges to c , then by continuity of S , c is a fixed point of S . Suppose that $\{x_n\}$ does not converge to c . We shall prove that c is a fixed point of T . Let $n \geq m \geq 1$. Then with $a_i = \alpha_i(x_n, c_m)$, we have

$$d(S(x_n), T(c_m)) \leq a_1 d(x_n, S(x_n)) + a_2 d(c_m, T(c_m)) + a_3 d(x_n, T(c_m)) + a_4 d(c_m, S(x_n)) + a_5 d(x_n, c_m).$$

Since

$$\begin{aligned} d(S(x_n), T(c_m)) &\geq d(x_n, T(c_m)) - d(S(x_n), x_n), \\ d(c_m, S(x_n)) &\leq d(c_m, x_n) + d(x_n, S(x_n)), \\ d(c_m, T(c_m)) &\leq d(x_n, c_m) + d(x_n, T(c_m)), \end{aligned}$$

we have

$$d(x_n, T(c_m)) \leq (a_2 + a_4 + a_5)d(x_n, c_m) + (a_2 + a_3)d(x_n, T(c_m)) + (1 + a_1 + a_4)d(x_n, S(x_n)).$$

So

$$(12) \quad \begin{aligned} d(x_n, T(c_m)) &\leq \frac{a_2 + a_4 + a_5}{1 - a_2 - a_3} d(x_n, c_m) + \frac{1 + a_1 + a_4}{1 - a_2 - a_3} d(x_n, S(x_n)) \\ &\leq d(x_n, c_m) + 2d(x_n, S(x_n)) \leq r_m(c_m) + s_m, \end{aligned}$$

where $s_m = \sup\{2d(x_n, S(x_n)) : n \geq m\}$. Suppose to the contrary that $T(c) \neq c$. Then by continuity of T , there exists s in $(0, M]$ and a subsequence $\{c_{k(m)}\}$ of $\{c_m\}$ such that

$$d(c_{k(m)}, T(c_{k(m)})) \geq s \quad \text{for each } m,$$

where M is the diameter of K . Since B is uniformly convex,

$$\delta(\varepsilon) = \inf\{1 - \|x + y\|/2 : \|x\| \leq 1, \|y\| \leq 1, \|x - y\| \geq \varepsilon\}$$

is positive for each ε in $(0, 2]$. So by (10), (11) and (12),

$$\left\| x_n - \frac{c_{k(m)} + T(c_{k(m)})}{2} \right\| \leq (r_{k(m)}(c_{k(m)}) + s_{k(m)})(1 - \delta(s/M))$$

for each $n \geq k(m)$. Since $\{s_m\}$ converges to 0,

$$\sup \left\{ \left\| x_n - \frac{c_{k(m)} + T(c_{k(m)})}{2} \right\| : n \geq k(m) \right\} < r_{k(m)}(c_{k(m)})$$

for large m , a contradiction to the choice of $r_m(c_m)$'s.

We remark here that when $S=T$ and $\alpha_1=\alpha_2=0$, the condition $\inf\{d(T(x), x) : x \in K\}$ is automatically satisfied.

THEOREM 6. *Let K be a bounded closed convex subset of a uniformly convex Banach space B . Let \mathcal{F} be a commuting family of continuous self-mappings T on K which satisfy the conditions of Theorem 5 with $S=T$. Then the set of all common fixed points of \mathcal{F} is nonempty, closed and convex.*

PROOF. Let T be an element of \mathcal{F} and let F_T be the set of all fixed points of T . By Theorem 5, F_T is nonempty. By continuity of T , F_T is closed. Let x, y be distinct elements of F_T , let $t \in (0, 1)$ and let $z = (1-t)x + ty$. Then with the notation of Theorem 5 and with $a_i = \alpha_i(z, x)$,

$$\begin{aligned} d(T(z), x) &= d(T(z), T(x)) \\ &\leq a_1 d(z, T(z)) + (a_3 + a_5) d(z, x) + a_4 d(x, T(z)). \end{aligned}$$

Since $d(z, T(z)) \leq d(z, x) + d(x, T(z))$, we have

$$d(T(z), x) \leq \frac{a_1 + a_3 + a_5}{1 - a_1 - a_4} d(z, x) \leq d(z, x) = td(x, y).$$

Similarly, $d(T(z), y) \leq (1-t)d(x, y)$. Since B is strictly convex, $T(z)=z$. So F_T is convex. This proves the theorem for a single mapping. Consider the family $\mathcal{A} = \{F_f : f \in \mathcal{F}\}$. Obviously the intersection of the members of \mathcal{A} is closed and convex. For any f, g in \mathcal{F} , f, g commute and therefore $f(F_g)$ is included in F_g ; thus applying this theorem to f (restricted to F_g), we conclude that F_f intersects F_g . Repeating this argument, we know that \mathcal{A} has the finite intersection property. Since B is reflexive, every bounded closed convex subset of B is weakly compact. So \mathcal{A} has nonempty intersection.

If we restrict $\alpha_1=\alpha_2=\alpha_3=\alpha_4=0$ in Theorem 5, then the above theorem is reduced to the well-known result of Browder [2] and Göhde [5].

ADDED IN PROOF. For Theorem 6 and more references, one should see the paper by K. Goebel, W. A. Kirk and T. N. Shimi: *A fixed point theorem in uniformly convex spaces*. With a different approach, Theorem 5 is proved with $S=T$ and without the condition $\inf\{d(x, T(x)) : x \in K\} = 0$.

REFERENCES

1. D. W. Boyd and J. S. W. Wong, *On nonlinear contractions*, Proc. Amer. Math. Soc. **20** (1969), 458–464. MR **39** #916.
2. F. E. Browder, *Nonexpansive nonlinear operators in a Banach space*, Proc. Nat. Acad. Sci. U.S.A. **54** (1965), 1041–1044. MR **32** #4574.
3. ———, *On the convergence of successive approximation for nonlinear functional equations*, Nederl. Akad. Wetensch. Proc. Ser. A **71**=Indag. Math. **30** (1968), 27–35. MR **37** #5743.
4. M. Edelstein, *The construction of an asymptotic center with a fixed point property*, Bull. Amer. Math. Soc. **78** (1972), 206–208. MR **45** #1005.
5. D. Göhde, *Zum Prinzip der Kontraktiven Abbildung*, Math. Nachr. **30** (1965), 251–258. MR **32** #8129.
6. P. R. Meyer, *A converse to Banach's contraction theorem*, J. Res. Nat. Bur. Standards **71** B (1967), 73–76.

SUMMER RESEARCH INSTITUTE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA