

A NOTE ON A LEMMA OF ZARISKI AND HIGHER DERIVATIONS

WEI-EIHN KUAN

ABSTRACT. A sufficient condition is given for an α -adic complete ring R to be a power series ring over a subring.

1. Introduction. We prove in this note the following theorem: Let R be a ring and let α be an ideal in R such that $\bigcap_{i=0}^{\infty} \alpha^i = 0$ and R is complete with respect to the α -adic topology. Assume that there exists a higher derivation $D = \{D_i\}_{i=0}^{\infty}$ of R such that $D_1(x) = 1$ for some $x \in \alpha$. Let $E = D_0 - xD_1 + \cdots + (-1)^n x^n D_n + \cdots$. If $E(x) = 0$, then there exists a subring R_1 of R such that $R = R_1[[x]]$, and x is analytically independent over R_1 .

This result generalizes Zariski's original lemma [5, Lemma 4, p. 526], and [1, Theorem 6, p. 412], a version of Zariski's lemma when R is of positive characteristic, and also removes the condition that R is an integral domain as we mentioned at the end of [1, p. 414]. [5, Lemma 4, p. 256] played a very important role in the study of analytic product of an affine algebraic variety V along a given subvariety W of V in A. Seidenberg's paper on differential ideals [4].

In the last section, we generalize a lemma of M. Miyanishi [2, p. 194] slightly, and give some remarks on his proofs of his lemma and Proposition 1.3 [2, p. 194].

2. Preliminaries. Throughout this note, all rings are commutative with identity. A derivation D of a ring R is an additive group homomorphism from R to R such that $D(a \cdot b) = aDb + bDa$ for all a and b in R . A higher derivation $D = \{D_i\}_{i=0}^{\infty}$ of a ring R is a sequence of additive group homomorphisms from R to R such that

(1) $D_0 =$ identity map on R , and

(2) $D_n(a \cdot b) = \sum_{i=0}^n D_i(a) \cdot D_{n-i}(b)$ for all $n = 1, 2, \dots$, and for all a and b in R . (Note D_1 is always a derivation of R .) *Leibniz formula.*

Let R be a ring and let α be an ideal in R . Then R has a topological structure with $\{\alpha^i\}_{i=0}^{\infty}$ as the fundamental system of neighborhoods of the

Received by the editors February 26, 1973.

AMS (MOS) subject classifications (1970). Primary 13J10, 13J05, 13B10.

Key words and phrases. Higher derivations, derivations, α -adic ring, power series ring.

© American Mathematical Society 1974

zero in R . This is the so called α -adic topology in R . Both addition and multiplication are continuous in α -adic topology. R is a Hausdorff space if and only if $\bigcap_{i=0}^{\infty} \alpha^i = 0$. A mapping $f: R \rightarrow R$ is continuous if there exists a subsequence $\{\alpha^{v_n}\}_{n=0}^{\infty}$ of the sequence $\{\alpha^n\}_{n=0}^{\infty}$ such that $f(\alpha^{v_n}) \subset \alpha^n$, for each natural number n .

LEMMA 1. Let R be a ring. Let $D = \{D_i\}_{i=0}^{\infty}$ be a higher derivation of R . Then

(1) For $a_1, \dots, a_n \in R$, and for each natural number m ,

$$\begin{aligned} D_m(a_1 \cdots a_n) &= \sum_{i=1}^n a_1 \cdots \hat{a}_i \cdots a_n D_m(a_i) \\ &\quad + \sum_{i=1}^{m-1} D_i(a_1 \cdots a_{n-1}) \cdot D_{m-i}(a_n) \\ &\quad + \sum_{i=1}^{m-1} \sum_{j=2}^{n-1} a_{j+1} \cdots a_n D_i(a_1 \cdots a_{j-1}) D_{m-i}(a_j), \end{aligned}$$

where $a_1 \cdots \hat{a}_i \cdots a_n = a_1 \cdots a_{i-1} a_{i+1} \cdots a_n$.

(2) For each ideal α in R , and for each natural number m , $D_m(\alpha^n) \subset \alpha^{n-m}$ for each α^n , i.e. D_m is continuous with respect to the α -adic topology.

(3) For each ideal α in R and for each $x \in \alpha$, let $E_l = \sum_{i=0}^l (-1)^i x^i D_i$ for each natural number l . $E_l(\alpha^n) \subset \alpha^n$ for each α^n , i.e. E_l is a continuous additive group homomorphism of R with respect to the α -adic topology.

PROOF. Straightforward.

LEMMA 2. Let R be a ring. Let α be an ideal of R such that $\bigcap_{i=0}^{\infty} \alpha^i = 0$. Assume R is complete with respect to the α -adic topology. Let $x \in \alpha$, and let $E_l = \sum_{i=0}^l (-1)^i x^i D_i$. Then the sequence $\{E_l\}_{l=0}^{\infty}$ is uniformly convergent, and $E = \sum_{i=0}^{\infty} (-1)^i x^i D_i = \lim_{l \rightarrow \infty} E_l$ is a continuous endomorphism of R .

PROOF. The sequence $\{E_l(a)\}$ is a Cauchy sequence for each $a \in R$. In fact $E_i(a) - E_j(a) \in \alpha^n$ for $i, j > n$; i.e., for each given α^n , there exists a natural number $N (=n)$ such that $E_i(a) - E_j(a) \in \alpha^n$ for $i, j > N$. Hence $\{E_l(a)\}$ converges in R . Since the natural number N above is independent of $a \in R$, therefore $\{E_l\}_{l=0}^{\infty}$ converges uniformly in R . Hence $E = \lim_{l \rightarrow \infty} E_l$ is continuous. In fact for each α^n and for each $a \in \alpha^n$, there exists a natural number $N (=n)$ which is independent of a such that $E(a) = (E(a) - E_N(a)) + E_N(a) \in \alpha^n$. Thus $E(\alpha^n) \subset \alpha^n$.¹

Since $E(a+b) = \lim_{l \rightarrow \infty} E_l(a+b) = \lim_{l \rightarrow \infty} E_l(a) + \lim_{l \rightarrow \infty} E_l(b) = E(a) + E(b)$, therefore $E(a+b) = E(a) + E(b)$. Also for each natural number l ,

¹ I thank the referee for his remark that $E(\alpha^n) \subset \alpha^n$.

$E_i(a) \cdot E_i(b) \equiv E_i(ab) \pmod{x^{i+1}}$. Thus $E_i(a)E_i(b) - E_i(ab) \in \mathfrak{a}^{i+1}$. Therefore

$$\lim_{i \rightarrow \infty} E_i(a) \cdot E_i(b) = \lim_{i \rightarrow \infty} E_i(ab).$$

But $\lim_{i \rightarrow \infty} E_i(a) \cdot E_i(b) = \lim_{i \rightarrow \infty} E_i(a) \cdot \lim_{i \rightarrow \infty} E_i(b)$. So $E(ab) = E(a)E(b)$.

3. THEOREM 1. *Let R be a ring and let \mathfrak{a} be a proper ideal in R such that $\bigcap_{i=0}^{\infty} \mathfrak{a}^i = 0$, and R is complete with respect to the \mathfrak{a} -adic topology. Assume that there exists a higher derivation $D = \{D_i\}_{i=0}^{\infty}$ of R such that $D_1(x) = 1$ for some $x \in \mathfrak{a}$. Let $E = D_0 - xD_1 + \dots + (-1)^n x^n D_n + \dots$. If $E(x) = 0$, then there exists a subring R_1 of R such that $R = R_1[[x]]$, and x is analytically independent over R_1 ; i.e. if $\sum_{i=0}^{\infty} a_i x^i = 0$ where $a_i \in R_1$ then $a_i = 0$ for all $i = 1, 2, \dots$.*

PROOF. $E(x) = 0$ implies $E^{-1}(0) = Rx$. Indeed, if $a \in E^{-1}(0)$, then $E(a) = 0$; i.e. $a - xD_1(a) + \dots + (-1)^n x^n D_n(a) + \dots = 0$. Thus $a = xD_1(a) + \dots + (-1)^n x^n D_n(a) + \dots$ is in Rx . So $E^{-1}(0) \subset Rx$. The other inclusion is obvious. Next, we observe that $E^2 = E$. In fact for each $a \in R$,

$$E(a) = a - xD_1a + \dots$$

and

$$\begin{aligned} E^2(a) &= E(a) - E(xD_1a + \dots) \\ &= E(a) - E(x) \cdot E(D_1a - xD_2a + \dots) = E(a), \end{aligned}$$

so $E^2(a) = E(a)$ for all $a \in R$. Let $R_1 = E(R)$ then E is an identity map on R_1 . Let a be an arbitrary element in R . $E(a) = a - xD_1a + \dots + (-1)^n x^n D_na + \dots$ implies that $a = E(a) + a_1x$ for some $a_1 \in R$. Thus $a = E(a) + xE(a_1) + a_2x^2$ for some $a_2 \in R$, and so on. Therefore we have $a = E(a) + xE(a_1) + \dots + x^n E(a_n) + \dots$ for some $a_1, a_2, \dots, a_n, \dots$ in R , and $a \in R_1[[x]]$. Hence $R = R_1[[x]]$. Finally we suppose $b_0 + b_1x + \dots + b_nx^n + \dots = 0$ where $b_0, b_1, \dots, b_n, \dots$ are in R_1 . We prove inductively $b_0 = b_1 = \dots = b_n = \dots = 0$. Since E is identity on R_1 , $E(b_0 + b_1x + \dots + b_nx^n + \dots) = 0$ implies $b_0 = E(b_0) = 0$. Assume $b_0 = b_1 = \dots = b_i = 0$. We have $b_{i+1}x^{i+1} + b_{i+2}x^{i+2} + \dots = 0$. By Lemma 1 and $D_1(x) = 1$, we have $D_nx^n \equiv 1 \pmod{x}$ and $D_nx^{n+j} \equiv 0 \pmod{x}$ for all natural numbers n and j . Thus

$$\begin{aligned} 0 &= D_{i+1}(b_{i+1}x^{i+1} + b_{i+2}x^{i+2} + \dots) \\ &= D_{i+1}(b_{i+1}x^{i+1}) + D_{i+1}(b_{i+2}x^{i+2} + \dots) \\ &\equiv b_{i+1}D_{i+1}(x^{i+1}) \pmod{x} \equiv b_{i+1} \pmod{x}. \end{aligned}$$

Therefore $b_{i+1} + c_{i+1}x = 0$ for some $c_{i+1} \in R$. So $0 = E(b_{i+1} + c_{i+1}x) = b_{i+1}$. Therefore x is analytically independent over R_1 .

If R contains the field of rational numbers as a subring, then every derivation D of R gives rise to a higher derivation of R , namely, $\{D_0, D, D^2/2!, \dots, D^n/n!, \dots\}$ where D^n is the n th successive derivation of D , and D_0 is the identity map in R . Let \mathfrak{a} be an ideal of R such that R is a complete Hausdorff space with respect to the \mathfrak{a} -adic topology. Assume $Dx=1$ for some $x \in \mathfrak{a}$. Then the endomorphism $E=e^{-xD}=\sum (-1)^n x^n D^n/n!$ always maps x to zero. Let $R_1=E(R)$, then $D(R_1)=0$. Thus we have the following

COROLLARY 1. *Let R be a ring containing the field of rational numbers as a subring. Let \mathfrak{a} be an ideal in R such that R is a complete Hausdorff space with respect to the \mathfrak{a} -adic topology. Assume there exists a derivation D of R such that $Dx=1$ for some $x \in \mathfrak{a}$. Then there exists a subring R_1 of R such that (1) D is zero on R_1 and (2) $R=R_1[[x]]$ and x is analytically independent over R_1 .*

In the following, a semilocal (local) ring \mathfrak{D} is a Noetherian ring with finitely many (unique) maximal ideals. Let \mathfrak{m} be the intersection of the maximal ideals of \mathfrak{D} . It is well known that $\bigcap_{i=0}^{\infty} \mathfrak{m}^i=0$. In this case we use \mathfrak{m} -adic topology for R . As a corollary to Corollary 1, we have the original lemma of Zariski.

COROLLARY 2. *Let $(\mathfrak{D}, \mathfrak{m})$ be a complete semilocal ring of characteristic zero. Let D be a derivation of \mathfrak{D} . Assume that there exists an element x in \mathfrak{m} of \mathfrak{D} such that Dx is a unit in \mathfrak{D} . Then \mathfrak{D} contains a ring \mathfrak{D}_1 of representatives of the (complete) semilocal ring $\mathfrak{D}/\mathfrak{D}x$ having the following properties: (a) D is zero on \mathfrak{D}_1 ; (b) x is analytically independent on \mathfrak{D}_1 ; (c) \mathfrak{D} is the power series ring $\mathfrak{D}_1[[x]]$.*

PROOF. Replace D by $(1/Dx)D$ and apply Corollary 1.

We also get [1, Theorem 6, p. 412] as a corollary to the theorem,

COROLLARY 3. *Let $(\mathfrak{D}, \mathfrak{m})$ be a complete local ring. Let $x \in \mathfrak{m}$ and let $D=\{D_i\}_{i=0}^{\infty}$ be a higher derivation of \mathfrak{D} such that D_1x is a unit in \mathfrak{D} , and $D_i x=0$ for $i>1$. Then there exists a subring \mathfrak{D}_1 of \mathfrak{D} such that: (a) \mathfrak{D}_1 is a complete local ring, (b) x is analytically independent over \mathfrak{D}_1 , and (c) $\mathfrak{D}=\mathfrak{D}_1[[x]]$.*

PROOF. Let $D_1x=\varepsilon^{-1}$, where ε is a unit in \mathfrak{D} . Replacing $\{D_i\}_{i=0}^{\infty}$ by $\{\varepsilon^i D_i\}_{i=0}^{\infty}$, we may assume $D_1x=1$. Since $D_i x=0$ for $i>1$, therefore $E x=x-xD_1x=0$ where $E=D_0-xD_1+\dots+(-1)^n x^n D_n+\dots$. Thus the theorem is applicable.

REMARKS. (1) Theorem 1 and Corollary 1 hold under the assumption that D_1x is a unit. The proofs are easily modified.

(2) If R has \mathfrak{a} as its sole maximal ideal and is a complete Hausdorff space with respect to the \mathfrak{a} -adic topology, then $E(x)=0$ and $x \neq 0$ implies that $D_1(x)$ is a unit.

(3) If R is a complete Hausdorff integral domain with respect to the \mathfrak{a} -adic topology, then $E(x)=0$ and $x \neq 0$ implies that $D_1(x)$ is a unit.

4. Though the following theorem could be easily proved by a similar technique used in the proof of Theorem 1, we would like to prove it as a corollary to Corollary 1.

THEOREM 2. *Let R be a ring containing the field of rational numbers as a subring. Let \mathfrak{a} be an ideal in R such that $\bigcap_{i=0}^{\infty} \mathfrak{a}^i = 0$. Assume that there exists a derivation D of R such that (1) for each $\gamma \in R$, $D^i(\gamma) = 0$ for sufficiently large i , and (2) $D(x) = 1$ for some $x \in \mathfrak{a}$. Then (a) there exists a subring R_1 of R such that $R = R_1[x]$ and x is algebraically independent over R_1 ; (b) D is trivial on R_1 .*

PROOF. Let \hat{R} be the completion of R with respect to the \mathfrak{a} -adic topology. Then \hat{R} is a complete Hausdorff space with respect to the topology defined by the filtration $\hat{\mathfrak{a}} \supset (\mathfrak{a}^2)^\wedge \supset \dots \supset (\mathfrak{a}^n)^\wedge \supset \dots$, where $(\mathfrak{a}^i)^\wedge$ is the closure of \mathfrak{a}^i in \hat{R} . Thus $\bigcap_{i=0}^{\infty} (\mathfrak{a}^i)^\wedge = 0$. Note that $(\mathfrak{a}^i)^\wedge \supseteq \mathfrak{a}^i \hat{R}$ in general and equality holds if \mathfrak{a} is finitely generated. Let $\gamma_1, \gamma_2, \dots, \gamma_n, \dots$ be a Cauchy sequence in R ; Lemma 1 implies that $\{D^j(\gamma_i)\}_{i=0}^{\infty}$ is also a Cauchy sequence for each j . Define $\hat{D}^j(\gamma) = \lim_{i \rightarrow \infty} D^j(\gamma_i)$. Then it is easy to check that $\{\hat{D}_0, \hat{D}, \dots, \hat{D}^n/n!, \dots\}$ forms a higher derivation in \hat{R} . Moreover $\hat{D}^j(\mathfrak{a}^i)^\wedge \subset (\mathfrak{a}^{i-j})^\wedge$. Indeed let $\hat{\gamma} \in (\mathfrak{a}^i)^\wedge$ and let $\gamma_1, \dots, \gamma_n, \dots$ be a sequence in \mathfrak{a}^i such that $\lim_{n \rightarrow \infty} \gamma_n = \hat{\gamma}$. Then $\hat{D}^j(\hat{\gamma}) = \lim_{n \rightarrow \infty} D^j(\gamma_n)$. Since $D^j(\gamma_n) \in \mathfrak{a}^{i-j}$, therefore $\hat{D}^j(\hat{\gamma}) \in (\mathfrak{a}^{i-j})^\wedge$. Lemma 1 and Lemma 2 are easily verified. Since the kernel of the natural ring homomorphism from R to \hat{R} is $\bigcap_{i=0}^{\infty} \mathfrak{a}^i = 0$, R is viewed as a subring of \hat{R} . \hat{D}^i restricted to R is D^i so $\hat{D}x = 1$ and

$$\begin{aligned} \hat{E}(x) &= x - (x/1!) \hat{D}(x) + \dots + (-1)^n (x^n/n!) \hat{D}^n(x) + \dots \\ &= x - (x/1!) D(x) + (x^2/2!) D^2(x) + \dots + (-1)^n (x^n/n!) D^n(x) + \dots \\ &= 0. \end{aligned}$$

Thus it follows from Theorem 1 that $\hat{R} = \hat{R}_1[[x]]$, where \hat{R}_1 is a subring of \hat{R} and x is analytically independent over \hat{R}_1 . Let $R_1 = R \cap \hat{R}_1$. Then $R = R_1[x]$. Indeed, let $\gamma \in R$. Then $\gamma = a_0 + a_1x + \dots + a_nx^n + \dots$ where $a_i \in \hat{R}_1$ for all a_i . Since there exists a natural number N such that $D^N(\gamma) = 0$. Therefore

$$0 = \hat{D}^N(\gamma) = N! a_N + \frac{1}{2}(N + 1)! a_{N+1}x + \dots$$

Hence $a_i = 0$ for all $i \geq N$, and $\gamma = a_0 + a_1x + \dots + a_{N-1}x^{N-1}$. It follows that $R \subset \hat{R}_1[x]$. Applying D^{N-1} to γ , we have $D^{N-1}(\gamma) = (N-1)! a_{N-1}$.

Therefore $a_{N-1} \in R$. Applying D^{N-2} to $(y - a_{N-1}x^{N-1}) = a_0 + a_1x + \dots + a_{N-2}x^{N-2}$, we get $a_{N-2} \in R$ and so on. Consequently, a_0, \dots, a_{N-1} are all in $\hat{R}_1 \cap R = R_1$. So $R = R_1[x]$, and x is of course algebraically independent over R_1 . (b) follows from Corollary 1.

We would like to thank Professor M. Miyanishi for communicating to us the following result which we also observed independently.

PROPOSITION 1. *Let R be an integral domain of characteristic 0. Assume there is a derivation of R such that $D^i(a) = 0$ for each $a \in R$ and for sufficiently large i . Then $Dx = 0$ for all units x in R .*

PROOF. Let x be a unit in R , and let $y \in R$ be such that $xy = 1$. Then $x Dy + y Dx = 0$. Suppose $Dx \neq 0$. Thus $Dy \neq 0$. Let i be the natural number such that $D^i x = 0$ and $D^m x \neq 0$ for $m < i$, also let j be the natural number such that $D^j y = 0$ and $D^p y \neq 0$ for $p < j$. By Leibniz's formula,

$$0 = D^n(xy) = \sum_{k=0}^n \binom{n}{k} D^k(x) D^{n-k}(y).$$

Taking $n = i + j - 2$ and assuming $i \leq j$ we get $D^{i-1}(x) \cdot D^{i-1}(y) = 0$. Hence either $D^{i-1}(x) = 0$ or $D^{i-1}(y) = 0$, a contradiction.

Proposition 1 completes the proof of [2, Lemma 1.4, p. 194]. One could not use [2, Proposition 1.4, p. 194] to yield a proof to the last part of [2, Proposition 1.3, p. 193]. But Proposition 1 corrects that part of the proof.

[4, Theorem 5, p. 30] locates differential ideals; [2, Proposition 1.3, p. 193] becomes interesting, we give it a slight generalization.

THEOREM 3. *Let R be an integral domain of characteristic 0 with a unique maximal ideal \mathfrak{m} such that $\bigcap_{i=0}^{\infty} \mathfrak{m}^i = 0$, i.e. (R, \mathfrak{m}) is a local domain which may not be Noetherian. If there is a derivation D of R such that $D^i(a) = 0$ for each $a \in R$ and for sufficiently large i , then \mathfrak{m} is differential, i.e. $D(\mathfrak{m}) \subset \mathfrak{m}$, and \bar{D} induced by D on R/\mathfrak{m} is trivial.*

PROOF. Suppose $D\mathfrak{m} \not\subset \mathfrak{m}$. Then there is $x \in \mathfrak{m}$ such that $Dx = u^{-1}$, where u is a unit in R . Then $uDx = 1$. Replacing D by uD , we have $(uD)^i(a) = u^i D^i(a)$ by Proposition 1. Thus $(uD)^i(a) = 0$ for sufficiently large i . It follows from Theorem 2 that $R = R_1[x]$, a contradiction. The last part follows from Proposition 1.

Observing the fact that in a polynomial ring $A[x]$ the units in $A[X]$ are of the form $a_0 + a_1x + \dots + a_nx^n$ such that a_0 is a unit in A and a_2, \dots, a_n are nilpotent in A , we give two examples countering Proposition 1 when R is not an integral domain.

EXAMPLE 1. Let $R=Z/(4)[X]$, where Z is the domain of integers and X is an indeterminate over $Z/(4)$. Let D be a derivation of R such that $DX=1+2X+2X^2$. $(DX)^2=1$, $D^2X=2\neq 0$, $D^i(a)=0$ for each $a \in R$ and for large integers i .

EXAMPLE 2. Let $R=(Q[t])[X]$, where Q is the field of rational numbers, $t^2=0$ and X is an indeterminate over $Q[t]$. Let D be a derivation of R such that $DX=1+tX$. Then DX is a unit $((DX) \cdot (1-tX)=1)$ $D^2X=t\neq 0$, and $Dt=0$, $D^i(a)=0$ for each $a \in R$ and for large i .

In the setting of Theorem 2, when R is an integral domain, if there is a derivation D such that DX is a unit u for some $x \in \alpha$, then there is a $y \in \alpha$ ($y=x/u$) such that $Dy=1$. What can one say in a more general case? Both Example 1 and Example 2 give negative answers. Using an idea of Professor M. Rosenlicht [3, Theorem 1, p. 721] we prove the following theorem.

THEOREM 4. Let R be a ring, which contains the field of rational numbers, with an ideal α such that $\bigcap_{i=0}^{\infty} \alpha^i=0$ and R is complete with respect to the α -adic topology. Assume there is a derivation D of R such that Dx is a unit for some $x \in \alpha$. Then there exists an element $y \in \alpha$ such that $Dy=1$.

PROOF. If Dx is a unit then $D(x/Dx)-1 \in Rx \subset \alpha$. If we can construct a Cauchy sequence $\{x_0=x/Dx, x_1, \dots, x_n, \dots\}$ such that $x_i \in xR$ and $Dx_i-1 \in Rx^{i+1}$, then putting $y=\lim_{i \rightarrow \infty} x_i$, and since α is also closed, we have $y \in \alpha$ and $Dy=\lim_{i \rightarrow \infty} Dx_i=1$. The proposed construction goes inductively as follows: Since $D(Rx^{i+1}) \subset Rx^i$, D induces a surjective R -homomorphism $\bar{D}^{(i)}: Rx^{i+1} \rightarrow Rx^i/Rx^{i+1}$ such that

$$\bar{D}^{(i)}(x^{i+1}) = (i+1)x^i + Rx^{i+1}.$$

Therefore there exists $z_i \in Rx^{i+1}$ such that

$$Dz_i \equiv Dx_{i-1} - 1 \pmod{Rx^{i+1}}$$

for $i=1, 2, \dots$. Thus $D(x_{i-1}-z_i)-1 \in Rx^{i+1}$. Putting $x_i=x_{i-1}-z_i$, we have a sequence $\{x_0=x/Dx, x_1, x_2, \dots\}$ such that $Dx_i-1 \in Rx^{i+1} \subset \alpha^{i+1}$ for $i=0, 1, 2, \dots$. For a given α^n , there exists a positive integer N ($=n$) such that for $i, j > N$, $x_i-x_j \in \alpha^n$. Therefore $\{x_0, x_1, \dots\}$ is a Cauchy sequence as desired.

ADDED IN PROOF. The author recently discovered that Theorem 2 can be derived from Taylor's lemma, see Y. Nouaze and P. Gabriel's *Idéaux premiers de l'algèbre enveloppante d'une algèbre de Lie nilpotente*, J. Algebra 6 (1967), 77-99.

REFERENCES

1. W. C. Brown and W. Kuan, *Ideals and higher derivations in commutative rings*, *Canad. J. Math.* **24** (1972), 400–415. MR **45** #3388.
2. M. Miyanishi, *Some remarks on the actions of additive group schemes*. *J. Math. Kyoto Univ.* **10** (1970), 189–205. MR **43** #7437.
3. M. Rosenlicht, *Canonical forms for local derivations*, *Pacific J. Math.* **42** (1972), 721–732.
4. A. Seidenberg, *Differential ideals in rings of finitely generated type*, *Amer. J. Math.* **89** (1967), 22–42. MR **35** #2902.
5. O. Zariski, *Studies in equisingularity. I. Equivalent singularities of plane algebroid curves*, *Amer. J. Math.* **87** (1965), 507–536. MR **31** #2243.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING,
MICHIGAN 48823