

A FINITELY GENERATED RESIDUALLY FINITE GROUP WITH AN UNSOLVABLE WORD PROBLEM

STEPHEN MESKIN

ABSTRACT. The group described in the title is obtained as a quotient of a center-by-metabelian group constructed by P. Hall.

It is well known that a residually finite group which is finitely presented has a solvable word problem (V. H. Dyson [1], A. W. Mostowski [6]). It is also known (see the last sentence in V. H. Dyson [1]) although perhaps not well enough (see W. Magnus [5, p. 307]) that a residually finite group which is only finitely generated may have an unsolvable word problem. Herein we exhibit such a group which moreover is soluble, indeed it is center-by-metabelian. This is in some sense a minimal soluble example since all finitely generated metabelian groups have a solvable word problem. (This is an easy consequence of the fact that finitely generated free metabelian groups satisfy the maximal condition on normal subgroups (P. Hall [2]) and thus any finitely generated metabelian group can be almost finitely presented in the sense of Mostowski [6] and is residually finite (P. Hall [3]).)

Credit must be given to Peter Neumann who pointed out the appropriate example to be modified.

Our example uses a recursive function f with a nonrecursive range in a manner similar to the way Higman [5] exhibited a finitely presented group with an unsolvable word problem.

Let G be the center-by-metabelian group constructed by P. Hall in ([2, p. 434]). In the notation of P. Hall, G is generated by a and b and defined by

$$[b_i, b_j, b_k] = 1 \quad \text{where } b_i = a^{-i}ba^i, \quad i, j, k = 0, \pm 1, \dots,$$

$$c_{ij} = c_{i+k, j+k} \quad \text{where } c_{ij} = [b_j, b_i], \quad j > i, \quad i, j, k = 0, \pm 1, \dots$$

The center C of G is free abelian with basis

$$d_1, d_2, \dots \quad \text{where } d_r = c_{i, i+r}, \quad r = 1, 2, \dots, \quad i = 0, \pm 1, \dots$$

Received by the editors March 12, 1973.

AMS (MOS) subject classifications (1970). Primary 20E25, 20F10; Secondary 20E15.

Key words and phrases. Finitely generated, residually finite, unsolvable word problem, center-by-metabelian.

© American Mathematical Society 1974

Let N be the subgroup of G generated by $\{d_{f(n)}^{f(n)!} | n=0, 1, \dots\}$. Clearly N is normal in G and a presentation of G/N is obtained from the presentation of G above by adding the relations

$$d_{f(n)}^{f(n)!} = 1, \quad n = 0, 1, 2, \dots$$

Since f is recursive G/N is recursively presented. On the other hand one cannot decide for an arbitrary integer n whether or not $n \in \text{range } f$. So one cannot decide whether or not $d_n^{n!} = 1$ in G/N , i.e. G/N has an unsolvable word problem.

It only remains to show that G/N is residually finite. Since C/N is a direct sum of cycles it is clear that $N = \bigcap_{m=1}^{\infty} C^m N$ and so it suffices to show that for any m , $G/C^m N$ is residually finite. However m divides $r!$ for all but a finite number of integers r thus $C^m N/C^m$ is finite and so it suffices to show that G/C^m is residually finite.

To see that G/C^m is residually finite we let R_k be the normal subgroup of G generated by a^{2k+1} . Then $G/R_k C^m$ can be constructed in a manner analogous to the way P. Hall constructed G .

We define B to be the group generated by the elements $\beta_0, \beta_1, \dots, \beta_{2k}$ subject to the following defining relations.

(i)
$$[\beta_i, \beta_j, \beta_k] = 1, \quad i, j, k = 0, 1, \dots, 2k.$$

This makes B nilpotent of class 2 and B' free abelian with basis

$$\begin{aligned} \gamma_{ij} &= [\beta_j, \beta_i], \quad j - i \equiv 1, 2, \dots, k \pmod{2k + 1}; \\ \text{(ii)} \quad \gamma_{ij} &= \gamma_{st} \quad \text{whenever } j - i \equiv t - s \pmod{2k + 1}. \end{aligned}$$

Now B' has as a basis

$$\delta_r = \gamma_{i, i+r}, \quad r = 1, 2, \dots, k, \quad i = 0, 1, \dots, 2k$$

($i + r$ is taken mod $2k + 1$).

(iii)
$$\delta_r^m = 1, \quad r = 1, 2, \dots, k.$$

This gives B' exponent m .

Clearly the map $\beta_i \rightarrow \beta_{i+1}$ ($i+1$ is taken mod $2k+1$) is an automorphism of B of order $2k+1$. It is easy to see that $G/R_k C^m$ is the extension of B by this automorphism. It may be defined as the group generated by α and $\beta = \beta_0$ with the above defining relations and

$$\alpha^{2k+1} = 1, \quad \alpha^{-1} \beta_i \alpha = \beta_{i+1}, \quad i = 0, 1, \dots, 2k - 1.$$

Now it is clear for any number of reasons that $G/R_k C^m$ is residually finite—for example it is polycyclic. Furthermore if $g \notin C^m$ then g can be written as a product of a power of a and a finite number of β_i 's and

d_r 's. Thus there is a k large enough so that $g \notin R_k C^m$ and it follows that G/C^m is residually finite.

REFERENCES

1. V. H. Dyson, *The word problem and residually finite groups*, Notices Amer. Math. Soc. **11** (1964), 743, Abstract 616-7.
2. P. Hall, *Finiteness conditions for soluble groups*, Proc. London Math. Soc. (3) **4** (1954), 419–436. MR **17**, 344.
3. ———, *On the finiteness of certain soluble groups*, Proc. London Math. Soc. (3) **9** (1959), 595–622. MR **22** #1618.
4. G. Higman, *Subgroups of finitely presented groups*, Proc. Roy. Soc. Ser. A **262** (1961), 455–475. MR **24** #A152.
5. W. Magnus, *Residually finite groups*, Bull. Amer. Math. Soc. **75** (1969), 305–316. MR **39** #2865.
6. A. W. Mostowski, *On the decidability of some problems in special classes of groups*, Fund. Math. **59** (1966), 123–135. MR **37** #292.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT 06268