

ON A THEOREM OF A. PEŁCZYŃSKI

JOHN L. B. GAMLEN¹

ABSTRACT. If Y is a weakly complete Banach space, and X is a Banach space with separable dual, then every continuous linear operator from $C_X(K)$ to Y must be weakly compact. Here $C_X(K)$ denotes the space of continuous functions on the compact Hausdorff space K , having values in X .

In 1953, A. Grothendieck [5] proved that if Y is a weakly complete Banach space, then every continuous linear map from $C(K)$ to Y must be weakly compact. (Here K is an arbitrary compact Hausdorff space.) Later A. Pełczyński [7] weakened the assumption on Y , to the requirement that c_0 is not isomorphic to any subspace of Y . In 1962, Pełczyński [8, p. 645] obtained a result which implies that $C(K)$ may be replaced in the above by $C_X(K)$, the space of continuous X -valued functions on K , where X is a reflexive Banach space. (See also J. Batt and E. J. Berg [2, p. 237], where a different but related proof is given.) Necessary and sufficient conditions on X for this to still work are not known. However, in order for every continuous linear map $T: C_X(K) \rightarrow Y$ to be weakly compact for a given Y , it is obviously necessary that every continuous operator from X to Y be weakly compact. But if X has a separable dual space, this holds for weakly complete Y , since Cantor's diagonal argument allows us to extract from every bounded sequence in X a subsequence which is weakly Cauchy. This suggests our result below.

THEOREM. *If Y is a weakly complete Banach space, and X is a Banach space whose separable subspaces have separable duals, then every continuous linear operator from $C_X(K)$ to Y must be weakly compact.*

This does not quite include the result of Pełczyński because of our slightly stronger assumption that Y is weakly complete. The interesting thing is that in the above theorem we cannot replace weak completeness of Y by the assumption that Y has no subspace isomorphic to c_0 . The counterexample is the well-known space J of R. C. James [6], which has

Received by the editors March 26, 1973 and, in revised form, August 13, 1973.

AMS (MOS) subject classifications (1970). Primary 46E40; Secondary 46G10, 28A45, 46B15.

Key words and phrases. Weakly compact operators.

¹ The author is partially supported by NRC Grant A7552.

separable bidual, but is not reflexive. (The identity map on J fails to be weakly compact, even though J contains no copy of c_0 .)

Batt and Berg's proof of Pełczyński's result proceeds via weak compactness of the adjoint map. This involves a weak compactness theorem in the space of X' -valued measures, which will not work in our case, as it depends on reflexivity of X . However this approach was used in [1] to prove weak compactness of a map from a C^* -algebra to a Banach space containing no copy of c_0 . (The details are, needless to say, quite different.)

PROOF OF THE THEOREM. We first deal with the case that K and X (hence also X') are separable. By the representation theorem of [2, pp. 225–228], we may represent our map $T: C_X(K) \rightarrow Y$ as an integral with respect to a measure μ taking values in the space $[X, Y]$ of bounded operators between X, Y , and having semivariation absolutely continuous with respect to some positive regular measure λ on K . Thus the adjoint T' maps Y' into a space of measures with values in X' , all of which are absolutely continuous with respect to λ . For such measures the Radon-Nikodym theorem is well known to hold, so we may embed the range of T' in $L^1(X', \lambda)$, the space of λ -integrable, X' -valued functions, by the formula:

$$\langle f, T'y' \rangle = \langle Tf, y' \rangle = \int f d(y'\mu) = \int f(d(y'\mu)/d\lambda) d\lambda,$$

for $f \in C_X(K)$, $y' \in Y'$.

Associated with the element $y'\mu$ in $T'Y'$ is the function $d(y'\mu)/d\lambda$ in $L^1(X', \lambda)$, and it is easy to see that the embedding of $T'Y'$ into L^1 is norm increasing. Since $L^1(X', \lambda)$ is separable, we conclude that $T'Y'$ is a separable subspace of the dual of $C_X(K)$. This fact enables us to use Cantor's diagonal argument to extract a subsequence $\{f_{n_m}\}$ such that for every y' in Y' the sequence $\{\langle f_{n_m}, T'y' \rangle\} = \{\langle Tf_{n_m}, y' \rangle\}$ is Cauchy. Thus by weak completeness of Y , $\{Tf_{n_m}\}$ converges weakly in Y , proving T is weakly compact.

The reduction to the case X, K , are separable is standard; see for example [2].

REFERENCES

1. C. A. Akemann, P. G. Dodds, and J. L. B. Gamlen, *Weak compactness in the dual space of a C^* -algebra*, J. Functional Analysis **10** (1972), 446–451.
2. J. Batt and E. J. Berg, *Linear bounded transformations on the space of continuous functions*, J. Functional Analysis **4** (1969), 215–239. MR **40** #1798.
3. C. Bessaga and A. Pełczyński, *On bases and unconditional convergence of a series in a Banach space*, Studia Math. **17** (1958), 151–164. MR **22** #5872.
4. ———, *A generalization of results of R. C. James concerning absolute bases in Banach spaces*, Studia Math. **17** (1958), 165–174. MR **22** #5874.
5. A. Grothendieck, *Sur les applications linéaires faiblement compactes d'espaces du type $C(K)$* , Canad. J. Math. **5** (1953), 129–173. MR **15**, 438.

6. R. C. James, *A non-reflexive space isometric with its second conjugate space*, Proc. Nat. Acad. Sci. U.S.A. **37** (1951), pp. 174–177. MR **13**, 356.

7. A. Pełczyński, *Projections in certain Banach spaces*, Studia. Math. **19** (1960), 209–228. MR **23** #A3441.

8. ———, *Banach spaces on which every unconditionally converging operator is weakly compact*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **10** (1962), 641–648. MR **26** #6785.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CONNECTICUT
06520

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA,
CANADA