

COHESIVE SETS: COUNTABLE AND UNCOUNTABLE

RICHARD A. SHORE¹

ABSTRACT. We show that many uncountable admissible ordinals (including some cardinals) as well as all countable admissible ordinals have cohesive subsets. Exactly which cardinals have cohesive subsets, however, is shown to depend on set-theoretic assumptions such as $V=L$ or a large cardinal axiom.

The study of recursion theory on the ordinals was initiated by Takeuti and then generalized by several others to all admissible ordinals. The analogy with ordinary recursion theory has been quite striking and particularly so for the theory of degrees of unsolvability. Many major theorems, especially ones about recursively enumerable degrees have been successfully generalized to all admissible ordinals. Thus for example, Sacks and Simpson [3] have introduced the finite injury priority argument into α -recursion theory to construct two incomparable α -r.e. degrees. Indeed even an infinite injury argument has been successfully adapted to this general setting to prove that the α -r.e. degrees are dense for every admissible α [5]. In general it seems fair to say that although the proofs are often somewhat different and usually more complicated than those in ordinary recursion theory, the theorems about degrees (at least the r.e. ones) seem to carry over.

The situation changes drastically when one turns from degrees to sets even if one restricts one's attention to the recursively enumerable sets. Of course the phenomena of nonregularity poses many interesting problems along these lines [4], [7] but a more striking example is provided by the notion of maximal set. (An α -r.e. set is maximal if and only if its complement is α -infinite but cannot be split into two α -infinite pieces by an α -r.e. set.) It is a well-known theorem of Friedberg that such sets exist in ordinary recursion theory [2]. On the other hand, Lerman and Simpson [1] have shown that there are no maximal r.e. sets for any uncountable admissible.

Finally, if we drop the requirement of recursive enumerability, the situation becomes even further removed from that of ordinary recursion

Received by the editors December 11, 1972.

AMS (MOS) subject classifications (1970). Primary 02F27; Secondary 02K05.

Key words and phrases. Admissible ordinal, cohesive set.

¹ During the preparation of this paper the author was partially supported by NSF Grant GP-28700.

© American Mathematical Society 1974

theory. Thus, Lerman and Simpson [1] ask if there are, for example, any uncountable cardinals α with cohesive subsets. (An α -infinite set is cohesive if and only if it cannot be split into two α -infinite pieces by any α -r.e. set.) We here answer this question affirmatively. However, their general question as to which admissibles have cohesive subsets turns out to be more complicated. The methods of [1] show, for example, that if $V=L$ then \aleph_1 (like each other successor cardinal) has no cohesive set. On the other hand, we will see that a large cardinal assumption implies that ω_1^L as well as \aleph_1 (indeed all true cardinals) have cohesive sets. Thus we have an example of a recursion theoretic question about sets which in general is not even absolute. Appropriately enough our methods of proof are (with the addition of definability considerations) much like those associated with Erdős' style partition relations and other large cardinal arguments.

We now turn to the theorems.

THEOREM 1. *If L_α is Σ_3 -admissible, cofinal with ω and satisfies the power set axiom, then there is a cohesive subset of α .*

PROOF. We inductively build a tree $(T, <_T)$ whose nodes x are members of α and have associated with them sets $E_x \subseteq \alpha$, consisting of all successors of x in T . To begin, we let 0 be the unique node of rank 0 and set $E_0 = \alpha - \{0\}$. At each level $\beta < \alpha$ we consider every node x of rank β and split E_x into two pieces, $E_x \cap R_\beta$ and $E_x - R_\beta$. (R_β is the β th α -r.e. set.) We then appoint the least elements, y_1 and y_2 , of these sets (if they are nonempty) as the immediate successors of x (of rank $\beta + 1$). We also set $E_{y_1} = (E_x \cap R_\beta) - \{y_1\}$ and $E_{y_2} = (E_x - R_\beta) - \{y_2\}$. At limit levels λ we naturally take all chains $\{x_i\}_{i < \lambda}$ in L_α and set the least element y of $\bigcap_{i < \lambda} E_{x_i}$ (if there is one) as the successor (of rank λ) of $\{x_i\}_{i < \lambda}$ and put $E_y = \bigcap_{i < \lambda} E_{x_i} - \{y\}$. We let T_β^α denote the successors of x of rank β .

The crucial fact about this tree is that every node x such that E_x is unbounded in α has successors y at every level of the tree less than α such that E_y is also unbounded in α . Of course every $x \in \alpha$ is on this tree. Next we note that T_β^α is the image of $(2^\beta)^{L_\alpha}$ under a partial Σ_2 function and so, by our assumptions, a member of L_α . We then note that the partial map defined on T_β^α by sending z to $\bigcup E_z$ if $\bigcup E_z < \alpha$ is Σ_3 . Thus if for some $\beta < \alpha$ there were no elements of T_β^α as required we could map T_β^α unboundedly into α by a Σ_3 function, contradicting the Σ_3 admissibility of α . To verify these assertions, first define $F: (2^\beta)^{L_\alpha} \rightarrow T_\beta^\alpha$ by $F(a) = y$ if and only if $y \in T_\beta^\alpha$ and $(\forall \delta < \beta)(y \in R_\delta \leftrightarrow \delta \in a)$. It now suffices to show that $y \in T_\beta^\alpha$ is a Σ_2 predicate. Say rank $x = \gamma$ and note that $y \in T_\beta^\alpha \equiv x <_T y$ and rank $y = \beta \equiv x < y \& (\forall \delta < \gamma)(x \in R_\delta \leftrightarrow y \in R_\delta) \& (\exists f: \beta \rightarrow \gamma)(f \text{ is one-one$

and order preserving and $(\forall \eta < \beta)(\forall \delta < \eta)(f(\eta) \in R_\delta \leftrightarrow y \in R_\delta)) \& \sim (\exists f : \beta + 1 \rightarrow y)(f \text{ is one-one and order preserving and } (\forall \eta < \beta + 1)(\forall \delta < \eta)(f(\eta) \in R_\delta \leftrightarrow y \in R_\delta))$. Finally, the function taking y of rank β to $\bigcup E_y$ is the Σ_3 -uniformization of the Π_2 -relation $S(y, z)$ given by $(\forall w)(y \leq_T w \rightarrow w \in z) \equiv (\forall w)(y \in w \& (\forall \delta < \beta)(w \in R_\delta \leftrightarrow y \in R_\delta) \rightarrow w \in z)$.

Armed with this fact we can now build a path of length α through T in ω many steps. Let $\{\beta_i\}_{i < \omega}$ be cofinal in α . Begin with $x_0 = 0$ and so of course E_{x_0} is unbounded in α . We can now successively choose x_i of rank β_i such that E_{x_i} is unbounded in α . Clearly, $\{x_i\}_{i < \omega}$ traces out a path P of length α in T . Moreover, $P = \{x \mid (\exists i < \omega)(x <_T x_i)\}$ is our desired cohesive set. For if we consider any α -r.e. set R_γ we see that either all elements of P of rank $> \gamma$ fall into R_γ or out of R_γ . As the elements of P of rank $\leq \gamma$ form a bounded Σ_2 set, it is α -finite, and so therefore are its intersections with R_γ and the complement of R_γ . \square

To see that this theorem answers the question of [1] we shall show that there are Σ_3 admissible cardinals cofinal with ω . We begin with a definable Σ_3 -skolem function F for L . Let $A_0 = \emptyset$, $A_{i+1} = (\bigcup (F''[A_i] \cap \text{ORD}))^+$ and $A = \bigcup_{i < \omega} A_i$. A is clearly a cardinal with the desired properties.

Turning now to the more general question of which admissibles have cohesive subsets we note first that we can adapt the above proof to establish

THEOREM 2. *Every countable admissible α has a cohesive subset.*

PROOF. Just arrange the α -r.e. sets in an ω -sequence R_i and build a tree as above. We can then of course get a path $\{y_i\}_{i < \omega}$ in T with E_{y_i} unbounded in α just by following a path determined by this requirement. We now form our cohesive set $\{x_i \mid i < \omega\}$ by choosing from each E_{y_i} an element x_i greater than β_i where $\{\beta_i\}_{i < \omega}$ is unbounded in α . This set is clearly cohesive. \square

We next note that, of course, if α is weakly compact in L the tree of Theorem 1 has an α -path and so has a cohesive subset. (This fact was already mentioned in [1].) Now the existence of a Ramsey cardinal, for example, guarantees that every true cardinal is weakly compact in L [6]. Since ω_1^L is then of course countable we see that a large cardinal assumption implies that ω_1^L as well as every real cardinal has a cohesive subset. On the other hand, the methods of [1] show that if \aleph_1 is a successor cardinal of L it has no cohesive subset. Thus we have our independence result. (Of course, no large cardinals are needed just to get a model with ω_1^L countable.)

In [1], Lerman and Simpson also indicated that if $V=L$, no cardinal α with Σ_2 -cofinality less than α has a cohesive subset. Our final theorem shows that this is best possible in the sense that there are (provably in

ZF) cardinals of L which are Σ_2 - but not Σ_3 -regular which have cohesive subsets. It also shows that there are real and constructible cardinals of all cofinalities with cohesive subsets.

THEOREM 3. *If α is a cardinal of L and for some $\beta > \alpha$, L_α is an elementary substructure of L_β with respect to Σ_2 formulas ($L_\alpha <_2 L_\beta$), then α has a cohesive subset and indeed a constructible one.*

PROOF. Let γ be the cofinality (in L) of α and let $\{\beta_i\}_{i < \gamma}$ be an unbounded increasing sequence in α . We build our cohesive set $\{x_i\}_{i < \gamma}$ by induction following the path traced by α itself. Say we have $\{x_i\}_{i < \lambda}$. Let $\delta = \max\{\bigcup_{i < \lambda} x_i, \beta_\lambda\}$. Let $f: \delta \rightarrow 2$ be given by $f(\eta) = 0$ if and only if $L_\beta \models \varphi_\eta(\alpha)$ where φ_η is the formula defining R_η over L_α . As α is a cardinal of L , $f \in L_\alpha$. Moreover, since $L_\alpha <_2 L_\beta$ we see that

$$L_\alpha \models (\exists x > \delta)(\forall \eta < \delta)(\varphi_\eta(x) \leftrightarrow f(\eta) = 0).$$

Let x_λ be such an x . $\{x_i | i < \gamma\}$ is now easily seen to be a cohesive subset of α . \square

To see that this theorem does indeed supply the desired example, consider a Σ_2 -Skolem function F for L . As before we set $A_0 = \emptyset$, $A_{\beta+1} = \bigcup (F''[A_\beta] \cap \text{ORD})^+$ and $A_\lambda = \bigcup_{\beta < \lambda} A_\beta$. If this is carried out in L (i.e. successor cardinal in the sense of L is intended) then for each limit λ , A_λ is Σ_2 -admissible but not Σ_3 -admissible and has a cohesive subset by the theorem ($L_{A_\lambda} <_2 L_{A_{\lambda+\omega}}$). On the other hand if we do the construction in the real world we get A_λ to be a true cardinal with cofinality that of λ . Of course the theorem still assures us that A_λ has a cohesive subset.

BIBLIOGRAPHY

1. M. Lerman and S. G. Simpson, *Maximal sets in α -recursion theory*, Israel J. Math. **4** (1973), 236–247.
2. H. Rogers, Jr., *Theory of recursive functions and effective computability*, McGraw-Hill, New York, 1967. MR **37** #61.
3. G. E. Sacks and S. G. Simpson, *The α -finite injury method*, Ann. Math. Logic **4** (1972), 343–367.
4. R. A. Shore, *On splitting an α -recursively enumerable set* (to appear).
5. ———, *The α -recursively enumerable degrees are dense* (in preparation).
6. J. H. Silver, *Some applications of model theory in set theory*, Ann. Math. Logic **3** (1971), 45–110.
7. S. G. Simpson, *Admissible ordinals and recursion theory*, Ph.D. Thesis, M.I.T., 1971.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637