

ENTIRENESS OF THE ENDOMORPHISM RINGS OF ONE-DIMENSIONAL FORMAL GROUPS¹

JONATHAN LUBIN

ABSTRACT. If, for a one-dimensional formal group of height h which is defined over the integers in a local field of characteristic zero, all the coefficients in degree less than p^h lie in an unramified extension of the p -adic numbers, then the endomorphism ring of the formal group is integrally closed.

In this note, \mathbb{Q}_p and \mathbb{Z}_p denote the field of p -adic numbers and the ring of p -adic integers, respectively; K and B denote a finite field extension of \mathbb{Q}_p and the integral closure of \mathbb{Z}_p in K , respectively; \bar{K} will be a fixed algebraic closure of K , and v the unique extension to \bar{K} of the (additive) p -adic valuation on \mathbb{Q}_p , normalized so that $v(p) = 1$; and finally, M will be the maximal ideal in the integral closure of B in \bar{K} , or, equivalently, the set of all elements z of \bar{K} for which $v(z) > 0$. All formal groups considered will be commutative and one dimensional.

In [3] the following proposition appeared:

Theorem 3.3.1. *If F is a one-dimensional formal group defined over B , of height $h < \infty$, and if the coefficients of F in terms of total degree less than p^h all lie in an unramified extension of \mathbb{Q}_p , then $\text{End}_B(F)$ is integrally closed in its fraction-field.*

It was soon pointed out to me by A. Frölich and A. Trojan that the proof in [3] was incorrect. Later [4] I proved the weaker result that if F itself is defined over an unramified extension of \mathbb{Q}_p , then $\text{End}_B(F)$ is integrally closed. That proof made essential use of the fact [4, Theorem 1.5] that if F , G , and H are formal groups defined over B , with $f \in \text{Hom}_B(F, G)$ and $g \in \text{Hom}_B(F, H)$ such that $\ker(f) \subset \ker(g)$, then there is some $h \in \text{Hom}_B(G, H)$ for which $h \circ f = g$. In this note I will use that fact together with the theory of the Newton polygon of a power series, as described for instance in [2], to show that Theorem 3.3.1 of [3] is correct as stated there.

The proofs below are for the category of formal groups, i.e., formal \mathbb{Z}_p -modules; the generalization, in the spirit of [5] or [1], to the category of

Received by the editors August 22, 1974.

AMS (MOS) subject classifications (1970). Primary 14L05; Secondary 14L20, 14G20.

¹Research supported by NSF Grant GP-29082.

formal A -modules, for A the ring of integers in a finite field extension of \mathbb{Q}_p , is a comparatively easy exercise.

Now let F be a fixed one-dimensional formal group defined over B , and of finite height h , with the property that all its coefficients in terms of total degree less than p^h lie in an unramified extension of \mathbb{Q}_p . Then according to Lemma 3.2.2 of [3], F is B -isomorphic to a formal group which is linear modulo degree p^h . We may assume from now on that F itself has this shape. It is a consequence of this, since B is of characteristic zero, that any endomorphism of F is also linear modulo degree p^h .

The points of finite order of F , in M , form a group W which is the disjoint union of $\{0\}$ with all the sets $X_m = \ker([p^m]_F) - \ker([p^{m-1}]_F)$, $m \geq 1$. We can now use the theory of the Newton polygon to show that if $w \in X_m$, then $v(w) = (p^h - 1)^{-1} p^{h(1-m)}$. Indeed, since $[p]_F(x) \equiv px + ux^{p^h} \pmod{(x^{p^{h+1}})}$, for some unit u of B , the Newton polygon of $[p]_F(x)$ has its first vertex at $(1, 1)$ and its next vertex at $(p^h, 0)$, so that the nonzero roots w of $[p]_F$ in M have $v(w) = 1/(p^h - 1)$. Inductively, if all elements y of X_{m-1} have $v(y) = (p^h - 1)^{-1} p^{h(2-m)}$, we use the fact that any w in X_m is a root of $-y + [p]_F(x)$ for some such y ; the Newton polygon of this power series has no vertices between $(0, v(y))$ and $(p^h, 0)$. The slope of this segment of the polygon is $-v(y)/p^h$, and this is the only segment of the polygon with negative slope. Thus $v(w) = v(y)/p^h$, completing the induction.

Now let f be a B -endomorphism of F . It will turn out that if f is not an automorphism, there is some $g \in \text{End}_B(F)$ such that $f = [p]_F \circ g$; in other words, $\text{End}_B(F)$ is a discrete valuation ring with prime element $[p]_F$, and hence certainly integrally closed in its fraction field.

Suppose now that the B -endomorphism f of F is not an automorphism, and not zero. Then $\ker(f) \neq \{0\}$, and in fact $\{0\} \neq \ker([p]_F) \cap \ker(f)$, since all elements of $\ker(f)$ are annihilated by some power of p . The fact that f has some nonzero roots in M implies that the first segment of the Newton polygon is not horizontal; the fact that f is linear modulo (x^{p^h}) implies that the right-hand endpoint of this segment has abscissa at least p^h . Nonzero elements of W with greatest v -value are just the elements of X_1 . So since f has some roots in X_1 , and at least $p^h - 1$ nonzero roots of greatest v -value, it follows that f has $p^h - 1$ roots in X_1 ; we have $\ker([p]_F) \subset \ker(f)$. This completes the proof.

REFERENCES

1. Lawrence Cox, *Formal A-modules*, Bull. Amer. Math. Soc. 79 (1973), 690–694. MR 48 #3973.
2. M. Lazard, *Les zéros des fonctions analytiques d'une variable sur un corps valué complet*, Inst. Hautes Études Sci. Publ. Math. No. 14 (1962), 47–75. MR 27 #2497.
3. Jonathan Lubin, *One-parameter formal Lie groups over p-adic integer rings*,

Ann. of Math. (2) 80 (1964), 464–484; Correction, *ibid.* (2) 84 (1966), 372. MR 29 #5827; 34 #179.

4. ———, *Finite subgroups and isogenies of one-parameter formal Lie groups*, *Ann. of Math.* (2) 85 (1967), 296–302. MR 35 #189.

5. ———, *Formal A -modules defined over A* , *Symposia Mathematica*, vol. III (INDAM, Rome, 1968/69), Academic Press, London, 1970, pp. 241–245. MR 42 #260.

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RHODE ISLAND 02912